

2. Diagnosis and Classification of Diabetes: Standards of Care in Diabetes—2026

Diabetes Care 2026;49(Suppl. 1):S27–S49 | <https://doi.org/10.2337/dc26-S002>

The American Diabetes Association (ADA) “Standards of Care in Diabetes” includes the ADA’s current clinical practice recommendations and is intended to provide the components of diabetes care, general treatment goals and guidelines, and tools to evaluate quality of care. Members of the ADA Professional Practice Committee for Diabetes, an interprofessional expert committee, are responsible for updating the Standards of Care annually, or more frequently as warranted. For a detailed description of ADA standards, statements, and reports, as well as the evidence-grading system for ADA’s clinical practice recommendations and a full list of Professional Practice Committee members, please refer to Introduction and Methodology. Readers who wish to comment on the Standards of Care are invited to do so at professional.diabetes.org/SOC.

Diabetes mellitus is a group of metabolic disorders of carbohydrate metabolism in which glucose is both underutilized as an energy source and overproduced due to inappropriate gluconeogenesis and glycogenolysis, resulting in hyperglycemia (1). Diabetes can be diagnosed by demonstrating increased concentrations of glucose in venous plasma or increased A1C in the blood. Diabetes is classified conventionally into several clinical categories (e.g., type 1 or type 2 diabetes, gestational diabetes mellitus, and other specific types derived from other causes, such as monogenic diabetes, exocrine pancreatic disorders, and high-risk medications) (2).

DIAGNOSTIC TESTS FOR DIABETES

Diabetes may be diagnosed based on A1C or plasma glucose criteria. Plasma glucose criteria include either the fasting plasma glucose (FPG), 2-h plasma glucose (2-h PG) during a 75-g oral glucose tolerance test (OGTT), or random glucose accompanied by classic symptoms of hyperglycemia (e.g., polyuria, polydipsia, and unexplained weight loss) or hyperglycemic crises (i.e., diabetic ketoacidosis [DKA] and/or hyperglycemic hyperosmolar state [HHS]) (Table 2.1).

Recommendations

- 2.1a Diagnose diabetes based on A1C or plasma glucose criteria. Plasma glucose criteria include either fasting plasma glucose (FPG), 2-h plasma glucose (2-h PG) during a 75-g oral glucose tolerance test (OGTT), or random glucose accompanied by classic hyperglycemic symptoms or crises (Table 2.1). B - 2.1b In the absence of unequivocal hyperglycemia (e.g., hyperglycemic crises), diagnosis requires confirmatory testing (Table 2.1). B

Duality of interest information for each contributor is available at <https://doi.org/10.2337/dc26-SDIS>.

Suggested citation: American Diabetes Association Professional Practice Committee for Diabetes. 2. Diagnosis and classification of diabetes: Standards of Care in Diabetes—2026. Diabetes Care 2026;49(Suppl. 1):S27–S49

© 2025 by the American Diabetes Association. Readers may use this work for educational, noncommercial purposes if properly cited and unaltered. The version of record may be linked at <https://diabetesjournals.org/care>, but ADA permission is required to post this work on any third-party site or platform. This publication and its contents may not be reproduced, distributed, or used for text or data mining, machine learning, or similar technologies without prior written permission. Requests to reuse, adapt, or distribute this work may be sent to permissions@diabetes.org. More information is available at <https://diabetesjournals.org/journals/pages/license>.

2026 February 01 on guest by

http://diabetesjournals.org/care/article-pdf/49/Supplement_1/S27/848854/dc26s002.pdf

Diagnosis and Classification of Diabetes

Diabetes Care Volume 49, Supplement 1, January 2026

Table 2.1—Criteria for the diagnosis of diabetes in nonpregnant individuals

A1C $\geq 6.5\%$ (≥ 48 mmol/mol). The test should be performed in a laboratory using a method that is NGSP certified and standardized to the DCCT assay.* OR FPG ≥ 126 mg/dL (≥ 7.0 mmol/L). Fasting is defined as no caloric intake for at least 8 h.* OR 2-h PG ≥ 200 mg/dL (≥ 11.1 mmol/L) during OGTT. The test should be performed as described by the WHO, using a glucose load containing the equivalent of 75 g anhydrous glucose dissolved in water.* OR In an individual with classic symptoms of hyperglycemia or hyperglycemic crisis, a random plasma glucose ≥ 200 mg/dL (≥ 11.1 mmol/L). Random is any time of the day without regard to time since previous meal.

DCCT, Diabetes Control and Complications Trial; FPG, fasting plasma glucose; OGTT, oral glucose tolerance test; NGSP, National Glycohemoglobin Standardization Program; WHO, World Health Organization; 2-h PG, 2-h plasma glucose. *In the absence of unequivocal hyperglycemia, diagnosis requires two abnormal results from different tests, which may be obtained at the same time (e.g., A1C and FPG), or the same test at two different time points.

Use of Fasting Plasma Glucose or 2-Hour Plasma Glucose for Screening and Diagnosis of Diabetes

In the clinical scenario where a person has classic symptoms of hyperglycemia (e.g., polyuria, polydipsia, unexplained weight loss) or presents with hyperglycemic crisis, measurement of random plasma glucose is sufficient to diagnose diabetes (symptoms of hyperglycemia or hyperglycemic crisis plus random plasma glucose ≥ 200 mg/dL [≥ 11.1 mmol/L]). In these cases, knowledge of the plasma glucose level is critical because, in addition to confirming that symptoms are due to diabetes, it will inform management decisions. In this context, testing A1C helps determine the chronicity of hyperglycemia. However, in an individual without symptoms, FPG or 2-h PG can be used for screening and diagnosis of diabetes. In nonpregnant individuals, FPG (or A1C) is typically preferred for routine screening due to the ease of administration (Table 2.3); however, the 2-h PG (OGTT) testing protocol is significantly more sensitive than the other two tests and is preferentially recommended for screening for some conditions (e.g., cystic fibrosis–related diabetes or post-transplantation diabetes mellitus). In the absence of classic symptoms of hyperglycemia, repeat testing is required to confirm the diagnosis regardless of the test used (see CONFIRMING THE DIAGNOSIS, below).

Major advantages of glucose monitoring are its low cost and availability. Disadvantages include the high diurnal variation in glucose and the 8-h fasting requirements (Table 2.3). Recent physical activity, illness, or acute stress can affect glucose concentrations. Glycolysis is also an important and underrecognized concern with glucose testing. Glucose concentrations will be falsely low if samples are not processed and analyzed promptly.

People should follow a mixed eating pattern with at least 150 g of carbohydrates on the 3 days prior to OGTT (12–14). Antecedent carbohydrate restriction in the days prior to OGTT can falsely elevate postchallenge glucose levels, potentially resulting in a false-positive OGTT (12).

Table 2.2—Criteria defining prediabetes in nonpregnant individuals

A1C 5.7–6.4% (39–47 mmol/mol) OR FPG 100 mg/dL (5.6 mmol/L) to 125 mg/dL (6.9 mmol/L) (IFG) OR 2-h PG during 75-g OGTT 140 mg/dL (7.8 mmol/L) to 199 mg/dL (11.0 mmol/L) (IGT)

For all three tests, risk is continuous, extending below the lower limit of the range and becoming disproportionately greater at the higher end of the range. FPG, fasting plasma glucose; IFG, impaired fasting glucose; IGT, impaired glucose tolerance; OGTT, oral glucose tolerance test; 2-h PG, 2-h plasma glucose.

Recommendations

2.2a The A1C test should be performed using a method that is certified by the National Glycohemoglobin.

2026 February 01 on guest by

http://diabetesjournals.org/care/article-pdf/49/Supplement_1/S27/848854/dc26s002.pdf from Downloaded

diabetesjournals.org/care

Diagnosis and Classification of Diabetes

Table 2.3—Considerations related to the use and interpretation of laboratory measurements of glucose and A1C

|| Glucose | A1C || ----- |

----- | | Cost | Inexpensive and available in most laboratories across the world | More expensive than glucose and not as widely available globally | | Time frame of hyperglycemia | Acute measure | Chronic measure of glucose exposure over the past ~2–3 months | | Preanalytic stability | Poor; plasma must be separated immediately or samples must be kept on ice to prevent glycolysis | Good | | Sample | Measurement can vary depending on sample type (plasma, serum, whole blood) and source (capillary, venous, arterial) | Requires whole-blood sample | | Assay standardization | Not standardized | Well standardized | | Fasting | Fasting or timed samples required | Nonfasting test; no participant preparation is needed | | Within-person variability | High | Low | | Acute factors that can affect levels | Food intake, stress, recent illness, activity | Unaffected by recent food intake, stress, illness, activity | | Other individual factors that can affect test results | Diurnal variation, medications, alcohol, smoking, bilirubin | Altered erythrocyte turnover (e.g., anemia, iron status, splenectomy, blood loss, transfusion, hemolysis, glucose-6-phosphate dehydrogenase deficiency, erythropoietin), HIV, cirrhosis, kidney failure, dialysis, pregnancy | | Test interferences | Depends on specific assay: sample handling/processing time, hemolysis, severe hypertriglyceridemia, severe hyperbilirubinemia | Depends on specific assay: hemoglobin variants, severe hypertriglyceridemia, severe hyperbilirubinemia |

Data are from Selvin (236).

2.2b

Point-of-care A1C testing for diabetes screening and diagnosis should be restricted to devices approved for diagnosis by the U.S. Food and Drug Administration at Clinical Laboratory Improvement Amendments–certified laboratories that perform testing of moderate complexity or higher by trained personnel. B

2.3

Evaluate for the possibility of a problem or interference with either test when there is consistent and substantial discordance between blood glucose values and A1C test results. B

2.4

In conditions associated with an altered relationship between A1C and glycemia, such as some hemoglobin variants, pregnancy, glucose-6-phosphate dehydrogenase deficiency, HIV, and conditions that may alter red blood cell turnover, plasma glucose criteria should be used to diagnose diabetes. B

The A1C test should be performed using a method that is certified by the National Glycohemoglobin Standardization Program (NGSP) (ngsp.org) and standardized or traceable to the Diabetes Control and Complications Trial (DCCT) reference assay. Outside the U.S., some assays are NGSP certified but many more are International Federation of Clinical Chemistry (IFCC) certified (a similarly stringent process) (1).

Point-of-care A1C assays may be NGSP certified and cleared by the U.S. Food and Drug Administration (FDA) for use in monitoring glycemic management in people with diabetes in both Clinical Laboratory Improvement Amendments (CLIA)–regulated and CLIA-waived settings. FDA-approved point-of-care A1C testing can be used in laboratories or sites that are CLIA certified and inspected and meet the CLIA quality standards. These standards include specified personnel requirements (including documented annual competency assessments) and participation three times per year in an approved proficiency testing program (1,15,16).

A1C has several advantages compared with FPG and OGTT, including greater convenience (fasting is not required), greater preanalytical stability, and fewer day-to-day perturbations during stress, changes in nutrition, or illness. However, it should be noted that there is lower sensitivity of A1C at the designated cut point compared with that of 2-h PG as well as limited access in some parts of the world (Table 2.3).

A1C reflects glucose bound to hemoglobin over the life span of the erythrocyte (~120 days) and is thus a “weighted” average that is more heavily affected by recent blood glucose exposure. Thus, clinically meaningful changes in A1C can be seen in ~120 days. A1C is an indirect measure of glucose exposure, and factors that affect hemoglobin concentrations or erythrocyte turnover can affect A1C (e.g., thalassemia or folate deficiency) (Table 2.3). A1C may not be a suitable diagnostic test in people with anemia, people treated with

erythropoietin, or people undergoing hemodialysis or HIV treatment (1,17). Some A1C assays can be affected by hemoglobin variants. For instance, an A1C assay without interference from hemoglobin variants.

2026 February 01 on guest by

http://diabetesjournals.org/care/article-pdf/49/Supplement_1/S27/848854/dc26s002.pdf from Downloaded

Diagnosis and Classification of Diabetes Diabetes Care Volume 49, Supplement 1, January 2026

Hemoglobin variants should be used in individuals with sickle cell trait. An updated list of A1C assays with interferences is available at ngsp.org/interf.asp. Another genetic variant, X-linked glucose-6-phosphate dehydrogenase G202A, carried by 11% of Black individuals in the U.S., is associated with a decrease in A1C of about 0.8% in homozygous men and 0.7% in homozygous women compared with levels in individuals without the variant (18). There is controversy regarding racial differences in A1C. Studies have found that Black individuals have slightly higher A1C levels (approximately 0.3%) than non-Hispanic White or Hispanic people (19–22). A more precise understanding of the genetic determinants of A1C in diverse populations (18) is the focus of ongoing investigations (23,24). While some genetic variants might be more common in certain race or ancestry groups, it is important that we do not use race or ancestry as proxies for poorly understood genetic differences. Reassuringly, studies have shown that the association of A1C with risk for complications appears to be similar in Black and non-Hispanic White populations (25).

Confirming the Diagnosis

In the absence of a clear clinical diagnosis (e.g., individual with classic symptoms of hyperglycemia or hyperglycemic crisis and random plasma glucose ≥ 200 mg/dL [≥ 11.1 mmol/L]), confirmatory tests are necessary to establish the diagnosis. This can be accomplished by two abnormal screening test results, measured either at the same time (26) or at two different time points. If using samples at two different time points, it is recommended that the second test, which may be either a repeat of the initial test or a different test, be performed in a timely manner. For example, if the A1C is 7.0% (53 mmol/mol) and a repeat result is 6.8% (51 mmol/mol), the diagnosis of diabetes is confirmed. Two different tests (such as A1C and FPG) both having results above the diagnostic threshold when collected at the same time or at two different time points would also confirm the diagnosis. On the other hand, if an individual has discordant results from two different tests, then the test result that is above the diagnostic cut point should be repeated, with careful consideration of factors that may affect measured A1C or glucose levels. The diagnosis is made based on the confirmatory screening test. For example, if an individual meets the diagnostic criterion for A1C (two results $\geq 6.5\%$ [≥ 48 mmol/mol]) but not FPG (≥ 126 mg/dL [≥ 7.0 mmol/L]), that person should nevertheless be considered to have diabetes.

Classification

Diabetes is classified conventionally into several clinical categories, although these are being reconsidered based on genetic, metabolomic, and other characteristics and pathophysiology (1):

1. Type 1 diabetes (due to autoimmune β -cell destruction, usually leading to absolute insulin deficiency, including latent autoimmune diabetes in adults)
2. Type 2 diabetes (due to a nonautoimmune progressive loss of adequate β -cell insulin secretion, frequently on the background of insulin resistance)
3. Specific types of diabetes due to other causes, e.g., monogenic diabetes syndromes, diseases of the exocrine pancreas, and drug- or chemical-induced diabetes
4. Gestational diabetes mellitus (diabetes diagnosed in the second or third trimester of pregnancy that was not clearly overt diabetes prior to gestation or other types of diabetes occurring throughout pregnancy, such as type 1 diabetes)

Occasionally, people with type 2 diabetes may present with DKA (33) in the context of ketosis-prone diabetes, particularly members of certain racial, ethnic, and ancestral groups (e.g., Black and Hispanic/Latino adults) (27). The initial requirement for insulin replacement therapy in individuals presenting with ketosis-prone diabetes is generally transient. It is important for health care professionals to realize that classification of diabetes type is not always straightforward at presentation and that misdiagnosis can occur in up to 40% of adults with new type 1 diabetes (e.g., adults with type 1 diabetes misdiagnosed as having type 2 diabetes). In comparison, individuals with maturity-onset diabetes of the young (MODY) may be misdiagnosed as having type 1 diabetes (32). Although difficulties in distinguishing diabetes type may occur in all age groups at onset, the diagnosis generally becomes more obvious in people with marked loss of β -cell function as they rapidly

require insulin therapy (Fig. 2.1). Although not prospectively validated, the AABCC approach is a useful tool to help distinguish.

2026 February 01 on guest by

http://diabetesjournals.org/care/article-pdf/49/Supplement_1/S27/848854/dc26s002.pdf from Downloaded

diabetesjournals.org/care

Diagnosis and Classification of Diabetes

Flowchart for investigation of suspected type 1 diabetes in newly diagnosed adults, based on data from White European populations

Adult with suspected type 1 diabetes1

| Test islet autoantibodies2 ||| |-----|
-----|-----| | Islet autoantibody positive | Islet autoantibody
negative ||| Type 1 diabetes | Age ||| <35 years | >35 years | | Are there features of monogenic diabetes?3 |
Unclear classification7 ||| Make clinical decision as to how person with diabetes should be treated ||| Trial
of noninsulin therapy may be appropriate8 ||| Yes | No ||| Test C-peptide4 | Are there features of type 2
diabetes?5 ||| >200 pmol/L | <200 pmol/L ||| No | <200 pmol/L ||| 200–600 pmol/L | >600 pmol/L ||| Genetic
testing for monogenic diabetes where available6 | Type 1 diabetes ||| Indeterminate9 ||| Type 2 diabetes |||

Figure 2.1—Flowchart for investigation of suspected type 1 diabetes in newly diagnosed adults, based on data from White European populations. 1No single clinical feature confirms type 1 diabetes in isolation. 2Glutamic acid decarboxylase (GAD) should be the primary antibody measured and, if negative, should be followed by islet tyrosine phosphatase 2 (IA-2) and/or zinc transporter 8 (ZnT8) where these tests are available. In individuals who have not been treated with insulin, antibodies against insulin may also be useful. In those diagnosed at <35 years of age who have no clinical features of type 2 diabetes or monogenic diabetes, a negative result does not change the diagnosis of type 1 diabetes, since 5–10% of people with type 1 diabetes do not have antibodies. 3Monogenic diabetes is suggested by the presence of one or more of the following features: A1C <58 mmol/mol (<7.5%) at diagnosis, one parent with diabetes, features of a specific monogenic cause (e.g., renal cysts, partial lipodystrophy, maternally inherited deafness, and severe insulin resistance in the absence of obesity), and monogenic diabetes prediction model probability >5% (diabetesgenes.org/exeter-diabetes-app/Modycalculator). 4A C-peptide test is only indicated in people receiving insulin treatment. A random sample (with concurrent glucose) within 5 h of eating can replace a formal C-peptide stimulation test in the context of classification. If the result is ≥600 pmol/L (≥1.8 ng/mL), the circumstances of testing do not matter. If the result is <600 pmol/L (<1.8 ng/mL) and the concurrent glucose is <4 mmol/L (<70 mg/dL) or the person may have been fasting, consider repeating the test. Results showing very low levels (e.g., <80 pmol/L [<0.24 ng/mL]) do not need to be repeated. Where a person is insulin treated, C-peptide must be measured prior to insulin discontinuation to exclude severe insulin deficiency. Do not test C-peptide within 2 weeks of a hyperglycemic emergency. 5Features of type 2 diabetes include increased BMI (≥25 kg/m²), absence of weight loss, absence of ketoacidosis, and less marked hyperglycemia. Less discriminatory features include non-White ethnicity, family history, longer duration and milder severity of symptoms prior to presentation, features of metabolic syndrome, and absence of a family history of autoimmunity. 6If genetic testing does not confirm monogenic diabetes, the classification is unclear and a clinical decision should be made about treatment. 7Type 2 diabetes should be strongly considered in older individuals. In some cases, investigation for pancreatic or other types of diabetes may be appropriate. 8A person with possible type 1 diabetes who is not treated with insulin will require careful monitoring and education so that insulin can be rapidly initiated in the event of glycemic deterioration. 9C-peptide values 200–600 pmol/L (0.6–1.8 ng/mL) are usually consistent with type 1 diabetes or maturity-onset diabetes of the young but may occur in insulin-treated type 2 diabetes, particularly in people with normal or low BMI or after long duration. Reprinted and adapted from Holt et al. (32).

2026 February 01 on guest by

http://diabetesjournals.org/care/article-pdf/49/Supplement_1/S27/848854/dc26s002.pdf from Downloaded

Diagnosis and Classification of Diabetes Diabetes Care Volume 49, Supplement 1, January 2026

In both type 1 and type 2 diabetes, genetic and environmental factors can result in the progressive loss of β -cell mass and/or function that manifests clinically as hyperglycemia. Once hyperglycemia occurs, people with all forms of diabetes are at risk for developing the same chronic complications, although rates of progression may differ. The identification of individualized therapies for diabetes in the future will be informed by better characterization of the many paths to β -cell demise or dysfunction (35). Ongoing research efforts aim to combine clinical, pathophysiological, and genetic characteristics to more precisely define the subsets of diabetes that are clustered into the current nomenclature with the goal of devising personalized treatment approaches (24).

Regarding the pathophysiology of type 1 diabetes, prospective studies showed that the persistent presence of two or more islet autoantibodies is a near-certain predictor of clinical diabetes (36). In at-risk cohorts followed from birth or a very young age, seroconversion rarely occurs before 6 months of age and there is a peak in seroconversion between 9 and 24 months of age (37–39). The rate of progression from preclinical to symptomatic stages of type 1 diabetes is dependent on the age at first detection of an autoantibody, number of autoantibodies, autoantibody specificity, and autoantibody titer. Glucose and A1C levels may rise well before the clinical onset of diabetes (e.g., changes in FPG and 2-h PG can occur about 6 months before diagnosis) (40), making diagnosis feasible under ideal situations of serial monitoring of individuals at high risk of type 1 diabetes before the onset of DKA. The current paradigm of the progression of type 1 diabetes across three distinct stages (Table 2.4) also serves as a framework for research and regulatory decision-making (35,41).

Adapted from Skyler et al. (35). FPG, fasting plasma glucose; IFG, impaired fasting glucose; IGT, impaired glucose tolerance; 2-h PG, 2-h plasma glucose. Alternative additional stage 2 diagnostic criteria of 30-, 60-, or 90-min plasma glucose on oral glucose tolerance test ≥ 200 mg/dL (≥ 11.1 mmol/L) and confirmatory testing in those aged ≥ 18 years have been used in clinical trials (82). Dysglycemia can be defined by one or more criteria as outlined in the table.

Recommendations

1. Screen for presymptomatic type 1 diabetes by testing autoantibodies against insulin (IA), glutamic acid decarboxylase (GAD), islet antigen 2 (IA-2), or zinc transporter 8 (ZnT8). B 2. Autoantibody-based screening for presymptomatic type 1 diabetes should be offered to those with a family history of type 1 diabetes or otherwise known elevated genetic risk. B 3. Individuals with screening results positive for one or more islet autoantibodies should be evaluated for stage 3 (overt) type 1 diabetes (using A1C, urinalysis, and/or plasma glucose), which would require prompt clinical management and education. B 4. Individuals with multiple confirmed islet autoantibodies and without overt type 1 diabetes have a high risk for progression to stage 3 type 1 diabetes and should be referred to a specialized center for metabolic staging (Table 2.4), education, and consideration of prevention trials or approved treatments (e.g., teplizumab). B 5. Individuals with a single confirmed IA-2 autoantibody should be monitored similarly to individuals with multiple islet autoantibodies, as IA-2 autoantibody positivity is an independent risk factor for progression. B 6. Individuals with a single confirmed islet autoantibody should undergo repeat antibody testing every 6 months to 3 years (depending on age) to assess for persistence or seroconversion. E 7. Standardized islet autoantibody tests are recommended for classification of diabetes in adults who have phenotypic risk factors that overlap with those for type 1 diabetes (e.g., younger age at diagnosis, unintentional weight loss, ketoacidosis, or short time to insulin treatment). E

Immune-Mediated Diabetes

Autoimmune type 1 diabetes accounts for 5–10% of diabetes and is caused by autoimmune destruction of the pancreatic β -cells. Autoimmune markers include islet cell autoantibodies and autoantibodies to glutamic acid decarboxylase (GAD) (e.g., GAD65), insulin, the tyrosine phosphatases islet antigen 2 (IA-2) and IA-2b, and zinc transporter 8 (ZnT8). Clinical studies are being conducted to test various methods of preventing or delaying type 1 diabetes in those with evidence of islet autoimmunity (trialnet.org/our-research/prevention-studies) (36–38,43,46,47).

The disease has strong HLA associations, with linkage to the DQB1 and DRB1 haplotypes, and genetic screening has been used in some research studies to identify high-risk populations. Specific alleles in these genes can be either predisposing (e.g., DRB1*0301-DQB1*0201 [DR3-DQ2] and DRB1*0401-DQB1*0302 [DR4-DQ8]) or protective (e.g., DRB1*1501 and DQA1*0102-DQB1*0602).

Evidence from pediatric cohorts shows that among children with a persistent single islet autoantibody, the 10-year cumulative risk of progression to clinical type 1 diabetes is ~14–15%, with much of the risk accruing in the first 2 years after seroconversion and the highest risk being associated with younger age and single IA-2 autoantibody positivity (36). In adults, progression risk is lower and decreases with age, and prospective data to define an optimal cadence are limited; therefore, intervals are extrapolated from pediatric patterns and clinical diabetes expertise.

The practical implication of this information is that the four islet autoantibodies (GAD autoantibody, IA-2 autoantibody, insulin autoantibody, ZnT8 autoantibody), a random venous or capillary blood glucose, and an A1C test should be repeated at intervals of 6 months to 3 years based on age. In children ≤ 3 years, test every 6 months for 3 years and then annually for 3 more years. In children and adolescents 3–18 years old, test annually and consider discontinuing after 3 years if no progression to multiple antibodies or onset of dysglycemia. In adults, test every 3 years or annually if additional risk factors are present, such as another autoimmune disease, elevated genetic risk score, or first-degree relative with type 1 diabetes (51).

Early diagnostic testing (Table 2.4 - stage 3 type 1 diabetes should be performed promptly in any individual with one or more islet autoantibodies to avoid delays in treatment, which can occur with long wait times for specialty referrals. Identifying overt hyperglycemia early can help prevent DKA at stage 3.

In adults with stage 3 type 1 diabetes, isolated GAD autoantibody positivity is common and has been associated with slower loss of endogenous insulin production after diagnosis (32). In early stages of type 1 diabetes, single GAD65 autoantibody positivity is also associated with slower progression to stage 3 compared with IA-2 or ZnT8A autoantibody positivity (1,53).

The rate of β -cell destruction is quite variable, being rapid in some individuals (particularly but not exclusively in infants and children) and slow in others (mainly but not exclusively adults) (40,54). Children and adolescents often present with DKA as the first manifestation of the disease, and rates in the U.S. have increased over the past 20 years (27–29).

2026 February 01 on guest by

http://diabetesjournals.org/care/article-pdf/49/Supplement_1/S27/848854/dc26s002.pdf from Downloaded

Diagnosis and Classification of Diabetes Diabetes Care Volume 49, Supplement 1, January 2026

that can rapidly change to severe hyperglycemia and/or DKA with infection or other stress. Adults may retain sufficient β -cell function to prevent DKA for many years; such individuals may have remission characterized by decreased insulin needs for months or years, but they eventually become insulin dependent and are at risk for DKA (30,31,55–57). At this later stage of the disease, there is little or no insulin secretion, as manifested by low or undetectable levels of plasma C-peptide. Autoimmune destruction of β -cells has multiple genetic factors and is also related to environmental factors that are still poorly defined. Although individuals do not classically have obesity when they present with type 1 diabetes, obesity is increasingly common in the general population; as such, obesity should not preclude testing for type 1 diabetes. People with type 1 diabetes are also prone to other autoimmune disorders, such as Hashimoto thyroiditis, Graves disease, celiac disease, Addison disease, vitiligo, autoimmune hepatitis, myasthenia gravis, and pernicious anemia (see section 4, “Comprehensive Medical Evaluation and Assessment of Comorbidities”).

Immune Checkpoint Inhibitor–Induced Autoimmune Diabetes

The introduction of immunotherapy, specifically immune checkpoint inhibitors (ICIs), for cancer treatment has led to unexpected adverse events, including immune system activation precipitating autoimmune disease. Fulminant onset of autoimmune diabetes can occur with DKA and low or undetectable levels of C-peptide as a marker of endogenous β -cell function (64,65). Fewer than half of these individuals have islet autoantibodies, and most do not experience a phase of partial remission, which is commonly seen in type 1 diabetes, supporting the emerging view of a partially distinct pathophysiology. A study reported the risk of progression to type 1 diabetes from the time of seroconversion to autoantibody positivity in three pediatric cohorts from Finland, Germany, and the U.S. Of the 585 children who developed more than two autoantibodies, nearly 70% developed type 1 diabetes within 10 years and 84% within 15 years (36). These findings are highly significant, because while the German group was recruited from offspring of parents with type 1 diabetes, the Finnish and American groups were recruited from the general population. Remarkably, the findings in all three groups were the same, suggesting that the same sequence of events led to clinical disease in both “sporadic” and familial cases of type 1 diabetes. Indeed, the risk of type 1 diabetes increases as the number of relevant autoantibodies detected increases (46,72,73).

In The Environmental Determinants of Diabetes in the Young (TEDDY) study, type 1 diabetes developed in 21% of 363 subjects with at least one autoantibody at 3 years of age (74). Such testing, coupled with education about diabetes symptoms and close follow-up, has been shown to enable earlier diagnosis and to prevent DKA (75,76). In several cohort studies, up to 50% of children with only a single autoantibody revert to being islet autoantibody negative during follow-up (77,78). Therefore, it is recommended that the first autoantibody-positive test be confirmed with a second test within 3 months, preferably in a laboratory that meets the performance standards set by the Islet Autoantibody Standardization Program (IASP) (51).

Idiopathic Type 1 Diabetes

Some forms of type 1 diabetes have no known etiologies. Individuals have permanent insulinopenia and are prone to DKA but have no evidence of β -cell autoimmunity. However, only a minority of people with type 1 diabetes fall into this category.

Screening for Type 1 Diabetes Risk

Type 1 diabetes genetic risk scores have been used in newborn screening to identify those at risk for future presentation of the disease. In a simulation...

2026 February 01 on guest by

http://diabetesjournals.org/care/article-pdf/49/Supplement_1/S27/848854/dc26s002.pdf Downloaded

diabetesjournals.org/care

using one such genetic risk score, the majority of those who would go on to develop type 1 diabetes, >77%, could be identified within just 10% of the general population, identifying a subset who may most benefit from

autoantibody testing (79). As many genetic risk studies have been performed in populations of European ancestry and discriminatory ability may differ in those of different ancestry, more large case-control cohorts from non-European populations are still needed (80).

Screening programs are available in Europe (e.g., Fr1da and gppad.org), Australia (e.g., type1screen.org), and the U.S. (e.g., trialnet.org, askhealth.org, and cascadekids.org). General population-based screening programs may offer broader testing where high-quality, validated assays and resources for appropriate follow-up of results are available, with several countries considering making such testing part of standard care. In 2023, Italy introduced nationwide screening for type 1 diabetes and celiac disease in the general population aged 1–17 years (81). Individuals who test autoantibody positive should be provided with or referred for counseling about the risk of developing diabetes, diabetes symptoms, and DKA prevention and should be.

PREDIABETES AND TYPE 2 DIABETES

Recommendations

1. Screening for risk of prediabetes and type 2 diabetes with an assessment of risk factors or validated risk calculator should be done in asymptomatic adults. B 2. Testing for prediabetes or type 2 diabetes in asymptomatic people should be considered in adults of any age with overweight or obesity who have one or more risk factors (Table 2.5). B 3. For all other people, screening should begin at age 35 years. B 4. In people without prediabetes or diabetes after screening, repeat screening recommended at a minimum of 3-year intervals is reasonable, sooner with symptoms or change in risk (e.g., weight gain). C 5. To screen for prediabetes and type 2 diabetes, FPG, 2-h PG during 75-g OGTT, and A1C are each appropriate (Table 2.1 and Table 2.2). B 6. When using OGTT as a screening tool for prediabetes or diabetes, adequate carbohydrate intake (at least 150 g/day) should be assured for 3 days prior to testing. C 7. Risk-based screening for prediabetes or type 2 diabetes should be considered after the onset of puberty or after 10 years of age, whichever occurs earlier, in children and adolescents with overweight (BMI $\geq$ 85th percentile) or obesity (BMI $\geq$ 95th percentile) and who have one or more risk factors for diabetes. (See Table 2.6 for evidence grading of risk factors.) B 8. Consider screening people for prediabetes or diabetes if they are on certain medications, such as statins, thiazide diuretics, and some HIV medications, as these agents are known to increase the risk of these conditions. C 9. In people who are prescribed second-generation antipsychotic medications, screen for prediabetes and diabetes at baseline and repeat 12–16 weeks after medication initiation or sooner, if clinically indicated, and annually thereafter. B 10. People with HIV should be screened for diabetes and prediabetes with an FPG test before starting antiretroviral therapy, at the time of switching antiretroviral therapy, and 3–6 months after starting or switching antiretroviral therapy. If initial screening results are normal, FPG should be checked annually. E

Table 2.5—Criteria for screening for diabetes or prediabetes in asymptomatic adults

- | 1. Testing should be considered in adults with overweight or obesity (BMI $\geq$ 25 kg/m² or $\geq$ 23 kg/m² in individuals of Asian ancestry) who have one or more of the following risk factors: | | -----

| -----
----- | | • First-degree relative with diabetes | • High-risk race, ethnicity, and ancestry (e.g., African American, Latino, Native American, Asian American) | | • History of cardiovascular disease | • Hypertension ($\geq$ 130/80 mmHg or on therapy for hypertension) | | • HDL cholesterol level $<$ 35 mg/dL ($<$ 0.9 mmol/L) and/or triglyceride level $>$ 250 mg/dL ($>$ 2.8 mmol/L) | • Individuals with polycystic ovary syndrome | | • Physical inactivity | • Other clinical conditions associated with insulin resistance (e.g., severe obesity, acanthosis nigricans, metabolic dysfunction-associated steatotic liver disease) |
2. People with prediabetes (A1C $\geq$ 5.7% [$\geq$ 39 mmol/mol], IGT, or IFG) should be tested yearly.
3. People who were diagnosed with GDM should have testing at least every 1–3 years.
4. For all other people, testing should begin at age 35 years.
5. If results are normal, testing should be repeated at a minimum of 3-year intervals, with consideration of more frequent testing depending on initial results and risk status.

6. Individuals in other high-risk groups (e.g., people with HIV, exposure to high-risk medicines, evidence of periodontal disease, history of pancreatitis) should also be closely monitored.

GDM, gestational diabetes mellitus; IFG, impaired fasting glucose; IGT, impaired glucose tolerance.

Prediabetes is the term used to identify glucose or A1C levels that do not meet the criteria for diabetes yet fall in an intermediate range between normoglycemia and diabetes (25,83). People with prediabetes are defined by the presence of IFG and/or IGT and/or A1C 5.7–6.4% (39–47 mmol/mol) (Table 2.2). Prediabetes is a significant risk factor for progression to diabetes as well as cardiovascular disease and several other cardiometabolic outcomes. Criteria for screening.

2026 February 01 on guest by

http://diabetesjournals.org/care/article-pdf/49/Supplement_1/S27/848854/dc26s002.pdf from Downloaded

Diagnosis and Classification of Diabetes # Diabetes Care Volume 49, Supplement 1, January 2026

Table 2.6—Risk-based screening for type 2 diabetes or prediabetes in asymptomatic children and adolescents in a clinical setting

Screening should be considered in youth* who have overweight (≥ 85 th percentile) or obesity (≥ 95 th percentile) A and who have one or more additional risk factors:

- Maternal history of diabetes or GDM during the child's gestation
- A - Family history of type 2 diabetes in first- or second-degree relative
- A - High-risk race, ethnicity, and ancestry
- A (see Table 2.5)
- Signs of insulin resistance or conditions associated with insulin resistance (acanthosis nigricans, hypertension, dyslipidemia, polycystic ovary syndrome, large- or small-for-gestational-age birth weight)
- B

GDM, gestational diabetes mellitus. *After the onset of puberty or after 10 years of age, whichever occurs earlier. If tests are normal, repeat testing at a minimum of 3-year intervals (or more frequently if BMI is increasing or risk factor profile is deteriorating) is recommended. Reports of type 2 diabetes before age 10 years exist, and this can be considered with numerous risk factors.

For details regarding individuals with prediabetes most likely to benefit from a formal behavioral or lifestyle intervention, see section 3, "Prevention or Delay of Diabetes and Associated Comorbidities."

Type 2 Diabetes

Type 2 diabetes accounts for 90–95% of all diabetes cases and encompasses individuals who generally have relative (rather than absolute) insulin deficiency in association with insulin resistance (i.e., decreased biological responses to insulin).

Prediabetes is associated with obesity (especially abdominal or visceral obesity), dyslipidemia with high triglycerides and/or low HDL cholesterol, and hypertension. The presence of prediabetes should prompt a comprehensive screening for cardiovascular risk factors.

Diagnosis of Prediabetes

IFG is defined as FPG levels from 100 to 125 mg/dL (from 5.6 to 6.9 mmol/L) (76,82) and IGT as 2-h PG levels during 75-g OGTT from 140 to 199 mg/dL (from 7.8 to 11.0 mmol/L) (10). It should be noted that the World Health Organization and several diabetes organizations define the IFG lower limit at 110 mg/dL (6.1 mmol/L). The ADA also initially endorsed this IFG lower limit in 1997 (10). However, in 2003 the ADA adopted the new range of 100–125 mg/dL (5.6–6.9 mmol/L) to better define IFG so that the population at risk for developing diabetes with IFG would be comparable to that with IGT (11).

As with the glucose measures, several prospective studies that used A1C to predict the progression to diabetes demonstrated a strong, continuous curvilinear association between A1C and risk of diabetes. In a systematic review of 44,203 individuals from 16 cohort studies with a follow-up interval averaging 5.6 years (range 2.8–12 years), those with A1C between 5.5% and 6.0% (between 37 and 42 mmol/mol) had a substantially increased risk of diabetes (5-year incidence from 9 to 25%). Those with an A1C range of 6.0–6.5% (42–48 mmol/mol) had a 5-year risk of developing diabetes between 25% and 50% and a relative risk 20 times higher than that with A1C of 5.0% (31 mmol/mol) (84).

In a community-based study of Black and non-Hispanic White adults without diabetes, baseline A1C was a stronger predictor of subsequent diabetes and cardiovascular events than fasting glucose (85). Another analysis also indicates that A1C at baseline was a strong predictor of the development of glucose-defined diabetes during the Diabetes Prevention Program (DPP) and its follow-up (7).

An A1C range of 5.7–6.4% (39–47 mmol/mol) identifies a group of individuals at high risk for diabetes and cardiovascular outcomes. These individuals should be informed of their increased risk for diabetes and cardiovascular disease and counseled about effective strategies to lower their risks (see section 3, “Prevention or Delay of Diabetes and Associated Comorbidities”).

Similar to glucose measurements, the continuum of risk is continuous and curvilinear: as A1C rises, the diabetes risk rises disproportionately (84). Aggressive interventions and vigilant follow-up should be pursued for those considered at very high risk (e.g., those with A1C >6.0% [>42 mmol/mol] and individuals with both IFG and IGT).

Table 2.5 outlines the criteria for screening for prediabetes. The ADA risk test is an additional option (i.e., an awareness tool for the layperson and the healthcare professional) to determine the appropriateness of screening for diabetes or prediabetes in asymptomatic adults (diabetes.org/diabetes-risk-test). For additional background regarding risk factors and screening.

2026 February 01 on guest by

http://diabetesjournals.org/care/article-pdf/49/Supplement_1/S27/848854/dc26s002.pdf from Downloaded

diabetesjournals.org/care # Diagnosis and Classification of Diabetes

developing macrovascular and microvascular complications (89). People with type 2 diabetes initially may have insulin levels that appear normal or elevated, yet the failure to normalize blood glucose reflects a relative defect in glucose-stimulated insulin secretion that is insufficient to compensate for insulin resistance. Insulin resistance may improve with weight reduction, physical activity, and/or pharmacologic treatment of hyperglycemia but is seldom restored to normal. Intensive lifestyle interventions and/or metabolic surgery can lead to diabetes remission (90–93) (see section 8, “Obesity and Weight Management for the Prevention and Treatment of Diabetes”).

The risk of developing type 2 diabetes increases with age, obesity, and lack of physical activity (94,95). It occurs more frequently in individuals with prediabetes, prior gestational diabetes mellitus, or polycystic ovary syndrome. It is also more common in people with hypertension or dyslipidemia and in certain racial, ethnic, and ancestral subgroups (Table 2.5). Finally, treatment with certain medications (e.g., glucocorticoids) can represent a significant risk factor for developing type 2 diabetes. In adults without traditional risk factors for type 2 diabetes and/or of younger age, consider islet autoantibody testing to exclude the diagnosis of type 1 diabetes (32) (Fig. 2.1).

Screening and Testing for Prediabetes and Type 2 Diabetes in Asymptomatic Adults

Screening for prediabetes and type 2 diabetes risk through a targeted assessment of risk factors (Table 2.5) or with an assessment tool, such as the ADA risk test (diabetes.org/diabetes-risk-test), is recommended to guide health care professionals on whether performing a diagnostic test (Table 2.1) is appropriate. Prediabetes and type 2 diabetes meet criteria for conditions in which early detection via screening is appropriate. Both conditions are common and impose significant clinical and public health burdens. There is often a long presymptomatic phase before the diagnosis, and simple tests to detect preclinical disease are readily available (96). The duration of glycemic burden is a strong predictor of adverse outcomes. When considering strategies to prevent the progression to diabetes in people with prediabetes, it is important to individualize the risk-to-benefit ratio of formal interventions and assess person-centered goals. Risk models have often found higher benefit of intervention in those at highest risk (97) (see section 3, “Prevention or Delay of Diabetes and Associated Comorbidities”) and reduced the risk of diabetes complications (98) (see section 10, “Cardiovascular Disease and Risk Management,” section 11, “Chronic Kidney Disease and Risk Management,” and section 12, “Retinopathy, Neuropathy, and Foot Care”). In the National Institutes of Health (NIH) Diabetes Prevention Program Outcomes Study (DPPOS) report, prevention of progression from prediabetes to diabetes (99) resulted in lower rates of developing retinopathy and nephropathy (100).

Despite the numerous benefits of screening and early diagnosis for prediabetes or diabetes, unfortunately many people in the U.S. and globally either remain undiagnosed or are diagnosed when complications have already arisen. Additional considerations regarding testing for type 2 diabetes and prediabetes in asymptomatic individuals are described below.

Age

Age is a major risk factor for diabetes. Testing should begin at no later than age 35 years for all people (101). Screening should be considered in adults of any age with overweight or obesity and one or more risk factors for diabetes.

Medications

Screening for prediabetes or diabetes should be performed in people treated with certain medications, such as glucocorticoids, statins (102), thiazide diuretics, some HIV medications (103), and second-generation antipsychotic medications (104), as these drug classes increase the risk of glucose abnormalities, often by promoting insulin resistance. Alone or in combination with other drugs, glucocorticoids are frequently used for the treatment of a diverse set of conditions including, but not limited to, autoimmune disease (e.g., rheumatoid arthritis), as part of the immunosuppression plan after transplant, or in conjunction with systemic anticancer therapy. The prevalence of glucocorticoid-induced diabetes (hyperglycemia in the absence of preexisting diabetes) depends on the drug's half-life, dose, and frequency of administration as well as the duration of treatment. In addition, the presence of common risk factors for diabetes (e.g., obesity) or concomitant use of other offending drugs (e.g., tacrolimus) may also play a role (105). Educating individuals treated with recurrent or long-term courses of glucocorticoids regarding symptoms of hyperglycemia is essential (107).

It is recommended that screening be performed by checking postprandial plasma glucose (possibly 1–2 h after meals) or random glucose rather than in the fasting state, as glucocorticoid-induced hyperglycemia generally results from insulin resistance. In fact, relying on fasting glucose as a screening test can lead to underdiagnosis and delayed treatment (105,108) (see also section 9, “Pharmacologic Approaches to Glycemic Treatment”).

The ideal frequency of glucose monitoring for screening is not established, as a direct comparison of different monitoring schedules and their diagnostic performance is not available. In the outpatient setting, a proposed pragmatic approach is to check glucose twice weekly and to intensify frequency to daily monitoring if glucose is ≥ 200 mg/dL (109). In the context of cancer treatment, other organizations advocate for daily capillary glucose monitoring and prompt institution of treatment for glucose ≥ 220 mg/dL on two occasions (108,110,111). People treated with second-generation antipsychotic medications require close monitoring because of an increased risk of type 2 diabetes associated with this class of medications, often clustering with dyslipidemia and weight gain. People treated with these agents should be screened for prediabetes or diabetes at baseline, re-screened 12–16 weeks after medication initiation, and annually thereafter (112). Repeat testing can occur sooner if clinically warranted.

People With HIV

People living with HIV are at higher risk for developing prediabetes and diabetes. In addition, some antiretroviral (ARV) therapies may further increase the risk.

2026 February 01 on guest by

http://diabetesjournals.org/care/article-pdf/49/Supplement_1/S27/848854/dc26s002.pdf from Downloaded

Diagnosis and Classification of Diabetes # Diabetes Care Volume 49, Supplement 1, January 2026

(113). Therefore, a screening protocol for community screening may be considered. on type 2 diabetes in children and prediabetes and type 2 diabetes is recommended (114). As the A1C test may underestimate glycemia in people with HIV, plasma glucose criteria are preferred to diagnose prediabetes and diabetes (17). Diabetes risk is increased with certain protease inhibitors (PIs) and nucleoside/nucleotide reverse transcriptase inhibitors (NRTIs). New-onset diabetes is estimated to occur in more than 5% of individuals infected with HIV on PIs, whereas more than 15% may have pre-diabetes (115). PIs are associated with insulin resistance and may also lead to apoptosis of pancreatic β -cells. NRTIs also affect fat distribution (both lipohypertrophy and lipoatrophy), which is associated with insulin resistance. For people with HIV and ARV-associated hyperglycemia, it may be appropriate to consider discontinuing the offending ARV agents if safe and effective

alternatives are available (116). Before making ARV substitutions, carefully consider the possible effect on HIV virological control and the potential adverse effects of new ARV agents. In some cases, glucose-lowering agents may still be necessary.

Testing Interval

The appropriate interval between screening tests is not known (117). The rationale for the 3-year interval is that with this interval, the number of false-positive tests that require confirmatory testing will be reduced, and individuals with false-negative tests will be retested before substantial time elapses and complications develop (117). In especially high-risk individuals such as those with previous values nearer to the diabetes diagnostic cut point or ongoing treatment with medications listed above, shorter intervals between screenings may be useful.

Screening and Testing for Prediabetes and Type 2 Diabetes in Children and Adolescents

The epidemiologic studies that formed the basis for the recommendations to use A1C and plasma glucose criteria to diagnose prediabetes and diabetes included only adult populations (122). However, ADA clinical guidance concluded that A1C, FPG, or 2-h PG also could be used to test for prediabetes or type 2 diabetes in children and adolescents (123). In the last decade, the incidence and prevalence of type 2 diabetes in children and adolescents has increased dramatically, especially in certain high-risk racial, ethnic, and ancestral subgroups (124). See Table 2.6 for recommendations on risk-based screening for type 2 diabetes or prediabetes in asymptomatic children and adolescents in a clinical setting (123). See Table 2.1 and Table 2.2 for the criteria for the diagnosis of diabetes and prediabetes, respectively, that apply to children, adolescents, and adults. See section 14, "Children and Adolescents," for additional information.

Community Screening

Ideally, screening should be carried out within a health care setting (including appropriately resourced pharmacies) because of the need for follow-up and treatment (118). Community screening outside a health care setting is generally not recommended because people with positive tests may not seek, or have access to, appropriate follow-up testing and care. However, in specific situations where an adequate referral system is established beforehand for positive tests, people receiving systemic anticancer treatment, as a direct effect of chemotherapeutic agents (e.g., busulfan) or the frequently associated glucocorticoid treatment (126), and represents a risk factor for poor cancer- and treatment-related outcomes (127,128).

Recommendations

1. People starting cancer treatment with immune checkpoint inhibitors (ICIs), including anti-PD-1 or anti-PDL-1 therapy (e.g., nivolumab, pembrolizumab, avelumab), phosphoinositidylinositol 3-kinase α (PI3K α) inhibitors (e.g., alpelisib, inavolisib), or mammalian target of rapamycin (mTOR) inhibitors (e.g., everolimus), should be educated regarding risks, symptoms, and signs of hyperglycemia and hyperglycemic crises. 2. In people treated with ICIs, fasting or random plasma glucose should be tested before initiating treatment, during each visit, or if symptoms and signs of hyperglycemia develop during or after treatment cessation. 3. In people treated with PI3K α inhibitors, fasting or random plasma glucose and A1C should be tested before initiating treatment, and random plasma glucose should be tested weekly for the first 2 weeks of treatment and then every 4 weeks during treatment. Consider testing A1C every 3 months during treatment. 4. In people treated with mTOR inhibitors, fasting or random plasma glucose should be tested before starting and at each visit throughout the duration of treatment. Consider testing A1C every 3 months during treatment.

Both acute and chronic endocrine disorders account for a large portion of adverse events reported with cancer treatment (125). Hyperglycemia occurs in 15–50% of people receiving systemic anticancer treatment, as a direct effect of chemotherapeutic agents (e.g., busulfan) or the frequently associated glucocorticoid treatment (126), and represents a risk factor for poor cancer- and treatment-related outcomes (127,128).

2026 February 01 on guest by

http://diabetesjournals.org/care/article-pdf/49/Supplement_1/S27/848854/dc26s002.pdf from Downloaded

diabetesjournals.org/care # Diagnosis and Classification of Diabetes

regarding ICIs, PD-1 (e.g., nivolumab) and PDL-1 inhibitors (e.g., durvalumab) and, to a lesser extent, anti-CTLA-4 antibodies (e.g., ipilimumab), are associated with new-onset autoimmune diabetes in addition to other immune-related adverse events. ICI-induced autoimmune diabetes occurs in 0.6–1.4% of treated people, often presents as DKA, and almost invariably requires lifelong insulin therapy (129). Despite a peak of incidence within 2–3 months from the initiation of treatment (67,130), new onset of ICI-induced diabetes can occur anytime during treatment and has been reported even after discontinuation of ICIs (65,130). Similar to people with newly diagnosed type 1 diabetes, people who develop ICI-induced diabetes should be promptly referred to an endocrinologist and receive appropriate diabetes self-management education. Thus, testing either fasting or random plasma glucose before initiating treatment and at each visit (or for intervening symptoms of hyperglycemia) is a reasonable approach to increase health care professionals' awareness of the metabolic risk of ICIs and reduce the risk of presentation with DKA. Further epidemiological insight and validation of HLA-based risk prediction scores may guide future targeted screening strategies.

Inhibitors targeting the α isoform of PI3K have been approved (e.g., alpelisib, inavolisib) for the treatment of advanced or metastatic breast cancer. Pivotal trials demonstrated that PI3K α inhibitors, when added to standard of care treatment, provided survival benefits but were also associated with the emergence of grade 3 or 4 hyperglycemia in up to 36% of participants, which was the most common reason for drug discontinuation (131,132). A retrospective analysis of adverse events accrued in a phase 3 randomized controlled trial (RCT) showed that the median time to onset for alpelisib-induced hyperglycemia was 13 days, with a range extending to approximately 1 year (133). The current data inform the recommendation to check fasting and random plasma glucose before initiating treatment and to monitor random plasma glucose weekly for the first 2 weeks of therapy and then every 4 weeks or sooner if symptoms of hyperglycemia develop in the interim. Testing A1C every 3 months can also be considered, while recognizing that alone A1C may not capture the early peak of hyperglycemia noted with PI3K α inhibitors. Inhibitors of mTOR (e.g., everolimus) are associated with hyperglycemia of variable severity but without a definitive peak of onset. In a large meta-analysis that included 3,900 participants treated with everolimus across nine RCTs, the incidence of all-grade hyperglycemia was as high as 27%, whereas grades 3 and 4 were reported in 2.5% of participants (134). In a single-center retrospective cohort study in people with cancer treated with everolimus, the estimated median time to onset of hyperglycemia was 5–6 months, but the incidence distribution extended from a few weeks to several years after the initiation of treatment (135). In the absence of a clear-cut temporal pattern for the onset of hyperglycemia, fasting or random plasma glucose should be checked before treatment initiation and at each visit during treatment with mTOR inhibitors, or sooner if symptoms of hyperglycemia develop in the interim, while monitoring A1C every 3 months can also be considered.

Finally, hyperglycemia is also a relatively frequent complication of Akt inhibitors (e.g., capivasertib) and antibody drug conjugates (e.g., enfortumab vedotin), which will not be reviewed.

PANCREATIC DIABETES OR DIABETES IN THE CONTEXT OF DISEASE OF THE EXOCRINE PANCREAS

Recommendations

- 2.23 Screen people for diabetes within 3–6 months following an episode of acute pancreatitis and annually thereafter. Screening for diabetes is recommended annually for people with chronic pancreatitis. E - 2.24a Annual screening for cystic fibrosis–related diabetes (CFRD) should begin by age 10 years in all people with cystic fibrosis, preferably using OGTT. B - 2.24b If an OGTT is not feasible, A1C can be used as an alternative method as part of a two-step screening strategy. Individuals with A1C values between 5.5% and 6.4% (37 and 47 mmol/mol, respectively) should undergo an OGTT within 3 months. C An A1C value of $\geq 6.5\%$ (≥ 48 mmol/mol) is consistent with a diagnosis of CFRD. B - 2.25 Beginning 5 years after the diagnosis of CFRD, annual monitoring for complications of diabetes is recommended. E

Pancreatic diabetes (also termed pancreatogenic diabetes or type 3c diabetes) includes both structural (e.g., destruction or removal of normal pancreatic tissue) and functional loss of insulin secretion in the context of exocrine pancreatic insufficiency and is commonly misdiagnosed as type 2 diabetes. The diverse set of etiologies includes pancreatitis (acute and chronic pancreatic inflammation and associated fibrosis leading to loss of functional exocrine and endocrine pancreatic function), cystic fibrosis, trauma or pancreatectomy, neoplasia, cystic fibrosis (addressed later in this section), hemochromatosis, fibrocalculous pancreatopathy, rare genetic disorders, and idiopathic forms (2); as such, pancreatic diabetes is the preferred umbrella term

(136).

Acute (even a single bout) and chronic pancreatitis can lead to postpancreatitis diabetes mellitus (137). A distinguishing feature is concurrent pancreatic exocrine insufficiency (consider screening individuals with acute and chronic pancreatitis for exocrine pancreatic insufficiency by measuring fecal elastase), pathological pancreatic imaging (endoscopic ultrasound, MRI, and computed tomography), and absence of type 1 diabetes–associated autoimmunity (138–141). Risk for microvascular complications appears to be similar to that of other forms of diabetes.

For people with pancreatitis and diabetes, glucose-lowering therapies potentially associated with increased risk of pancreatitis (i.e., incretin-based therapies) should be avoided. Early initiation of insulin therapy should be considered. In the context of pancreatectomy, islet autotransplantation can be considered for selected individuals with medically refractory chronic pancreatitis in specialized centers to preserve, fully or partially, endogenous islet function and insulin secretion (142,143).

2026 February 01 on guest by

http://diabetesjournals.org/care/article-pdf/49/Supplement_1/S27/848854/dc26s002.pdf from Downloaded

Diagnosis and Classification of Diabetes Diabetes Care Volume 49, Supplement 1, January 2026

Recessive mutations in the gene encoding the cystic fibrosis transmembrane conductance regulator (CFTR) protein. Cystic fibrosis–related diabetes (CFRD) is a common comorbidity in people living with cystic fibrosis that is distinct from type 1 or type 2 diabetes and is often accompanied by pancreatic exocrine insufficiency (144). CFRD occurs in about 20% of adolescents and 50% of adults (145) and is associated with worse nutritional status and increased morbidity and mortality (146). Although insufficient insulin secretion is the primary defect in CFRD, insulin resistance can also occur in the setting of intervening illnesses and treatment with glucocorticoids. Current guidelines recommend screening for CFRD in people living with cystic fibrosis starting at age 10 years. Although screening for diabetes before the age of 10 years can identify risk for progression to CFRD in those with IGT, no benefit has been established with respect to weight, height, BMI, or lung function (145). Based on its elevated predictive value (147), OGTT is the recommended screening test for CFRD in all people living with cystic fibrosis, if feasible, and should be repeated annually. However, engagement in current CFRD screening guidelines is poor, with only 30–40% of adults with cystic fibrosis having annual OGTTs (148,149).

In this context, other professional organizations have advocated for the use of A1C as part of a two-step screening strategy as an alternative to OGTT and to maximize the diagnostic yield of OGTT (150). Specifically, a retrospective study investigating the diagnostic performance of OGTT vis-a-vis A1C in 256 community-dwelling people living with cystic fibrosis, age 10–18 years, showed that A1C of 5.5% and 5.8% provided a sensitivity of 91% and 95%, respectively, to detect CFRD by OGTT (151). A1C 5.5% also afforded a negative predictive value for CFRD of approximately 98%, thus making A1C an effective first-line screening tool to identify people living with cystic fibrosis who should not undergo further testing with OGTT. In this framework alternative to the annual OGTT, only people living with cystic fibrosis with A1C 5.5–6.4% would proceed with OGTT. A1C $\geq 6.5\%$ (≥ 48 mmol/mol) is consistent with a diagnosis of CFRD (152–154) and should be followed by a confirmatory test (preferably a repeat A1C or an OGTT).

POSTTRANSPLANTATION DIABETES MELLITUS

Recommendations

1. After organ transplantation, screening for hyperglycemia should be done. A formal diagnosis of post-transplantation diabetes mellitus (PTDM) is best made once the individual is stable on an immunosuppressive plan (usually 3 months) and in the absence of acute infection (158,161–163,166).
2. OGTT is the recommended test for the diagnosis of PTDM as early as 3 months posttransplant (158). However, screening people with FPG and/or A1C can identify high-risk individuals who require further assessment and may reduce the number of overall OGTTs required.
3. Few RCTs have reported on the short- and long-term use of glucose-lowering medications in the setting of PTDM (163,167,168). Most studies have reported that transplant individuals with hyperglycemia and PTDM after transplantation have higher rates of rejection, infection, and rehospitalization (161,163,169). Insulin therapy is the agent of choice for the management of hyperglycemia and diabetes in the hospital setting and can be continued.

Several terms are used in the literature to describe the presence of diabetes following organ transplantation (157). New-onset diabetes after transplantation (NODAT) is one such designation that describes individuals who develop new-onset diabetes following transplant. NODAT is consistent with a diagnosis of CFRD (152–154) and should be followed by a confirmatory test (preferably a repeat A1C or an OGTT).

2026 February 01 on guest by

http://diabetesjournals.org/care/article-pdf/49/Supplement_1/S27/848854/dc26s002.pdf from Downloaded

diabetesjournals.org/care # Diagnosis and Classification of Diabetes

postdischarge. Noninsulin glucose-lowering therapies can also be used for long-term management. The choice of agent is usually made based on the side effect profile of the medication, possible interactions with the individual's immunosuppression plan, and potential cardiovascular and kidney benefits in individuals with PTDM (163). See section 9, "Pharmacologic Approaches to Glycemic Treatment," for further information.

Neonatal Diabetes

Diabetes occurring under 6 months of age is termed neonatal diabetes, and about 80–85% of cases can be found to have an underlying monogenic cause (32,172–175). Neonatal diabetes occurs much less often after 6 months of age, whereas autoimmune type 1 diabetes rarely occurs before 6 months of age. Neonatal diabetes can either be transient or permanent. Transient diabetes is most often due to overexpression of genes on chromosome 6q24, is recurrent in about half of cases, and may be treatable with medications other than insulin. Permanent neonatal diabetes is most commonly due to autosomal dominant mutations in the genes encoding the Kir6.2 subunit (KCNJ11) and SUR1 subunit (ABCC8) of the β -cell K_{ATP} channel. The ADA-European Association for the Study of Diabetes type 1 diabetes consensus report recommends that regardless of current age, individuals diagnosed under 6 months of age should have genetic testing (32). Correct diagnosis has critical implications, because 30–50% of people with K_{ATP}-related neonatal diabetes will exhibit improved blood glucose levels when treated with high-dose oral sulfonylureas instead of insulin. Insulin gene (INS) mutations are the second most common cause of permanent neonatal diabetes, with insulin therapy being the preferred treatment strategy. Additionally, diagnosis can lead to identification of other affected family members and can indicate potential extrapancreatic complications in affected individuals.

Recommendations

- 2.29a Regardless of current age, all people diagnosed with diabetes in the first 6 months of life should have genetic testing for neonatal diabetes. - 2.29b Children and young adults who do not have typical characteristics of type 1 or type 2 diabetes and have a family history of diabetes in successive generations (suggestive of an autosomal-dominant pattern of inheritance) should have genetic testing for maturity-onset diabetes of the young (MODY). - 2.29c In both instances, consultation with a center specializing in diabetes genetics is recommended to understand the significance of genetic mutations and how best to approach further evaluation, treatment, and genetic counseling.

Diagnosis of Monogenic Diabetes

The diagnosis of monogenic diabetes should be considered in children and adults diagnosed with diabetes in early adulthood with the following findings:

- • Diabetes diagnosed within the first 6 months of life (170,172) - • Diabetes without typical features of type 1 or type 2 diabetes (negative autoantibodies for type 1 diabetes precludes further testing for monogenic diabetes, but the presence of autoantibodies in people with monogenic diabetes has been reported.)

Maturity-Onset Diabetes of the Young

MODY is frequently characterized by onset of hyperglycemia at an early age (classically before age 25 years, although diagnosis may occur at older ages). MODY is characterized by impaired insulin secretion with minimal or no defects in insulin action (in the absence of coexistent obesity). It is inherited in an autosomal dominant pattern with abnormalities in at least 14 genes on different chromosomes identified to date (170). The most common forms are GCK-MODY (MODY2), HNF1A-MODY (MODY3), and HNF4A-MODY (MODY1). Correct diagnosis of monogenic forms of diabetes is critical because people who have them may be incorrectly diagnosed with type 1 or type 2 diabetes, leading to suboptimal, even potentially harmful, treatment plans and

delays in diagnosing other family members (170).

GESTATIONAL DIABETES MELLITUS

Recommendations

- 2.30 In individuals who are planning pregnancy, screen those with risk factors (Table 2.5) B and consider testing all individuals of childbearing potential for undiagnosed prediabetes or diabetes. E

2026 February 01 on guest by

http://diabetesjournals.org/care/article-pdf/49/Supplement_1/S27/848854/dc26s002.pdf from Downloaded

Diagnosis and Classification of Diabetes Diabetes Care Volume 49, Supplement 1, January 2026

Table 2.7—Most common causes of monogenic diabetes

Gene	Inheritance	Clinical features				
MODY	HNF1A	AD	HNF1A-MODY: progressive insulin secretory defect with presentation in adolescence or early adulthood; lowered renal threshold for glucosuria; large rise in 2-h PG level on OGTT (>90 mg/dL [>5 mmol/L]); low hs-CRP; sensitive to sulfonylureas	GCK	AD	GCK-MODY: higher glucose threshold (set point) for glucose-stimulated insulin secretion, causing stable, nonprogressive elevated fasting blood glucose; typically does not require treatment in nonpregnant individuals; microvascular complications are rare; small rise in 2-h PG level on OGTT (<54 mg/dL [<3 mmol/L])
HNF4A	AD	HNF4A-MODY: progressive insulin secretory defect with presentation in adolescence or early adulthood; may have large birth weight (macrosomia) and transient neonatal hypoglycemia; sensitive to sulfonylureas	HNF1B	AD	HNF1B-MODY: developmental renal disease (typically cystic); genitourinary abnormalities; atrophy of the pancreas; hyperuricemia; gout	Neonatal diabetes
Permanent	KCNJ11	AD	IUGR; possible developmental delay and seizures; responsive to sulfonylureas	ABCC8	AD	IUGR; rarely developmental delay; responsive to sulfonylureas
INS	AD	IUGR; insulin requiring	FOXP3	X-linked	Immunodysregulation, polyendocrinopathy, enteropathy X-linked (IPEX) syndrome: autoimmune diabetes, autoimmune thyroid disease, exfoliative dermatitis; insulin requiring	Transient
6q24 (PLAGL1, HYMA1)	AD	for paternal duplications	IUGR; macroglossia; umbilical hernia; may be treatable with medications other than insulin			

Adapted from Carmody et al. (171). AD, autosomal dominant; AR, autosomal recessive; IUGR, intrauterine growth restriction; OGTT, oral glucose tolerance test; UPD6, uniparental disomy of chromosome 6; 2-h PG, 2-h plasma glucose.

2.31a

Before 15 weeks of gestation, test individuals with risk factors (Table 2.5) and consider testing all individuals for undiagnosed diabetes at the first prenatal visit using standard diagnostic criteria if not screened preconception.

2.31b

Before 15 weeks of gestation, screen for abnormal glucose metabolism (defined as A1C 5.9–6.4% [41–47 mmol/mol] or FPG 110–125 mg/dL [6.1–6.9 mmol/L]) to identify individuals who are at higher risk of adverse pregnancy and neonatal outcomes and are at high risk of a later gestational diabetes mellitus (GDM) diagnosis.

2.32

Screen for GDM at 24–28 weeks of gestation in pregnant individuals not previously found to have diabetes or high-risk abnormal glucose metabolism detected earlier in the current pregnancy.

2.33

Screen individuals with GDM for prediabetes or diabetes at 4–12 weeks postpartum, using the 75-g OGTT and clinically appropriate non-pregnancy diagnostic criteria.

Individuals with a history of GDM should have lifelong screening for the development of prediabetes or diabetes every 1–3 years. Ideally, undiagnosed diabetes should be identified preconception in individuals with risk

factors or in high-risk populations, as they are likely to benefit from preconception care.

The preconception care of people with known preexisting diabetes results in lower A1C and reduced risk of birth defects, preterm delivery, perinatal mortality, small-for-gestational-age birth weight, and neonatal intensive care unit admission. If individuals are not screened prior to pregnancy, universal early screening at 15 weeks of gestation for undiagnosed diabetes may be considered over selective screening (Table 2.5), particularly in populations with high prevalence of risk factors and undiagnosed diabetes in people of childbearing age. Strong racial and ethnic disparities exist in the prevalence of undiagnosed diabetes. Therefore, early screening provides an initial step to identify and address these health disparities.

2026 February 01 on guest by

http://diabetesjournals.org/care/article-pdf/49/Supplement_1/S27/848854/dc26s002.pdf from Downloaded

diabetesjournals.org/care

Diagnosis and Classification of Diabetes

for identifying undiagnosed diabetes in early pregnancy are the same as those used in nonpregnant individuals (Table 2.1). Individuals found to have diabetes should be classified as having diabetes complicating pregnancy (most often type 2 diabetes, rarely type 1 diabetes or monogenic diabetes) and managed accordingly. Early abnormal glucose metabolism, defined as a fasting glucose threshold of 110 mg/dL (6.1 mmol/L) or an A1C of 5.9% (41 mmol/mol), may identify individuals who are at higher risk of adverse pregnancy and neonatal outcomes (pre-eclampsia, macrosomia, shoulder dystocia, and perinatal death) and are at high risk of a later GDM diagnosis (191–194). Some, but not all, studies showed that an A1C threshold of 5.7% (39 mmol/L) was associated with adverse perinatal outcomes (193,195–197). Professional society guidance regarding early-pregnancy diagnosis and treatment of hyperglycemia less than overt diabetes differs (see section 15, “Management of Diabetes in Pregnancy”). If early screening for undiagnosed diabetes was negative, individuals should be rescreened for GDM between 24 and 28 weeks of gestation, and individuals not previously screened should be screened for GDM.

Diagnosis

GDM carries risks for the mother, fetus, and neonate. The Hyperglycemia and Adverse Pregnancy Outcome (HAPO) study (208), a large-scale multinational cohort study completed by more than 23,000 pregnant individuals, demonstrated that risk of adverse maternal, fetal, and neonatal outcomes continuously increased as a function of maternal glycemia at 24–28 weeks of gestation, even within ranges previously considered normal for pregnancy. For most complications, there was no threshold for risk. These results have led to careful reconsideration of the diagnostic criteria for GDM.

GDM diagnosis (Table 2.8) can be accomplished with either of two strategies:

1. The “one-step” 75-g OGTT derived from the IADPSG criteria, or 2. The older “two-step” approach with a 50-g (nonfasting) screen followed by a 100-g OGTT for those who screen positive based on the work of Carpenter-Coustan’s interpretation of the older O’Sullivan and Mahan criteria (209).

| One-step strategy | | -----
----- | -----
----- | | Perform a 75-g OGTT, with plasma glucose measurement when an individual is fasting and at 1 and 2 h, at 24–28 weeks of gestation in individuals not previously diagnosed with diabetes. | The OGTT should be performed in the morning after an overnight fast of at least 8 h. | | The diagnosis of GDM is made when any of the following plasma glucose values are met or exceeded: | | Fasting: 92 mg/dL (5.1 mmol/L) | 1 h: 180 mg/dL (10.0 mmol/L) | 2 h: 153 mg/dL (8.5 mmol/L) | | Two-step strategy | | Step 1: | Perform a 50-g GLT (nonfasting), with plasma glucose measurement at 1 h, at 24–28 weeks of gestation in individuals not previously diagnosed with diabetes. | | If the plasma glucose level measured 1 h after the load is ≥130, 135, or 140 mg/dL (7.2, 7.5, or 7.8 mmol/L, respectively),* proceed to a 100-g OGTT. | | Step 2: | The 100-g OGTT should be performed when the individual is fasting. | | The diagnosis of GDM is made when at least two of the following four plasma glucose levels (measured fasting and at 1, 2, and 3 h during OGTT) are met or exceeded: | |

Different diagnostic criteria will identify different degrees of maternal hyperglycemia and maternal/fetal risk, leading experts to debate optimal strategies for the diagnosis of GDM.

One-Step Strategy

The IADPSG examined data from the HAPO study and defined diagnostic cut points for GDM as the average fasting, 1-h, and 2-h PG values during a 75-g OGTT in individuals at 24–28 weeks of gestation, wherein the cut points were those at which odds for adverse outcomes reached 1.75 times the estimated odds. This one-step strategy was anticipated to significantly increase the incidence of GDM (from 5 to 6% to 15–20%), primarily because only one abnormal value, not two, became sufficient to make the diagnosis (210). Many regional studies have seen a roughly one- to threefold increase in GDM cases using the IADPSG criteria (211). A study of pregnancy OGTTs with glucose levels blinded to caregivers found that 11 years after their pregnancies, individuals who would have been diagnosed with GDM by the one-step approach, as

2026 February 01 on guest by

http://diabetesjournals.org/care/article-pdf/49/Supplement_1/S27/848854/dc26s002.pdf from Downloaded

Diagnosis and Classification of Diabetes Diabetes Care Volume 49, Supplement 1, January 2026

Compared with those without GDM, were higher prevalence of IGT; higher 30-min, 1-h, and 2-h glucose levels during the OGTT; and reduced insulin sensitivity and oral disposition index in their offspring at 10–14 years of age compared with offspring of mothers without GDM. Associations of mother's fasting, 1-h, and 2-h values on the 75-g OGTT were continuous with a comprehensive panel of offspring metabolic outcomes. The ADA recommends the IADPSG diagnostic criteria to optimize gestational outcomes, because these criteria are the only ones based on pregnancy outcomes rather than end points such as prediction of subsequent maternal diabetes.

Expected benefits of using IADPSG criteria for offspring are inferred from intervention trials focusing on individuals with lower levels of hyperglycemia than those identified using older GDM diagnostic criteria. Those trials found modest benefits, including reduced rates of large-for-gestational-age births and preeclampsia. Of note, 80–90% of participants being treated for mild GDM in these two RCTs could be managed with lifestyle therapy alone. The OGTT glucose cutoffs in these two trials overlapped the thresholds recommended by the IADPSG, and in one trial, the 2-h PG threshold (140 mg/dL [7.8 mmol/L]) was lower than the cutoff recommended by the IADPSG (153 mg/dL [8.5 mmol/L]).

No RCTs of treating versus not treating GDM diagnosed by different criteria have been published to date. However, a randomized trial of testing for GDM at 24–28 weeks of gestation by the one-step method using IADPSG criteria versus the two-step method by Carpenter-Coustan criteria identified twice as many individuals with GDM using the one-step method. Despite treating more individuals for GDM using the one-step method, there was no difference in pregnancy and perinatal complications, though concerns were raised about sample size estimates and unanticipated suboptimal engagement with the screening and treatment protocol. For example, in the two-step group, 165 participants not counted as having GDM were treated for isolated elevated FPG >95 mg/dL (>5.3 mmol/L).

The two-step strategy: In 2013, the NIH convened a consensus development conference to consider diagnostic criteria for diagnosing GDM. The panel recommended continuing a two-step approach to screening that used a 1-h 50-g glucose loading test (GLT) followed by a 3-h 100-g OGTT for those who screened positive. The American College of Obstetricians and Gynecologists (ACOG) recommends any of the commonly used thresholds of 130, 135, or 140 mg/dL for the 1-h 50-g GLT.

Future considerations: Data exist to support each strategy, as demonstrated by differing recommendations by expert groups. A systematic review of economic evaluations of GDM screening found that the one-step method identified more cases of GDM and was more likely to be cost-effective than the two-step method. The decision of which strategy to implement must therefore be made based on the relative values placed on factors that have yet to be measured (e.g., willingness to change practice based on correlation studies rather than intervention trial results, available infrastructure, and importance of cost considerations).

The IADPSG criteria (one-step strategy) have been adopted internationally as the preferred approach. Data that compare population-wide outcomes with one-step versus two-step approaches have been inconsistent to

date. Pregnancies complicated by GDM per the IADPSG criteria, but not recognized as such, have outcomes comparable to pregnancies with diagnosed GDM by the more stringent two-step criteria. There remains strong consensus that establishing...

2026 February 01 on guest by

http://diabetesjournals.org/care/article-pdf/49/Supplement_1/S27/848854/dc26s002.pdf from Downloaded

diabetesjournals.org/care # Diagnosis and Classification of Diabetes

a uniform approach to diagnosing GDM will benefit people with GDM, caregivers, and policymakers. Longer-term outcome studies are currently underway.

References

1. Sacks DB, Arnold M, Bakris GL, et al. Guidelines and recommendations for laboratory analysis in the diagnosis and management of diabetes mellitus. *Diabetes Care* 2023;46:e151–e199
2. American Diabetes Association. Diagnosis and classification of diabetes mellitus. *Diabetes Care* 2014;37(Suppl. 1):S81–S90
3. International Expert Committee. International Expert Committee report on the role of the A1C assay in the diagnosis of diabetes. *Diabetes Care* 2009;32:1327–1334
4. Meijnikman AS, De Block CEM, Dirinck E, et al. Not performing an OGTT results in significant underdiagnosis of (pre)diabetes in a high risk adult Caucasian population. *Int J Obes (Lond)* 2017;41:1615–1620
5. Knowler WC, Barrett-Connor E, Fowler SE, et al. Reduction in the incidence of type 2 diabetes with lifestyle intervention or metformin. *N Engl J Med* 2002;346:393–403
6. Tuomilehto J, Lindstrom J, Eriksson JG, et al.; Finnish Diabetes Prevention Study Group. Prevention of type 2 diabetes mellitus by changes in lifestyle among subjects with impaired glucose tolerance. *N Engl J Med* 2001;344:1343–1350
7. Diabetes Prevention Program Research Group. HbA1c as a predictor of diabetes and as an outcome in the diabetes prevention program: a randomized clinical trial. *Diabetes Care* 2015;38:51–58
8. Echouffo-Tcheugui JB, Selvin E. Prediabetes and what it means: the epidemiological evidence. *Annu Rev Public Health* 2021;42:59–77
9. Chadha C, Pittas AG, Lary CW, et al.; D2d Research Group. Reproducibility of a prediabetes classification in a contemporary population. *Metabol Open* 2020;6:100031
10. Expert Committee on the Diagnosis and Classification of Diabetes Mellitus. Report of the Expert Committee on the Diagnosis and Classification of Diabetes Mellitus. *Diabetes Care* 1997;20:1183–1197
11. Expert Committee on the Diagnosis and Classification of Diabetes Mellitus. Report of the Expert Committee on the Diagnosis and Classification of Diabetes Mellitus. *Diabetes Care* 2003;26(Suppl. 1):S5–S20
12. Klein KR, Walker CP, McFerren AL, Huffman H, Frohlich F, Buse JB. Carbohydrate intake prior to oral glucose tolerance testing. *J Endocr Soc* 2021;5:bvab049
13. Conn JW. Interpretation of the glucose tolerance test. The necessity of a standard preparatory diet. *Am J Med Sci* 1940;199:555–563
14. Wilkerson HL, Butler FK, Francis JO. The effect of prior carbohydrate intake on the oral glucose tolerance test. *Diabetes* 1960;9:386–391
15. Centers for Medicare & Medicaid Services. CLIA Brochures. Accessed 12 August 2025. Available from https://www.cms.gov/regulations-and-guidance/legislation/clia/downloads/clia_compbrochure_508.pdf
16. Nathan DM, Griffin A, Perez FM, Basque E, Do L, Steiner B. Accuracy of a point-of-care hemoglobin A1c assay. *J Diabetes Sci Technol* 2019;13:1149–1153
17. Smith K, Deutsch AJ, McGrail C, et al. Multi-ancestry polygenic mechanisms of type 2 diabetes. *Nat Med* 2024;30:1065–1074
18. Steck AK, Vehik K, Bonifacio E, et al.; TEDDY Study Group. Predictors of progression from the appearance of islet autoantibodies to early childhood diabetes: The Environmental Determinants of Diabetes in the Young (TEDDY) study. *Diabetologia* 2015;58:980–987

2026 February 01 on guest by

http://diabetesjournals.org/care/article-pdf/49/Supplement_1/S27/848854/dc26s002.pdf from Downloaded

Diagnosis and Classification of Diabetes # Diabetes Care Volume 49, Supplement 1, January 2026

Diabetes in the Young (TEDDY). *Diabetes Care* 2015;38:808–813

induced by immune checkpoint inhibitors: a genetic risk score for use in newborn screening and systematic review. *Cancer Immunol Immunother* 2021;70:1527–1540

incident diagnosis. *Diabetes Care* 2019;42:200–207

1. McKeigue PM, Spiliopoulou A, McGurnaghan S, et al. Persistent C-peptide secretion in type 1 diabetes and its relationship to the genetic architecture of diabetes. *BMC Med* 2019;17:165
2. Sosenko JM, Palmer JP,

Rafkin-Mervis L, et al.; Diabetes Prevention Trial-Type 1 Study Group. Incident dysglycemia and progression to type 1 diabetes among participants in the Diabetes Prevention Trial-Type 1. *Diabetes Care* 2009;32:1603–1607

3. Krischer JP; Type 1 Diabetes TrialNet Study Group. The use of intermediate endpoints in the design of type 1 diabetes prevention trials. *Diabetologia* 2013;56:1919–1924

4. Triolo TM, Chase HP, Barker JM; DPT-1 Study Group. Diabetic subjects diagnosed through the Diabetes Prevention Trial-Type 1 (DPT-1) are often asymptomatic with normal A1C at diabetes onset. *Diabetes Care* 2009;32:769–773

5. Phillip M, Achenbach P, Addala A, et al. Consensus guidance for monitoring individuals with islet autoantibody-positive pre-stage 3 type 1 diabetes. *Diabetes Care* 2024;47:1276–1298

6. Sims EK, Cuthbertson D, Ferrat LA, et al. IA-2A positivity increases risk of progression within and across established stages of type 1 diabetes. *Diabetologia* 2025;68:993–1004

7. Ziegler A-G, Bonifacio E. Why is the presence of autoantibodies against GAD associated with a relatively slow progression to clinical diabetes? *Diabetologia* 2020;63:1665–1666

8. Greenbaum CJ, Beam CA, Boulware D, et al.; Type 1 Diabetes TrialNet Study Group. Fall in C-peptide during first 2 years from diagnosis: evidence of at least two distinct phases from composite Type 1 Diabetes TrialNet data. *Diabetes* 2012;61:2066–2073

9. Hope SV, Wienand-Barnett S, Shepherd M, et al. Practical classification guidelines for diabetes in patients treated with insulin: a cross-sectional study of the accuracy of diabetes diagnosis. *Br J Gen Pract* 2016;66:e315–e322

10. Mishra R, Hodge KM, Cousminer DL, Leslie RD, Grant SFA. A global perspective of latent autoimmune diabetes in adults. *Trends Endocrinol Metab* 2018;29:638–650

11. Buzzetti R, Zampetti S, Maddaloni E. Adult-onset autoimmune diabetes: current knowledge and implications for management. *Nat Rev Endocrinol* 2017;13:674–686

12. Ben-Skowronek I. IPEX syndrome: genetics and treatment options. *Genes (Basel)* 2021;12:323

13. Frommer L, Kahaly GJ. Autoimmune polyendocrinopathy. *J Clin Endocrinol Metab* 2019;104:4769–4782

14. Wang Y, Guo H, Wang G, Zhai J, Du B. COVID-19 as a trigger for type 1 diabetes. *J Clin Endocrinol Metab* 2023;108:2176–2183

15. Geng A, Tang H, Huang J, et al. The deacetylase SIRT6 promotes the repair of UV-induced DNA damage by targeting DDB2. *Nucleic Acids Res* 2020;48:9181–9194

16. Betsch C, Korn L, Sprengholz P, et al. Social and behavioral consequences of mask policies during the COVID-19 pandemic. *Proc Natl Acad Sci U S A* 2020;117:21851–21853

17. Dendrite Clinical Systems. CoviDIAB Registry. Accessed 12 August 2025. Available from <https://covidiab.e-dendrite.com/>

18. Zhao Z, Wang X, Bao X-Q, Ning J, Shang M, Zhang D. Autoimmune polyendocrine syndrome

19. Sharp SA, Rich SS, Wood AR, et al. Development and standardization of an improved type 1 diabetes

2026 February 01 on guest by

http://diabetesjournals.org/care/article-pdf/49/Supplement_1/S27/848854/dc26s002.pdf

from Downloaded

diabetesjournals.org/care # Diagnosis and Classification of Diabetes

glycemic status among middle-aged and older adults adults with severe mental illness who take preexisting diabetes mellitus: a systematic review in the United States, 2006-2014. *Diabetes Res Clin Pract* 2021;174:108726

97. Sussman JB, Kent DM, Nelson JP, Hayward RA. Improving diabetes prevention with benefit based tailored treatment: risk based reanalysis of Diabetes Prevention Program. *BMJ* 2015;350:h454

98. Palladino R, Tabak AG, Khunti K, et al. Association between pre-diabetes and microvascular and macrovascular disease in newly diagnosed type 2 diabetes. *BMJ Open Diabetes Res Care* 2020;8:e001061

99. Perreault L, Pan Q, Aroda VR, et al.; Diabetes Prevention Program Research Group. Exploring residual risk for diabetes and microvascular disease in the Diabetes Prevention Program Outcomes Study (DPPOS). *Diabet Med* 2017;34:1747–1755

100. Nathan DM, Bennett PH, Crandall JP, et al.; Research Group. Does diabetes prevention translate into reduced long-term vascular complications of diabetes? *Diabetologia* 2019;62:1319–1328

101. Chung S, Azar KMJ, Baek M, Lauderdale DS, Palaniappan LP. Reconsidering the age thresholds for type II diabetes screening in the U.S. *Am J Prev Med* 2014;47:375–381

102. Mansi IA, Sumithran P, Kinaan M. Risk of diabetes with statins. *BMJ* 2023;381:e071727

103. Eckhardt BJ, Holzman RS, Kwan CK, Baghdadi J, Aberg JA. Glycated hemoglobin A(1c) as screening for diabetes mellitus in HIV-infected individuals. *AIDS Patient Care STDS* 2012;26:197–201
104. Vancampfort D, Correll CU, Galling B, et al. Diabetes mellitus in people with schizophrenia, bipolar disorder and major depressive disorder: a systematic review and large scale meta-analysis. *World Psychiatry* 2016;15:166–174
105. Katsuyama T, Sada K-E, Namba S, et al. Risk factors for the development of glucocorticoid-induced diabetes mellitus. *Diabetes Res Clin Pract* 2015;108:273–279
106. Liu X-X, Zhu X-M, Miao Q, Ye H-Y, Zhang Z-Y, Li Y-M. Hyperglycemia induced by glucocorticoids in nondiabetic patients: a meta-analysis. *Ann Nutr Metab* 2014;65:324–332
107. Movahedi M, Beauchamp M-E, Abrahamowicz M, et al. Risk of incident diabetes mellitus associated with the dosage and duration of oral glucocorticoid therapy in patients with rheumatoid arthritis. *Arthritis Rheumatol* 2016;68:1089–1098
108. Cansu GB, Cansu DU, Taşkınran B, Bilge SY, Bilgin M, Korkmaz C. What is the optimal time for measuring glucose concentration to detect steroid-induced hyperglycemia in patients with rheumatic diseases? *Clin Biochem* 2019;67:33–39
109. Barker HL, Morrison D, Llano A, Sainsbury CAR, Jones GC. Practical guide to glucocorticoid induced hyperglycaemia and diabetes. *Diabetes Ther* 2023;14:937–945
110. Joint British Diabetes Societies for Inpatient Care. The Management of Glycaemic Control in People With Cancer, 2021. Accessed 12 August 2025. Available from link
111. Roberts A, James J, Dhatariya K; Joint British Diabetes Societies (JBDS) for Inpatient Care. Management of hyperglycaemia and steroid (glucocorticoid) therapy: a guideline from the Joint British Diabetes Societies (JBDS) for Inpatient Care group. *Diabet Med* 2018;35:1011–1017
112. Mangurian C, Newcomer JW, Vittinghoff E, et al. Diabetes screening among underserved populations. *J Gen Intern Med* 2019;34:1234–1240
113. Namara D, Schwartz JI, Tusubira AK, et al. The risk of hyperglycemia associated with use of dolutegravir among adults living with HIV in Kampala, Uganda: a case-control study. *Int J STD AIDS* 2022;33:1158–1164
114. Schambelan M, Benson CA, Carr A, et al.; International AIDS Society-USA. Management of metabolic complications associated with antiretroviral therapy for HIV-1 infection: recommendations of an International AIDS Society-USA panel. *J Acquir Immune Defic Syndr* 2002;31:257–275
115. Monroe AK, Glesby MJ, Brown TT. Diagnosing and managing diabetes in HIV-infected patients: current concepts. *Clin Infect Dis* 2015;60:453–462
116. Wohl DA, McComsey G, Tebas P, et al. Current concepts in the diagnosis and management of metabolic complications of HIV infection and its therapy. *Clin Infect Dis* 2006;43:645–653
117. Johnson SL, Tabaei BP, Herman WH. The efficacy and cost of alternative strategies for systematic screening for type 2 diabetes in the U.S. population 45–74 years of age. *Diabetes Care* 2005;28:307–311
118. Kaul P, Chu LM, Dover DC, Yeung RO, Eurich DT, Butalia S. Disparities in adherence to diabetes screening guidelines among males and females in a universal care setting: a population-based study of 1,380,697 adults. *Lancet Reg Health Am* 2022;14:100320
119. Tabaei BP, Burke R, Constance A, et al. Community-based screening for diabetes in Michigan. *Diabetes Care* 2003;26:668–670
120. Lalla E, Cheng B, Kunzel C, Burkett S, Lamster IB. Dental findings and identification of undiagnosed hyperglycemia. *J Dent Res* 2013;92:888–892
121. Herman WH, Taylor GW, Jacobson JJ, Burke R, Brown MB. Screening for prediabetes and type 2 diabetes in dental offices. *J Public Health Dent* 2015;75:175–182

122. Cowie CC, Rust KF, Byrd-Holt DD, et al. Prevalence of diabetes and high risk for diabetes using A1C criteria in the U.S. population in 1988–2006. *Diabetes Care* 2010;33:562–568
123. Arslanian S, Bacha F, Grey M, Marcus MD, White NH, Zeitler P. Evaluation and management of youth-onset type 2 diabetes: a position statement by the American Diabetes Association. *Diabetes Care* 2018;41:2648–2668
124. Wagenknecht LE, Lawrence JM, Isom S, et al.; SEARCH for Diabetes in Youth study. Trends in incidence of youth-onset type 1 and type 2 diabetes in the USA, 2002–18: results from the population-based SEARCH for Diabetes in Youth study. *Lancet Diabetes Endocrinol* 2023;11:242–250
125. Gebauer J, Higham C, Langer T, Denzer C, Brabant G. Long-term endocrine and metabolic consequences of cancer treatment: a systematic review. *Endocr Rev* 2019;40:711–767
126. Hwangbo Y, Lee EK. Acute hyperglycemia associated with anti-cancer medication. *Endocrinol Metab (Seoul)* 2017;32:23–29
127. Alenzi EO, Kelley GA. The association of hyperglycemia and diabetes mellitus and the risk of chemotherapy-induced neutropenia among cancer patients: a systematic review with meta-analysis. *J Diabetes Complications* 2017;31:267–272
128. Barone BB, Yeh H-C, Snyder CF, et al. Long-term all-cause mortality in cancer patients with diabetes. *J Clin Oncol* 2018;36:1234–1240
129. Perdigoto AL, Quandt Z, Anderson M, Herold KC. Checkpoint inhibitor-induced insulin-dependent diabetes: an emerging syndrome. *Lancet Diabetes Endocrinol* 2019;7:421–423
130. Akturk HK, Kahramangil D, Sarwal A, Hoffecker L, Murad MH, Michels AW. Immune checkpoint inhibitor-induced type 1 diabetes: a systematic review and meta-analysis. *Diabet Med* 2019;36:1075–1081
131. André F, Ciruelos E, Rubovszky G, et al.; SOLAR-1 Study Group. Alpelisib for PIK3CA-mutated, hormone receptor-positive advanced breast cancer. *N Engl J Med* 2019;380:1929–1940
132. Jhaveri KL, Im S-A, Saura C, et al. Overall survival with inavolisib in PIK3CA-mutated advanced breast cancer. *N Engl J Med* 2025;393:151–161
133. Rugo HS, André F, Yamashita T, et al. Time course and management of key adverse events during the randomized phase III SOLAR-1 study of PI3K inhibitor alpelisib plus fulvestrant in patients with HR-positive advanced breast cancer. *Ann Oncol* 2020;31:1001–1010
134. Xu KY, Shameem R, Wu S. Risk of hyperglycemia attributable to everolimus in cancer patients: a meta-analysis. *Acta Oncol* 2016;55:1196–1203
135. Spanjaard P, Petit JM, Schmitt A, Verges B, Bouillet B. Screening and management of metabolic complications of mTOR inhibitors in real-life settings. *Ann Endocrinol (Paris)* 2024;85:263–268
136. Ewald N, Bretzel RG. Diabetes mellitus secondary to pancreatic diseases (type 3c)—are we neglecting an important disease? *Eur J Intern Med* 2013;24:203–206
137. Hines OJ, Pandol SJ. Management of chronic pancreatitis. *BMJ* 2024;384:e070920
138. Woodmansey C, McGovern AP, McCullough KA, et al. Incidence, demographics, and clinical characteristics of diabetes of the exocrine pancreas (type 3c): a retrospective cohort study. *Diabetes Care* 2017;40:1486–1493
139. Makuc J. Management of pancreatogenic diabetes: challenges and solutions. *Diabetes Metab Syndr Obes* 2016;9:311–315
140. Andersen DK, Korc M, Petersen GM, et al. Diabetes, pancreatogenic diabetes, and pancreatic cancer. *Diabetes* 2017;66:1103–1110
141. Petrov MS, Basina M. Diagnosis of endocrine disease: diagnosing and classifying diabetes in diseases of the exocrine pancreas. *Eur J Endocrinol* 2021;184:R151–R163

142. Bellin MD, Gelrud A, Arreaza-Rubin G, et al. Total pancreatectomy with islet autotransplantation: summary of an NIDDK workshop. *Ann Surg* 2015;261:21–29
143. Quartuccio M, Hall E, Singh V, et al. Glycemic predictors of insulin independence after total pancreatectomy with islet autotransplantation. *J Clin Endocrinol Metab* 2017;102:801–809
144. Putman MS, Norris AW, Hull RL, et al. Cystic Fibrosis-Related Diabetes Workshop: research priorities spanning disease pathophysiology, diagnosis, and outcomes. *Diabetes Care* 2023;46:1112–1123
145. Ode KL, Ballman M, Battezzati A, et al. ISPAD Clinical Practice Consensus Guidelines 2022: management of cystic fibrosis-related diabetes in children and adolescents. *Pediatr Diabetes* 2022;23:1212–1228
146. Lewis C, Blackman SM, Nelson A, et al. Diabetes-related mortality in adults with cystic fibrosis. *Diabetes Care* 2023;46:1124–1130
- 2026 February 01 on guest by
http://diabetesjournals.org/care/article-pdf/49/Supplement_1/S27/848854/dc26s002.pdf Downloaded
- Diagnosis and Classification of Diabetes *Diabetes Care* Volume 49, Supplement 1, January 2026
- fibrosis. Role of genotype and sex. *Am J Respir Crit Care Med* 2015;191:194–200
163. Wallia A, Illuri V, Molitch ME. Diabetes care after transplant: definitions, risk factors, and clinical management. *Med Clin North Am* 2016;100:535–550
164. Kim HD, Chang J-Y, Chung BH, et al. Effect of everolimus with low-dose tacrolimus on development of new-onset diabetes after transplantation and allograft function in kidney transplantation: a multicenter, open-label, randomized trial. *Ann Transplant* 2021;26:e927984
165. Cheng C-Y, Chen C-H, Wu M-F, et al. Risk factors in and long-term survival of patients with post-transplantation diabetes mellitus: a retrospective cohort study. *Int J Environ Res Public Health* 2020;17:4581
166. Gulsoy Kirnap N, Bozkus Y, Haberal M. Analysis of risk factors for posttransplant diabetes mellitus after kidney transplantation: single-center experience. *Exp Clin Transplant* 2020;18:36–40
167. Galindo RJ, Fried M, Breen T, Tamler R. Hyperglycemia management in patients with posttransplantation diabetes. *Endocr Pract* 2016;22:454–465
168. Jenssen T, Hartmann A. Emerging treatments for post-transplantation diabetes mellitus. *Nat Rev Nephrol* 2015;11:465–477
169. Thomas MC, Mathew TH, Russ GR, Rao MM, Moran J. Early peri-operative glycaemic control and allograft rejection in patients with diabetes mellitus: a pilot study. *Transplantation* 2001;72:1321–1324
170. Greeley SAW, Polak M, Njølstad PR, et al. ISPAD Clinical Practice Consensus Guidelines 2022: the diagnosis and management of monogenic diabetes in children and adolescents. *Pediatr Diabetes* 2022;23:1188–1211
171. Carmody D, Stoy J, Greely Saw, Bell Gi, Philipson L. Chapter 2. A clinical guide to monogenic diabetes. In *Genetic diagnosis of endocrine disorders*, 2nd ed., Weiss RE, Refetoff S, Eds. Philadelphia, PA, Academic Press, 2016, pp. 21–30
172. De Franco E, Flanagan SE, Houghton JAL, et al. The effect of early, comprehensive genomic testing on clinical care in neonatal diabetes: an international cohort study. *Lancet* 2015;386:957–963
173. Sanyoura M, Letourneau L, Knight Johnson AE, et al. GCK-MODY in the US Monogenic Diabetes Registry: description of 27 unpublished variants. *Diabetes Res Clin Pract* 2019;151:231–236
174. Carmody D, Naylor RN, Bell CD, et al. GCK-MODY in the US National Monogenic Diabetes Registry: frequently misdiagnosed and unnecessarily treated. *Acta Diabetol* 2016;53:703–708
175. Timsit J, Saint-Martin C, Dubois-Laforgue D, Bellanné-Chantelot C. Searching for maturity-onset diabetes of the young (MODY): when and what for? *Can J Diabetes* 2016;40:455–461

176. Awa WL, Schober E, Wiegand S, et al. Reclassification of diabetes type in pediatric patients initially classified as type 2 diabetes mellitus: 15 years follow-up using routine data from the German/Austrian DPV database. *Diabetes Res Clin Pract* 2011;94:463–467
177. Shields BM, Hicks S, Shepherd MH, Colclough K, Hattersley AT, Ellard S. Maturity-onset diabetes of the young (MODY): how many cases are we missing? *Diabetologia* 2010;53:2504–2508
178. Chen Jovanovic L, Liang Y, Weng W, Hamilton M, Wintfeld N. Trends in the incidence of diabetes, its clinical sequelae, and associated costs in pregnancy. *Diabetes Metab Res Rev* 2015;31:707–716
179. Draznin B, Philipson LH, McGill JB. Atypical Diabetes: Pathophysiology, Clinical Presentations, and Treatment Options. Arlington, VA, American Diabetes Association Arlington,
180. Exeter Diabetes. MODY VA, 2018 Accessed 12 August 2025. Probability Calculator. Available from <https://www.diabetesgenes.org/exeter-diabetes-app/ModyCalculator>
181. Feig DS, Hwee J, Shah BR, Booth GL, Bierman AS, Lipscombe LL. Trends in incidence of diabetes in pregnancy and serious perinatal outcomes: a large, population-based study in Ontario, Canada, 1996–2010. *Diabetes Care* 2014;37:1590–1596
182. Peng TY, Ehrlich SF, Crites Y, et al. Trends and racial and ethnic disparities in the prevalence of pregestational type 1 and type 2 diabetes in Northern California: 1996–2014. *Am J Obstet Gynecol* 2017;216:177.e1–e177.e8
183. Wahabi HA, Fayed A, Esmaeil S, et al. Systematic review and meta-analysis of the effectiveness of pre-pregnancy care for women with diabetes for improving maternal and perinatal outcomes. *PLoS One* 2020;15:e0237571
184. Zhu W-W, Yang H-X, Wei Y-M, et al. Evaluation of the value of fasting plasma glucose in the first prenatal visit to diagnose gestational diabetes mellitus in China. *Diabetes Care* 2013;36:586–590
185. Mission JF, Catov J, Deihl TE, Feghali M, Scifres C. Early pregnancy diabetes screening and diagnosis: prevalence, rates of abnormal test results, and associated factors. *Obstet Gynecol* 2017;130:1136–1142
186. Cho NH, Shaw JE, Karuranga S, et al. IDF Diabetes Atlas: global estimates of diabetes prevalence for 2017 and projections for 2045. *Diabetes Res Clin Pract* 2018;138:271–281
187. Britton LE, Hussey JM, Crandell JL, Berry DC, Brooks JL, Bryant AG. Racial/ethnic disparities in diabetes diagnosis and glycemic control among women of reproductive age. *J Womens Health (Larchmt)* 2018;27:1271–1277
188. Robbins C, Boulet SL, Morgan I, et al. Disparities in preconception health indicators - Behavioral Risk Factor Surveillance System, 2013–2015, and Pregnancy Risk Assessment Monitoring System, 2013–2014. *MMWR Surveill Summ* 2018;67:1–16
189. Yuen L, Wong VW, Simmons D. Ethnic disparities in gestational diabetes. *Curr Diab Rep* 2018;18:68

2026 February 01 on guest by

http://diabetesjournals.org/care/article-pdf/49/Supplement_1/S27/848854/dc26s002.pdf Downloaded

diabetesjournals.org/care # Diagnosis and Classification of Diabetes

S49

1. Kattini R, Hummelen R, Kelly L. Early and adverse pregnancy outcomes. *N Engl J Med* diagnosing gestational diabetes mellitus. *NIH* 2008;358:1991–2002
2. Hughes RCE, Moore MP, Gullam JE, Mohamed K, Rowan J. An early pregnancy HbA1c $\geq 5.9\%$ (41 mmol/mol) is optimal for detecting diabetes and identifies women at increased risk of adverse pregnancy outcomes. *Diabetes Care* 2014;37:2953–2959
3. Shen L, Zhang S, Wen J, et al. Universal screening for hyperglycemia in early pregnancy and the risk of adverse pregnancy outcomes. *BMC Pregnancy Childbirth* 2025;25:203
4. Chen L, Pocobelli G, Yu O, et al. Early pregnancy hemoglobin A1C and pregnancy outcomes: a population-based study. *Am J Perinatol* 2019;36:1045–1053
5. Osmundson SS, Zhao BS, Kunz L, et al. First trimester hemoglobin A1c prediction of gestational diabetes. *Am J*

Perinatol 2016;33:977–982 6. McIntyre HD, Sacks DA, Barbour LA, et al. Issues with the diagnosis and classification of hyperglycemia in early pregnancy. *Diabetes Care* 2016;39:53–54 7. Cavagnoli G, Pimentel AL, Freitas PAC, Gross JL, Camargo JL. Factors affecting A1C in non-diabetic individuals: review and meta-analysis. *Clin Chim Acta* 2015;445:107–114 8. National Glycohemoglobin Standardization Program. Factors that interfere with HbA1c test results, 2024. Accessed 12 August 2025. Available from <https://ngsp.org/factors.asp> 9. Buchanan TA, Xiang A, Kjos SL, Watanabe R. What is gestational diabetes? *Diabetes Care* 2007;30(Suppl 2):S105–111 10. Noctor E, Crowe C, Carmody LA, et al.; ATLANTIC-DIP Investigators. Abnormal glucose tolerance post-gestational diabetes mellitus as defined by the International Association of Diabetes and Pregnancy Study Groups criteria. *Eur J Endocrinol* 2016;175:287–297 11. Kim C, Newton KM, Knopp RH. Gestational diabetes and the incidence of type 2 diabetes: a systematic review. *Diabetes Care* 2002;25:1862–1868 12. Liu Z, Zhang Q, Liu L, Liu W. Risk factors associated with early postpartum glucose intolerance in women with a history of gestational diabetes mellitus: a systematic review and meta-analysis. *Endocrine* 2023;82:498–512 13. Ratner RE, Christophi CA, Metzger BE, et al.; Diabetes Prevention Program Research Group. Prevention of diabetes in women with a history of gestational diabetes: effects of metformin and lifestyle interventions. *J Clin Endocrinol Metab* 2008;93:4774–4779 14. Aroda VR, Christophi CA, Edelstein SL, et al.; Diabetes Prevention Program Research Group. The effect of lifestyle intervention and metformin on preventing or delaying diabetes among women with and without gestational diabetes: the Diabetes Prevention Program outcomes study 10-year follow-up. *J Clin Endocrinol Metab* 2015;100:1646–1653 15. Vounzoulaki E, Khunti K, Abner SC, Tan BK, Davies MJ, Gillies CL. Progression to type 2 diabetes in women with a known history of gestational diabetes: systematic review and meta-analysis. *BMJ* 2020;369:m1361 16. Metzger BE, Lowe LP, Dyer AR, et al.; HAPO Study Cooperative Research Group. Hyperglycemia. *Diabetes Care* 2021;44:2197–2204

2026 February 01 on guest by

http://diabetesjournals.org/care/article-pdf/49/Supplement_1/S27/848854/dc26s002.pdf from Downloaded