

ISE 426

Optimization models and applications

Lecture 11 — October 2, 2014

- ▶ Basic feasible solutions
- ▶ simplex method
- ▶ sensitivity analysis

Reminders:


- ▶ Quiz on 10/14, practice on 10/09.

Simplex Method

Example

$$\begin{array}{llll} \text{maximize} & 3x_1 & + & 5x_2 \\ \text{subject to} & x_1 & & \leq 4 \\ & & 2x_2 & \leq 12 \\ & 3x_1 & + & 2x_2 \leq 18 \\ & x_1 & & \geq 0 \\ & & x_2 & \geq 0 \end{array}$$


The lines are the **constraint boundaries**.



Corner-Point Solutions

5 corner-point feasible (CPF) solutions:

1. (0,0)
2. (4,0)
3. (4,3)
4. (2,6)
5. (0,6)




The Key Idea

Taken together, the two properties mean we can find an optimal solution by:

1. Starting at any CPF solution
2. Moving to a better adjacent CPF solution, if one exists
3. Continuing until the current CPF solution has no adjacent CPF solutions that are better

This is the essence of the simplex method.

The Dual



$$\begin{array}{llllll}
 \text{minimize} & 4u_1 & + & 12u_2 & + & 18u_3 \\
 \text{subject to} & u_1 & & & + & 3u_3 & \geq & 3 \\
 & & & 2u_2 & + & 2u_3 & \geq & 5 \\
 & u_1 & & & & & \geq & 0 \\
 & & & u_2 & & & \geq & 0 \\
 & & & & & u_3 & \geq & 0
 \end{array}$$

Example

$$\begin{array}{llll} \max & 3x_1 & + & 5x_2 \\ \text{s.t.} & x_1 & & \leq 4 \\ & & 2x_2 & \leq 12 \\ & 3x_1 & + & 2x_2 \leq 18 \\ & x_1 & & \geq 0 \\ & & x_2 & \geq 0 \end{array}$$

$$\begin{array}{llllll} \min & 4u_1 & + & 12u_2 & + & 18u_3 \\ \text{s.t.} & u_1 & & & + & 3u_3 \geq 3 \\ & & & 2u_2 & + & 2u_3 \geq 5 \\ & u_1 & & & & \geq 0 \\ & & & u_2 & & \geq 0 \\ & & & & & u_3 \geq 0 \end{array}$$

Consider optimal $(x_1, x_2) = (2, 6)$, compute dual from complementary slackness:

$$u_1(x_1 - 4) = 0$$

$$u_2(+2x_2 - 12) = 0$$

$$u_3(3x_1 + 2x_2 - 18) = 0$$

$$x_1(u_1 + 3u_3 - 3) = 0$$

$$x_2(2u_2 + 2u_3 - 5) = 0$$

\Rightarrow

$$u_1(-2) = 0$$

$$u_2(0) = 0$$

$$u_3(0) = 0$$

$$2(u_1 + 3u_3 - 3) = 0$$

$$6(2u_2 + 2u_3 - 5) = 0$$

Example

$$\begin{array}{llll} \max & 3x_1 & + & 5x_2 \\ \text{s.t.} & x_1 & & \leq 4 \\ & & 2x_2 & \leq 12 \\ & 3x_1 & + & 2x_2 \leq 18 \\ & x_1 & & \geq 0 \\ & & x_2 & \geq 0 \end{array}$$

$$\begin{array}{llll} \min & 4u_1 & + & 12u_2 & + & 18u_3 \\ \text{s.t.} & u_1 & & & + & 3u_3 \geq 3 \\ & & 2u_2 & + & 2u_3 & \geq 5 \\ & u_1 & & & & \geq 0 \\ & & u_2 & & & \geq 0 \\ & & & & u_3 & \geq 0 \end{array}$$

Consider optimal $(x_1, x_2) = (2, 6)$, compute dual from complementary slackness:

$$\begin{array}{ll} u_1 = 0 & \\ 3u_3 = 3 & \\ 2u_2 + 2u_3 = 5 & \end{array} \quad \Rightarrow \quad \begin{array}{l} u_1 = 0 \\ u_2 = \frac{3}{2} \\ u_3 = 1 \end{array}$$

Example

$$\begin{array}{llll} \max & 3x_1 & + & 5x_2 \\ \text{s.t.} & x_1 & & \leq 4 \\ & & 2x_2 & \leq 12 \\ & 3x_1 & + & 2x_2 \leq 18 \\ & x_1 & & \geq 0 \\ & & x_2 & \geq 0 \end{array}$$

$$\begin{array}{llll} \min & 4u_1 & + & 12u_2 & + & 18u_3 \\ \text{s.t.} & u_1 & & & + & 3u_3 \geq 3 \\ & & 2u_2 & + & 2u_3 & \geq 5 \\ & u_1 & & & & \geq 0 \\ & & u_2 & & & \geq 0 \\ & & & u_3 & & \geq 0 \end{array}$$

Consider a **feasible** CPF $(x_1, x_2) = (4, 3)$, compute dual complementary solution:

$$u_1(x_1 - 4) = 0$$

$$u_2(+2x_2 - 12) = 0$$

$$u_3(3x_1 + 2x_2 - 18) = 0$$

$$x_1(u_1 + 3u_3 - 3) = 0$$

$$x_2(2u_2 + 2u_3 - 5) = 0$$

\Rightarrow

$$u_1(0) = 0$$

$$u_2(-4) = 0$$

$$u_3(0) = 0$$

$$2(u_1 + 3u_3 - 3) = 0$$

$$6(2u_2 + 2u_3 - 5) = 0$$

Example

$$\begin{array}{llll} \max & 3x_1 & + & 5x_2 \\ \text{s.t.} & x_1 & & \leq 4 \\ & & 2x_2 & \leq 12 \\ & 3x_1 & + & 2x_2 \leq 18 \\ & x_1 & & \geq 0 \\ & & x_2 & \geq 0 \end{array}$$

$$\begin{array}{llll} \min & 4u_1 & + & 12u_2 & + & 18u_3 \\ \text{s.t.} & u_1 & & & + & 3u_3 \geq 3 \\ & & 2u_2 & + & 2u_3 & \geq 5 \\ & u_1 & & & & \geq 0 \\ & & u_2 & & & \geq 0 \\ & & & & u_3 & \geq 0 \end{array}$$

Consider a **feasible** CPF $(x_1, x_2) = (4, 3)$, compute dual complementary solution:

$$\begin{array}{ll} u_2 = 0 \\ u_1 + 3u_3 = 3 \\ 2u_3 = 5 \end{array} \quad \Rightarrow \quad \begin{array}{l} u_1 = -\frac{9}{2} \\ u_2 = 0 \\ u_3 = \frac{5}{2} \end{array}$$

Example

$$\begin{array}{llll} \max & 3x_1 & + & 5x_2 \\ \text{s.t.} & x_1 & & \leq 4 \\ & & 2x_2 & \leq 12 \\ & 3x_1 & + & 2x_2 \leq 18 \\ & x_1 & & \geq 0 \\ & & x_2 & \geq 0 \end{array}$$

$$\begin{array}{llllll} \min & 4u_1 & + & 12u_2 & + & 18u_3 \\ \text{s.t.} & u_1 & & & + & 3u_3 \geq 3 \\ & & & 2u_2 & + & 2u_3 \geq 5 \\ & u_1 & & & & \geq 0 \\ & & & u_2 & & \geq 0 \\ & & & & & u_3 \geq 0 \end{array}$$

Consider a **feasible** CPF $(x_1, x_2) = (4, 3)$, compute dual complementary solution:

- ▶ $u_1 = -\frac{9}{2}$ is the **shadow price** for the constraint $x_1 \leq 4$ at the solution $x_1 = 4$.
- ▶ $u_1 < 0$ means that the objective function will improve if we allow $x < 4$ while we keep $3x_1 + 2x_2 = 18$.
- ▶ Consider $x_1 = 3$, then $x_2 = \frac{9}{2}$ and $3x_1 + 5x_2 = 3 * 4 + 5 * 3 + \frac{9}{2}$.

Example

$$\begin{array}{llll} \max & 3x_1 & + & 5x_2 \\ \text{s.t.} & x_1 & & \leq 4 \\ & & 2x_2 & \leq 12 \\ & 3x_1 & + & 2x_2 \leq 18 \\ & x_1 & & \geq 0 \\ & & x_2 & \geq 0 \end{array}$$

$$\begin{array}{llllll} \min & 4u_1 & + & 12u_2 & + & 18u_3 \\ \text{s.t.} & u_1 & & & + & 3u_3 \geq 3 \\ & & & 2u_2 & + & 2u_3 \geq 5 \\ & u_1 & & & & \geq 0 \\ & & u_2 & & & \geq 0 \\ & & & u_3 & & \geq 0 \end{array}$$

Consider a **feasible** CPF $(x_1, x_2) = (0, 6)$, compute dual complementary solution:

$$u_1(x_1 - 4) = 0$$

$$u_2(+2x_2 - 12) = 0$$

$$u_3(3x_1 + 2x_2 - 18) = 0$$

$$x_1(u_1 + 3u_3 - 3) = 0$$

$$x_2(2u_2 + 2u_3 - 5) = 0$$

\Rightarrow

$$u_1(-4) = 0$$

$$u_2(0) = 0$$

$$u_3(-6) = 0$$

$$0(u_1 + 3u_3 - 3) = 0$$

$$6(2u_2 + 2u_3 - 5) = 0$$

Example

$$\begin{array}{llll} \max & 3x_1 & + & 5x_2 \\ \text{s.t.} & x_1 & & \leq 4 \\ & & 2x_2 & \leq 12 \\ & 3x_1 & + & 2x_2 \leq 18 \\ & x_1 & & \geq 0 \\ & & x_2 & \geq 0 \end{array}$$

$$\begin{array}{llll} \min & 4u_1 & + & 12u_2 & + & 18u_3 \\ \text{s.t.} & u_1 & & & + & 3u_3 \geq 3 \\ & & 2u_2 & + & 2u_3 & \geq 5 \\ & u_1 & & & & \geq 0 \\ & & u_2 & & & \geq 0 \\ & & & & u_3 & \geq 0 \end{array}$$

Consider a **feasible** CPF $(x_1, x_2) = (0, 6)$, compute dual complementary solution:

$$\begin{aligned} u_1 &= 0 \\ 2u_2 &= 2 \\ u_3 &= 0 \end{aligned}$$

$$\begin{aligned} u_1 &= 0 \\ \Rightarrow u_2 &= \frac{5}{2} \\ u_3 &= 0 \\ u_1 + 3u_3 - 3 &= -3 \end{aligned}$$

Example

$$\begin{array}{llll} \max & 3x_1 & + & 5x_2 \\ \text{s.t.} & x_1 & & \leq 4 \\ & & 2x_2 & \leq 12 \\ & 3x_1 & + & 2x_2 \leq 18 \\ & x_1 & & \geq 0 \\ & & x_2 & \geq 0 \end{array}$$

$$\begin{array}{llllll} \min & 4u_1 & + & 12u_2 & + & 18u_3 \\ \text{s.t.} & u_1 & & & + & 3u_3 \geq 3 \\ & & & 2u_2 & + & 2u_3 \geq 5 \\ & u_1 & & & & \geq 0 \\ & & u_2 & & & \geq 0 \\ & & & u_3 & & \geq 0 \end{array}$$

Consider a **feasible** CPF $(x_1, x_2) = (0, 6)$, compute dual complementary solution:

- ▶ $s_1 = u_1 + 3u_3 - 3 = -3$ is the **reduced cost** for the variable $x_1 \geq 0$ at the solution $x_1 = 0$.
- ▶ $s_1 < 0$ means that the objective function will improve if we allow $x > 0$ while we keep $2x_2 = 12$.
- ▶ Consider $x_1 = 1$, then $x_2 = 6$ then $3x_1 + 5x_2 = 3 * 0 + 5 * 6 + 3$.

Changes in Objective Function Coefficients

- ▶ Suppose x^* is the optimal solution for an LP.
- ▶ Z^* is its optimal objective value.
- ▶ Suppose that some objective function coefficient c_j changes.

$$\begin{array}{ll}\max Z = & 3x_1 + 5x_2 \\ \text{s.t.} & x_1 \leq 4 \\ & 2x_2 \leq 12 \\ & 3x_1 + 2x_2 \leq 18 \\ & x_1, x_2 \geq 0\end{array}$$

- ▶ What if the 3 changed?

Changes in c , cont'd

$$\begin{array}{ll}\max Z = & 3x_1 + 5x_2 \\ \text{s.t.} & x_1 \leq 4 \\ & 2x_2 \leq 12 \\ & 3x_1 + 2x_2 \leq 18 \\ & x_1, x_2 \geq 0\end{array}$$

- ▶ If 3 changed to 0, what would the new solution be?
- ▶ If 3 changed to 30, what would the new solution be?
- ▶ If the 3 changed to $3 \pm \delta$, where δ is tiny, what would the new solution be?

Example: Maximization

$$\begin{array}{llll} \max Z = & 3x_1 & + & 5x_2 \\ \text{s.t.} & x_1 & & \leq 4 \\ & & 2x_2 & \leq 12 \\ & 3x_1 + 2x_2 & \leq & 18 \\ & x_1, & x_2 & \geq 0 \end{array}$$

- ▶ Optimal solution is $(x_1^*, x_2^*) = (2, 6)$, $Z^* = 36$.
- ▶ Suppose the 3 increased to 7.
- ▶ Which of the following is true?
 1. Z^* will increase.
 2. Z^* will decrease.
 3. Z^* will stay the same.
 4. We can't say.

Example: Maximization, cont'd

$$\begin{array}{lll} \max Z = & 3x_1 + 5x_2 \\ \text{s.t.} & x_1 & \leq 4 \\ & & 2x_2 \leq 12 \\ & 3x_1 + 2x_2 & \leq 18 \\ & x_1, x_2 & \geq 0 \end{array}$$

- ▶ $(x_1^*, x_2^*) = (2, 6)$, $Z^* = 36$.
- ▶ **3** increases to **7**.
- ▶ The local rate of increase is derived from $x_1^* \delta = 2 \times 4 = 8$.
- ▶ Which of the following is true?
 1. Z^* will increase by *exactly* 8.
 2. Z^* will increase by *at most* 8.
 3. Z^* will increase by *at least* 8.
 4. Z^* will increase, but we don't know by how much.

Changes in Constraint Right-Hand Sides

$$\begin{array}{llll} \max Z = & 3x_1 & + & 5x_2 \\ \text{s.t.} & x_1 & & \leq 4 \\ & & 2x_2 & \leq 12 \\ & 3x_1 & + & 2x_2 \leq 18 \\ & x_1 & , & x_2 \geq 0 \end{array}$$

- ▶ Suppose 12 increased.
- ▶ Would optimal solution change?
- ▶ Would optimal objective value change?
- ▶ Would optimal basis change?

Example





$$\begin{array}{llll} \max Z = & 3x_1 + 5x_2 & & \\ \text{s.t.} & x_1 & \leq & 4 \\ & & 2x_2 \leq & \textcolor{red}{12} \\ & 3x_1 + 2x_2 \leq & 18 & \\ & x_1, x_2 \geq & 0 & \end{array}$$

- ▶ Optimal solutions are $(x_1^*, x_2^*) = (2, 6)$,
 $(u_1^*, u_2^*, u_3^*) = (0, \frac{3}{2}, 1)$, $Z^* = 36$.
- ▶ Suppose the $\textcolor{red}{12}$ increased to $\textcolor{red}{16}$.
- ▶ Which of the following is true?
 1. Z^* will increase.
 2. Z^* will decrease.
 3. Z^* will stay the same.
 4. We can't say.

Example, cont'd

$$\begin{array}{llll} \max Z = & 3x_1 + 5x_2 & & \\ \text{s.t.} & x_1 & \leq & 4 \\ & & 2x_2 \leq & \textcolor{red}{12} \\ & 3x_1 + 2x_2 \leq & 18 & \\ & x_1, x_2 \geq & 0 & \end{array}$$

- ▶ $(x_1^*, x_2^*) = (2, 6)$, $(u_1^*, u_2^*, u_3^*) = (0, \frac{3}{2}, 1)$, $Z^* = 36$.
- ▶ $\textcolor{red}{12}$ increases to $\textcolor{red}{16}$.
- ▶ The local rate of increase come from the shadow price:
 $u_2^* \delta = \frac{3}{2} \times 4 = 6$.
- ▶ Which of the following is true?
 1. Z^* will increase by *exactly* 6.
 2. Z^* will increase by *at most* 6.
 3. Z^* will increase by *at least* 6.
 4. Z^* will increase, but we don't know by how much.

	Change c	Change b
Maximization	 <p>A graph showing the optimal value Z^* on the vertical axis versus the cost coefficient c_j on the horizontal axis. The curve starts at a constant value for small changes in c_j, then begins to rise with an increasing slope, indicating that the objective value increases at an increasing rate as c_j increases beyond a certain point.</p>	 <p>A graph showing the optimal value Z^* on the vertical axis versus the right-hand side value b_i on the horizontal axis. The curve starts at the origin and increases with a decreasing slope, eventually leveling off to a horizontal line, indicating that the objective value approaches a maximum as b_i increases.</p>
Minimization	 <p>A graph showing the optimal value Z^* on the vertical axis versus the cost coefficient c_j on the horizontal axis. The curve starts at a low value and increases with a decreasing slope, eventually leveling off to a horizontal line, indicating that the objective value approaches a minimum as c_j increases.</p>	 <p>A graph showing the optimal value Z^* on the vertical axis versus the right-hand side value b_i on the horizontal axis. The curve starts at a constant value for small changes in b_i, then begins to rise with an increasing slope, indicating that the objective value increases at an increasing rate as b_i increases beyond a certain point.</p>

Relationship to Complementary Slackness

- ▶ Suppose there is slack in the i th primal constraint.
 - ▶ Increasing the RHS would not change the optimal solution.
 - ▶ By complementary slackness, u_i^* must equal 0 (in the dual).
 - ▶ Using the statement on the previous slides, the optimal objective function changes by u_i^* , or 0.
- ▶ Suppose there is no slack in the i th primal constraint.
 - ▶ Increasing the RHS *would* change the optimal solution.
 - ▶ u_i^* *probably* (!) is greater than 0.
 - ▶ Using the statement above, the optimal objective function changes by u_i^* .
- ▶ This agrees with our interpretation of the dual values as *shadow prices*.

Relationship to Complementary Slackness, cont'd

- ▶ Now suppose there is slack in the j th dual constraint.
 - ▶ By complementary slackness, $x_j^* = 0$ (in the primal).
 - ▶ If we increase c_j slightly, we'll still want to set $x_j^* = 0$.
 - ▶ We argued that for each unit increase in c_j , Z^* changes by x_j^* (if optimal basis stays the same).
 - ▶ So Z^* increases by 0 when c_j increases.
- ▶ Suppose there is no slack in the j th dual constraint (reduced cost is 0).
 - ▶ $x_j^* > 0$ (probably).
 - ▶ If we increase c_j by 1, the objective value will go up by x_j^* .