

ISE 426

Optimization models and applications


Lecture 2 — September 1, 2015

Convexity; Relaxations; Lower and upper bounds.

- ▶ Winston, chapter 1, **or**
- ▶ Winston & Venkataramanan, chapter 1, **or**
- ▶ Hillier & Lieberman, chapter 2.


Convexity

Convex sets



Def.: A set $S \subseteq \mathbb{R}^n$ is **convex** if any two points x' and x'' of S are joined by a segment **entirely** contained in S :


Convex sets



Def.: A set $S \subseteq \mathbb{R}^n$ is **convex** if any two points x' and x'' of S are joined by a segment **entirely** contained in S :

$$\forall x', x'' \in S, \alpha \in [0, 1], \quad \alpha x' + (1 - \alpha)x'' \in S$$

Convex sets




Def.: A set $S \subseteq \mathbb{R}^n$ is **convex** if any two points x' and x'' of S are joined by a segment **entirely** contained in S :


$$\forall x', x'' \in S, \alpha \in [0, 1], \quad \alpha x' + (1 - \alpha)x'' \in S$$

The intersection of two convex sets is convex.


Definition




Definition




Definition




Definition



Definition



Definition



Convex functions

Def.: A function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is **convex** if, for any two points x' and $x'' \in \mathbb{R}^n$ and for any $\alpha \in [0, 1]$

$$f(\alpha x' + (1 - \alpha)x'') \leq \alpha f(x') + (1 - \alpha)f(x'')$$

Convex functions

Def.: A function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is **convex** if, for any two points x' and $x'' \in \mathbb{R}^n$ and for any $\alpha \in [0, 1]$

$$f(\alpha x' + (1 - \alpha)x'') \leq \alpha f(x') + (1 - \alpha)f(x'')$$

- The **sum** of convex functions is a convex function

Convex functions

Def.: A function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is **convex** if, for any two points x' and $x'' \in \mathbb{R}^n$ and for any $\alpha \in [0, 1]$

$$f(\alpha x' + (1 - \alpha)x'') \leq \alpha f(x') + (1 - \alpha)f(x'')$$

- ▶ The **sum** of convex functions is a convex function
- ▶ Multiplying a convex function by a positive scalar gives a convex function


Convex functions

Def.: A function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is **convex** if, for any two points x' and $x'' \in \mathbb{R}^n$ and for any $\alpha \in [0, 1]$


$$f(\alpha x' + (1 - \alpha)x'') \leq \alpha f(x') + (1 - \alpha)f(x'')$$

- ▶ The **sum** of convex functions is a convex function
- ▶ Multiplying a convex function by a positive scalar gives a convex function
- ▶ **linear** functions $\sum_{i=1}^k a_i x_i$ are convex, irrespective of the sign of a_i 's.


Definition




Definition



Definition



Definition



Local and global optima

A vector $x^l \in \mathbb{R}^n$ is a **local** optimum if

- ▶ there is a *neighbourhood*¹ N of x^l with no better point than x^l :

$$\forall x \in N, f_0(x) \geq f_0(x^l)$$


A vector $x^g \in \mathbb{R}^n$ is a **global** optimum if

- ▶ x^g
- ▶ there is no x better than x^g , i.e.,

$$f_0(x) \geq f_0(x^g) \quad \forall x$$

¹A *neighbourhood* of x^l can be defined as $N = \{x : \|x - x^l\|_2 \leq \epsilon\}$ for some ϵ .

Local optima, global optima



Positive Semidefinite Matrices

A square $n \times n$ matrix A is **Positive Semidefinite** (PSD) (denoted with $A \succeq 0$) if, for any n -vector, the following holds:

$$x^{\top} A x \geq 0$$

$$\sum_{i=1}^n \sum_{j=1}^n a_{ij} x_i x_j =$$

Positive Semidefinite Matrices

A square $n \times n$ matrix A is **Positive Semidefinite** (PSD) (denoted with $A \succeq 0$) if, for any n -vector, the following holds:

$$x^\top A x \geq 0$$

$$\begin{aligned} \sum_{i=1}^n \sum_{j=1}^n a_{ij} x_i x_j = & \\ a_{11}x_1^2 & + a_{12}x_1x_2 + \dots + a_{1n}x_1x_n + \\ a_{21}x_2x_1 & + a_{22}x_2^2 + \dots + a_{2n}x_2x_n + \\ \vdots & \vdots \quad \ddots \quad \vdots \\ a_{n1}x_nx_1 & + a_{n2}x_nx_1 + \dots + a_{nn}x_n^2 \\ & \geq 0 \end{aligned}$$

Some linear algebra

- ▶ A **minor** of a $m \times n$ matrix A is the determinant of a (square) submatrix of A obtained by removing some rows and columns of A .
- ▶ For a square matrix B , a **principal minor** of B is obtained by removing the same row and column indices from B

For example, $A = \begin{pmatrix} 1 & 2 & 3 & 4 \\ 2 & 5 & 6 & 7 \\ 3 & 6 & 8 & 9 \\ 4 & 7 & 9 & 0 \end{pmatrix}$.

Principal minors: (1) , (8) , $\begin{pmatrix} 1 & 2 \\ 2 & 5 \end{pmatrix}$, $\begin{pmatrix} 5 & 7 \\ 7 & 0 \end{pmatrix}$, $\begin{pmatrix} 1 & 2 & 4 \\ 2 & 5 & 7 \\ 4 & 7 & 0 \end{pmatrix}$

Positive (Semi)Definite Hessian Matrices

A symmetric matrix $A \in R^{n \times n}$ is **Positive Semidefinite** (PSD) (denote it with $A \succeq 0$) if all principal minors of A are nonnegative.

A symmetric matrix $A \in R^{n \times n}$ is **Positive Semidefinite** (PSD) if and only if $A = BB^T$ for some $B \in R^{n \times n}$.

A symmetric matrix $A \in R^{n \times n}$ is **Positive Semidefinite** (PSD) if for all $i = 1, \dots, n$ $a_{ii} \geq \sum_{j=1, j \neq i}^n |a_{ij}|$.

A symmetric matrix $A \in R^{n \times n}$ is **Positive Semidefinite** (PSD) if all its eigenvalues are nonnegative.

Positive (Semi)Definite Hessian Matrices

Given a function $f : \mathbb{R}^n \rightarrow \mathbb{R}$, consider its **Hessian**:

$$H_f(x) = \begin{pmatrix} \frac{\partial^2 f}{\partial x_1^2} & \frac{\partial^2 f}{\partial x_1 \partial x_2} & \cdots & \frac{\partial^2 f}{\partial x_1 \partial x_n} \\ \frac{\partial^2 f}{\partial x_2 \partial x_1} & \frac{\partial^2 f}{\partial x_2^2} & \cdots & \frac{\partial^2 f}{\partial x_2 \partial x_n} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial^2 f}{\partial x_n \partial x_1} & \frac{\partial^2 f}{\partial x_n \partial x_2} & \cdots & \frac{\partial^2 f}{\partial x_n^2} \end{pmatrix}$$

A function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is **convex** on set $\Omega \in \mathbb{R}^n$ if and only if the Hessian $H_f(x) \succeq 0$ for all $x \in \Omega$.

Examples

- ▶ The function $f(x) = x$ is **convex**
- ▶ The function $f(x_1, x_2) = x_1 + x_2$ is **convex**
- ▶ The function $f(x_1, x_2) = x_1^2 + x_2$ is **convex**
- ▶ The function $f(x_1, x_2) = 5x_1^2 + 3x_2^2$ is **convex**
- ▶ The function $f(x_1, x_2) = x_1^2 + x_2^2 - x_1x_2$ is **convex**

Examples

- ▶ The function $f(x) = x$ is **convex**
- ▶ The function $f(x_1, x_2) = x_1 + x_2$ is **convex**
- ▶ The function $f(x_1, x_2) = x_1^2 + x_2$ is **convex**
- ▶ The function $f(x_1, x_2) = 5x_1^2 + 3x_2^2$ is **convex**
- ▶ The function $f(x_1, x_2) = x_1^2 + x_2^2 - x_1x_2$ is **convex**

- ▶ The function $f(x_1, x_2) = x_1^2 + x_2^2 + 5x_1x_2$ is **nonconvex**
- ▶ The function $f(x_1, x_2) = x_1^2 - x_2^2$ is **nonconvex**
- ▶ The function $f(x_1, x_2) = x_1x_2$ is **nonconvex**
- ▶ The function $f(x) = \sin x$, for $x \in [0, 2\pi]$ is **nonconvex**
- ▶ The function $f(x) = -x^2$ is **nonconvex**


Convex constraints

- ▶ A constraint $g(x) \leq b$, with $g : \mathbb{R}^n \rightarrow \mathbb{R}$, defines a subset S of \mathbb{R}^n , that is,


$$S = \{x \in \mathbb{R}^n : g(x) \leq b\}$$

- ▶ The constraint $g(x) \leq b$ is **convex** if the set S is convex.
- ▶ If the *function* $g(x)$ is convex, the *constraint* $g(x) \leq b$ is convex.

Convex constraints



Convex constraints



Note: if the function $g(x)$ is convex, the constraint $g(x) \geq b$ may be nonconvex!

Convex and concave functions

Def.: A function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is **concave** if $-f(x)$ is convex.

- concave functions are useful with **maximization** problems:

$$\begin{aligned} \max \quad & f(x) \\ & g_i(x) \leq 0 \quad \forall i = 1, 2, \dots, m \\ & h_i(x) = 0 \quad \forall i = m + 1, m + 2, \dots, m + q \\ & x \in \mathbb{R}^n \end{aligned}$$

is a convex problem if $f(x)$ is **concave**, all $g_i(x)$ are convex, and all $h_i(x)$ are affine.

- concave functions are also useful for \geq constraints:

$$g_i(x) \geq 0$$

is a convex constraint if $g_i(x)$ is concave.

For what convex $g(x)$ the *constraint* $g(x) \geq b$ is always convex?

For what convex $g(x)$ the *constraint* $g(x) \geq b$ is always convex?

- ▶ linear constraints $\sum_{i=1}^k a_i x_i \begin{cases} \leq \\ = \\ \geq \end{cases} b$ are convex

The general optimization problem

Consider a vector $x \in \mathbb{R}^n$ of variables.

An optimization problem can be expressed as:

$$\begin{array}{ll} \mathbf{P} : & \text{minimize } f_0(x) \\ & \text{s.t. } f_1(x) \leq b_1 \\ & \quad f_2(x) \leq b_2 \\ & \quad \vdots \\ & \quad f_m(x) \leq b_m \end{array}$$

Feasible solutions, local and global optima

Define $F = \{x \in \mathbb{R}^n : f_1(x) \leq b_1, f_2(x) \leq b_2, \dots, f_m(x) \leq b_m\}$, that is, F is the **feasible set** of an optimization problem.

All points $x \in F$ are called **feasible solutions**.

A vector $x^l \in \mathbb{R}^n$ is a **local** optimum if

- ▶ $x^l \in F$
- ▶ there is a *neighbourhood*² N of x^l with no better point than x^l :

$$\forall x \in N \cap F, f_0(x) \geq f_0(x^l)$$

A vector $x^g \in \mathbb{R}^n$ is a **global** optimum if

- ▶ $x^g \in F$
- ▶ there is no $x \in F$ better than x^g , i.e.,

$$f_0(x) \geq f_0(x^g) \quad \forall x \in F$$

²A *neighbourhood* of x^l can be defined as $N = \{x : \|x - x^l\|_2 \leq \epsilon\}$ for some ϵ .

Convex problems

Def.: An optimization problem is **convex** if

- ▶ the objective function is convex
- ▶ **all** constraints are convex

Convex problems

Def.: An optimization problem is **convex** if

- ▶ the objective function is convex
- ▶ **all** constraints are convex

Convex optimization problems are *easy*:

If a problem P is convex, a local optimum x^* of P is also a global optimum of P.
--

(Hint) When modeling an optimization problem, it would be good if we found a convex problem.

Examples of convex problems

$$\begin{array}{ll}\mathbf{P} : & \min \quad x_1^2 + 2x_2^2 \\ & \text{s.t.} \quad x_1^2 + x_2^2 \leq 1 \\ & \quad \quad 0 \leq x_1 \leq 2 \\ & \quad \quad 1 \leq x_2 \leq 5\end{array}$$

Examples of nonconvex problems

$$\begin{array}{ll}\mathbf{P} : & \min \quad x_1^2 + 2x_2^2 \\ & \text{s.t.} \quad x_1^2 + x_2^2 = 5 \\ & \quad \quad 0 \leq x_1 \leq 2 \\ & \quad \quad 1 \leq x_2 \leq 5\end{array}$$

Examples of nonconvex problems

$$\begin{array}{ll}\mathbf{P} : & \min \quad x_1^2 + 2x_2^2 \\ & \text{s.t.} \quad x_1^2 + x_2^2 \leq 5 \\ & \quad \quad 0 \leq x_1 \leq 2 \\ & \quad \quad 1 \leq x_2 \leq 5 \\ & \quad \quad x_2 \in \mathbb{Z}\end{array}$$

Examples of "hidden" convex problems

$$\begin{array}{ll}\mathbf{P} : & \min \quad x_1 - 2x_2^2 \\ & \text{s.t.} \quad x_1^2 + x_2^2 \leq 1 \\ & \quad \quad x_2 = 0 \\ & \quad \quad 0 \leq x_1 \leq 5\end{array}$$

Additional examples of convex problems and functions

$$\begin{array}{ll}\mathbf{P} : & \min \quad |x_1| + |x_2| \\ & \text{s.t.} \quad x_1^2 + x_2^2 \leq 1 \\ & \quad \quad |2x_1 + 3x_2| \leq 10 \\ & \quad \quad 0 \leq x_1 \leq 5\end{array}$$

Your first Optimization model

Variables	r : radius of the can's base h : height of the can
Objective	$2\pi rh + 2\pi r^2$ (minimize)
Constraints	$\pi r^2 h = V$ $h > 0$ $r > 0$

Relaxations

Relaxation of an Optimization problem

Consider an optimization problem

$$\begin{array}{ll}\mathbf{P} : & \min f_0(x) \\ & \text{s.t. } f_1(x) \leq b_1 \\ & \quad f_2(x) \leq b_2 \\ & \quad \vdots \\ & \quad f_m(x) \leq b_m,\end{array}$$

Let F denote the set of points x that satisfy all constraints:

$$F = \{x \in \mathbb{R}^n : \begin{array}{l} f_1(x) \leq b_1, \\ f_2(x) \leq b_2, \\ \vdots \\ f_m(x) \leq b_m \end{array}\}$$

So we can write $\mathbf{P} : \min\{f_0(x) : x \in F\}$ for short.

Relaxation of an Optimization problem

Consider a problem $\mathbf{P} : \min\{f_0(x) : x \in F\}$.

A problem $\mathbf{P}' : \min\{f'_0(x) : x \in F'\}$ is a **relaxation** of \mathbf{P} if:

- ▶ $F' \supseteq F$
- ▶ $f'_0(x) \leq f_0(x)$ for all $x \in F$.³


If \mathbf{P}' is a relaxation of a problem \mathbf{P} , then the global optimum of \mathbf{P}' is \leq the global optimum of \mathbf{P} .

³We don't care what $f'_0(x)$ is outside of F .


Examples

- ▶ $\min\{f(x) : -1 \leq x \leq 1\}$ is a relaxation of $\min\{f(x) : x = 0\}$
- ▶ $\min\{f(x) : -1 \leq x \leq 1\}$ is a r. of $\min\{f(x) : 0 \leq x \leq 1\}$
- ▶ $\min\{f(x) : -1 \leq x \leq 1\}$ is **not** a r. of $\min\{f(x) : -2 \leq x \leq 1\}$
- ▶ $\min\{f(x) : g(x) \leq b\}$ is a r. of $\min\{f(x) : g(x) \leq b - 1\}$
- ▶ $\min\{f(x) - 1 : g(x) \leq b\}$ is a r. of $\min\{f(x) : g(x) \leq b\}$


Relaxation of an Optimization problem




Relaxation of an Optimization problem




Relaxation of an Optimization problem




Relaxation of an Optimization problem



Relaxation of an Optimization problem



Relaxation of an Optimization problem



Relaxation of an Optimization problem

Consider again a problem

$\mathbf{P} : \min\{f_0(x) : f_1(x) \leq b_1, f_2(x) \leq b_1, \dots, f_m(x) \leq b_m\}$, or

$\mathbf{P} : \min\{f_0(x) : x \in F\}$ for short.

- ▶ **deleting** a constraint from \mathbf{P} provides a relaxation of \mathbf{P} .
- ▶ **adding** a constraint $f_{m+1}(x) \leq b_{m+1}$ to a problem \mathbf{P} does the opposite:

$$F'' = \{x \in \mathbb{R}^n : \begin{array}{l} f_1(x) \leq b_1, \\ f_2(x) \leq b_2, \\ \dots, \\ f_m(x) \leq b_m, \\ f_{m+1}(x) \leq b_{m+1} \end{array}\} \subseteq F$$

and therefore

$$\min\{f_0(x) : x \in F''\} \geq \min\{f_0(x) : x \in F\}$$

Upper & Lower bounds

Lower and upper bounds

Consider an optimization problem $\mathbf{P} : \min\{f_0(x) : x \in F\}$:

- ▶ for any feasible solution $x \in F$, the corresponding objective function value $f_0(x)$ is an **upper bound**.
- ▶ the most interesting upper bounds are the **local optima**.
- ▶ a **lower bound** of \mathbf{P} is instead a value z such that

$$z \leq \min\{f_0(x) : x \in F\}.$$

Upper vs. Lower bounds

Situation #1:

You: “We found a solution that will only cost 372,000 \$.”

Boss: “Ok, that sounds good.”

Upper vs. Lower bounds

Situation #1:

You: “We found a solution that will only cost 372,000 \$.”

Boss: “Ok, that sounds good.”

Situation #2:

You: “We found a solution that will only cost 372,000 \$.”

Boss: “That’s too much, find something better.”

Upper vs. Lower bounds

Situation #1:

You: “We found a solution that will only cost 372,000 \$.”

Boss: “Ok, that sounds good.”

Situation #2:

You: “We found a solution that will only cost 372,000 \$.”

Boss: “That’s too much, find something better.”

...

You: “We found another solution that costs 354,000 \$.”

Boss: “Can’t you do better than that?”

Upper vs. Lower bounds

Situation #1:

You: "We found a solution that will only cost 372,000 \$."

Boss: "Ok, that sounds good."

Situation #2:

You: "We found a solution that will only cost 372,000 \$."

Boss: "That's too much, find something better."

...

You: "We found another solution that costs 354,000 \$."

Boss: "Can't you do better than that?"

You: "I can try again, but here's the proof that we can't go below 351,500."

Boss: "Ok then, that's a good solution."

What relaxations are for

- ▶ If \mathbf{P}' is a relaxation of a problem \mathbf{P} , then the global optimum of \mathbf{P}' is \leq the global optimum of \mathbf{P} .
 - ▶ Hence, any relaxation \mathbf{P}' of \mathbf{P} provides a **lower bound** on \mathbf{P} .
- \Rightarrow If a problem \mathbf{P} is difficult but a relaxation \mathbf{P}' of \mathbf{P} is easier to solve than \mathbf{P} itself, we can still try and solve \mathbf{P}' : (i) we get a lower bound and (ii) the solution of \mathbf{P}' may help solve \mathbf{P} .

To recap: the Knapsack problem

At a flea market in Rome, you spot n objects (old pictures, a vessel, rusty medals. . .) that you could re-sell in your antique shop for about double the price.

- ▶ You want these objects to pay for your flight ticket to Rome, which cost C .
- ▶ Also, you don't want a heavy backpack, so you want to buy the objects that will load it as little as possible.

How do you solve this problem?

The Knapsack problem

Each object $i = 1, 2, \dots, n$ has a price p_i and a weight w_i .

- ▶ Variables: one variable x_i for each $i = 1, 2, \dots, n$. This is a “yes/no” variable: either you take the i -th object or not.

The Knapsack problem

Each object $i = 1, 2, \dots, n$ has a price p_i and a weight w_i .

- ▶ Variables: one variable x_i for each $i = 1, 2, \dots, n$. This is a “yes/no” variable: either you take the i -th object or not.
- ▶ Constraint: total revenue must be at least C

The Knapsack problem

Each object $i = 1, 2, \dots, n$ has a price p_i and a weight w_i .

- ▶ Variables: one variable x_i for each $i = 1, 2, \dots, n$. This is a “yes/no” variable: either you take the i -th object or not.
- ▶ Constraint: total revenue must be at least C
(As you'll double the price when selling them at your store, the revenue for each object is exactly p_i)

The Knapsack problem

Each object $i = 1, 2, \dots, n$ has a price p_i and a weight w_i .

- ▶ Variables: one variable x_i for each $i = 1, 2, \dots, n$. This is a “yes/no” variable: either you take the i -th object or not.
- ▶ Constraint: total revenue must be at least C
(As you'll double the price when selling them at your store, the revenue for each object is exactly p_i)
- ▶ Objective function: the total weight

Your first (non-trivial) optimization model

$$\begin{array}{ll}\min & \sum_{i=1}^n w_i x_i \\ & \sum_{i=1}^n p_i x_i \geq C \\ & x_i \in \{0, 1\} \quad \forall i = 1, 2, \dots, n\end{array}$$

Is it convex?

Relaxing the Knapsack problem

$$\begin{array}{ll} \min & \sum_{i=1}^n w_i x_i \\ & x_i \in \{0, 1\} \quad \forall i = 1, 2, \dots, n \end{array}$$

Relaxing the Knapsack problem

$$\begin{array}{ll} \min & \sum_{i=1}^n w_i x_i \\ & x_i \in \{0, 1\} \quad \forall i = 1, 2, \dots, n \end{array}$$

This relaxation would give us $x_i = 0$ for all $i = 1, 2, \dots, n$, and a lower bound of $\sum_{i=1}^n w_i x_i = 0$. Not so great...

Relaxing the Knapsack problem

$$\begin{aligned} \min \quad & \sum_{i=1}^n w_i x_i \\ & x_i \in \{0, 1\} \quad \forall i = 1, 2, \dots, n \end{aligned}$$

This relaxation would give us $x_i = 0$ for all $i = 1, 2, \dots, n$, and a lower bound of $\sum_{i=1}^n w_i x_i = 0$. Not so great...

$$\begin{aligned} \min \quad & \sum_{i=1}^n w_i x_i \\ & \sum_{i=1}^n p_i x_i \geq C \\ & 0 \leq x_i \leq 1 \quad \forall i = 1, 2, \dots, n \end{aligned}$$

Relaxing the Knapsack problem

$$\begin{aligned} \min \quad & \sum_{i=1}^n w_i x_i \\ & x_i \in \{0, 1\} \quad \forall i = 1, 2, \dots, n \end{aligned}$$

This relaxation would give us $x_i = 0$ for all $i = 1, 2, \dots, n$, and a lower bound of $\sum_{i=1}^n w_i x_i = 0$. Not so great...

$$\begin{aligned} \min \quad & \sum_{i=1}^n w_i x_i \\ & \sum_{i=1}^n p_i x_i \geq C \\ & 0 \leq x_i \leq 1 \quad \forall i = 1, 2, \dots, n \end{aligned}$$

- ▶ Relaxing integrality of the variables gives a relaxation where we admit **fractions** of objects.
- ≈ we pulverized all objects and took some spoonfuls of each
- ▶ It doesn't make sense, but it's a relaxation, and it **does** give us a better lower bound.