

Optimization of Graph Squares - Proofs

Eddie Xu

`eddie.xu@student.unimelb.com.au`

July 28, 2026

Contents

1	Formulations	1
1.1	Integer Program	2
1.2	Linear Program	3
1.3	Max Linear Program	4
2	Two-Edge-Triangle Proof	4
2.1	Integer Program	4
2.2	Linear Program	5
2.3	Max Linear Program	5

1 Formulations

Let known constants $g_{ij} \in \{0, 1\}$ come from the adjacency matrix of the square A_G and let $M \in \mathbb{Z}_{>0}$ be a sufficiently large constant. The variables in the program are x_{ij} , y_{ikj} , z_{ij} where x_{ij} form the ij -th entry of the adjacency matrix of the root A_H .

1.1 Integer Program

Variables:

$$x_{ij} \in \mathbb{Z}_{\geq 0}, \quad \forall i, j \in \{1, \dots, n\}, \quad (1)$$

$$y_{ikj} \in \mathbb{Z}_{\geq 0}, \quad \forall i, k, j \in \{1, \dots, n\}, \quad (2)$$

$$z_{ij} \in \mathbb{Z}_{\geq 0}, \quad \forall i, j \in \{1, \dots, n\}. \quad (3)$$

Constraints:

$$(1) \text{ Indicator argument: } z_{ii} = \sum_{k=1}^n y_{iki} - \sum_{k=1}^n x_{ik}, \forall i \in \{1, \dots, n\}, \quad (4)$$

$$z_{ij} = \sum_{k=1}^n y_{ikj} + x_{ij}, \quad \forall i \neq j \in \{1, \dots, n\}, \quad (5)$$

$$(2) \text{ Indicator function: } z_{ij} \leq M g_{ij}, \quad \forall i, j \in \{1, \dots, n\}, \quad (6)$$

$$z_{ij} \geq g_{ij}, \quad \forall i, j \in \{1, \dots, n\}, \quad (7)$$

$$(3) \text{ Product: } y_{ikj} \leq x_{ik}, \quad \forall i, k, j \in \{1, \dots, n\}, \quad (8)$$

$$y_{ikj} \leq x_{kj}, \quad \forall i, k, j \in \{1, \dots, n\}, \quad (9)$$

$$y_{ikj} \geq x_{ik} + x_{kj} - 1, \quad \forall i, k, j \in \{1, \dots, n\} \quad (10)$$

$$(4) \text{ Upper Bound: } x_{ij} \leq 1, \quad \forall i, j \in \{1, \dots, n\}, \quad (11)$$

$$y_{ikj} \leq 1, \quad \forall i, k, j \in \{1, \dots, n\}, \quad (12)$$

$$(5) \text{ Symmetry: } x_{ij} = x_{ji}, \quad \forall i, j \in \{1, \dots, n\}, \quad (13)$$

$$(6) \text{ Diagonal: } x_{ii} = 0, \quad \forall i \in \{1, \dots, n\}. \quad (14)$$

1.2 Linear Program

$$x_{ij} \in \mathbb{R}_{\geq 0}, \quad \forall i, j \in \{1, \dots, n\}, \quad (15)$$

$$y_{ikj} \in \mathbb{R}_{\geq 0}, \quad \forall i, k, j \in \{1, \dots, n\}, \quad (16)$$

$$z_{ij} \in \mathbb{R}_{\geq 0}, \quad \forall i, j \in \{1, \dots, n\}, \quad (17)$$

Constraints:

$$(1) \text{ Indicator argument: } z_{ii} = \sum_{k=1}^n y_{iki} - \sum_{k=1}^n x_{ik}, \forall i \in \{1, \dots, n\}, \quad (18)$$

$$z_{ij} = \sum_{k=1}^n y_{ikj} + x_{ij}, \quad \forall i \neq j \in \{1, \dots, n\}, \quad (19)$$

$$(2) \text{ Indicator function: } z_{ij} \leq M g_{ij}, \quad \forall i, j \in \{1, \dots, n\}, \quad (20)$$

$$z_{ij} \geq g_{ij}, \quad \forall i, j \in \{1, \dots, n\}, \quad (21)$$

$$(3) \text{ Product: } y_{ikj} \leq x_{ik}, \quad \forall i, k, j \in \{1, \dots, n\}, \quad (22)$$

$$y_{ikj} \leq x_{kj}, \quad \forall i, k, j \in \{1, \dots, n\}, \quad (23)$$

$$y_{ikj} \geq x_{ik} + x_{kj} - 1, \quad \forall i, k, j \in \{1, \dots, n\} \quad (24)$$

$$(4) \text{ Upper Bound: } x_{ij} \leq 1, \quad \forall i, j \in \{1, \dots, n\}, \quad (25)$$

$$y_{ikj} \leq 1, \quad \forall i, k, j \in \{1, \dots, n\}, \quad (26)$$

$$(5) \text{ Symmetry: } x_{ij} = x_{ji}, \quad \forall i, j \in \{1, \dots, n\}, \quad (27)$$

$$(6) \text{ Diagonal: } x_{ii} = 0, \quad \forall i \in \{1, \dots, n\}. \quad (28)$$

1.3 Max Linear Program

$$x_{ij} \in \mathbb{R}_{\geq 0}, \quad \forall i, j \in \{1, \dots, n\}, \quad (29)$$

$$y_{ikj} \in \mathbb{R}_{\geq 0}, \quad \forall i, k, j \in \{1, \dots, n\}, \quad (30)$$

$$z_{ij} \in \mathbb{R}_{\geq 0}, \quad \forall i, j \in \{1, \dots, n\}, \quad (31)$$

Constraints:

$$(1) \text{ Indicator argument: } z_{ii} = \sum_{k=1}^n y_{iki} - \sum_{k=1}^n x_{ik}, \forall i \in \{1, \dots, n\}, \quad (32)$$

$$z_{ij} = \sum_{k=1}^n y_{ikj} + x_{ij}, \quad \forall i \neq j \in \{1, \dots, n\}, \quad (33)$$

$$(2) \text{ Indicator function: } z_{ij} \leq M g_{ij}, \quad \forall i, j \in \{1, \dots, n\}, \quad (34)$$

$$z_{ij} \geq g_{ij}, \quad \forall i, j \in \{1, \dots, n\}, \quad (35)$$

$$(3) \text{ Product: } y_{ikj} \leq x_{ik}, \quad \forall i, k, j \in \{1, \dots, n\}, \quad (36)$$

$$y_{ikj} \leq x_{kj}, \quad \forall i, k, j \in \{1, \dots, n\}, \quad (37)$$

$$y_{ikj} \geq x_{ik} + x_{kj} - 1, \quad \forall i, k, j \in \{1, \dots, n\} \quad (38)$$

$$(4) \text{ Upper Bound: } x_{ij} \leq 1, \quad \forall i, j \in \{1, \dots, n\}, \quad (39)$$

$$y_{ikj} \leq 1, \quad \forall i, k, j \in \{1, \dots, n\}, \quad (40)$$

$$(5) \text{ Symmetry: } x_{ij} = x_{ji}, \quad \forall i, j \in \{1, \dots, n\}, \quad (41)$$

$$(6) \text{ Diagonal: } x_{ii} = 0, \quad \forall i \in \{1, \dots, n\}, \quad (42)$$

Objective Function:

$$\max \sum_{i=1}^n \sum_{j=1}^n x_{ij}. \quad (43)$$

2 Two-Edge-Triangle Proof

Let x be an optimal solution to the linear relaxation defined by Equations (29)–(43). If $g_{ij} = 0$, then for every $k \in \{1, \dots, n\}$,

$$x_{ik} + x_{kj} = 1.$$

2.1 Integer Program

Lemma 1. *Consider the triangle with three vertices i, j and k where edges ik and kj are present and ij is absent in square G . Then for root G , the integer program defined by Equations (1)–(14) must satisfy*

$$x_{ik} + x_{kj} \leq 1.$$

Proof. As ij is absent in G , $g_{ij} = 0$ such that Equation (6) gives $z_{ij} \leq 0$ and Equation (7) gives $z_{ij} \geq 0$ such that $z_{ij} = 0$. Therefore Equation (5) gives

$$0 = \sum_{a=1}^n y_{iaj} + x_{ij} = \sum_{a=1}^n x_{ia} x_{aj} + x_{ij},$$

where the second equality follows from the product constraints Equations (8)–(14). Using Equation (1), x_{ij} is an integer and hence 0 only holds when $x_{ij} = 0$ and all $x_{ia}x_{aj} = 0$. This implies that at most one of x_{ia} , x_{aj} is equal to 1 and the other must be 0, and can be equivalently written as $x_{ia} + x_{aj} \leq 1$.

Taking $a = k$ gives $x_{ik} + x_{kj} \leq 1$ as required. $\square$

2.2 Linear Program

Lemma 2. *Consider the triangle with three vertices i , j and k where edges ik and kj are present and ij is absent in square G . Then for root G , the linear program defined by Equations (15)–(28) must satisfy*

$$x_{ik} + x_{kj} \leq 1.$$

Proof. The linear program is the relaxation of the integer program obtained by removing the integrality constraints on the variables while retaining all constraints. Since the proof of Lemma 1 relies only on these constraints, the inequality also holds for every variable forming a feasible solution to the linear program. $\square$

2.3 Max Linear Program

Proposition 1. *Let x be a feasible solution to the maximum linear relaxation defined by Equations (29)–(43). Suppose there exists vertices i , j and k such that*

$$g_{ik} = 1, \quad g_{kj} = 1, \quad g_{ij} = 0 \quad \text{and} \quad x_{ik} + x_{kj} < 1.$$

Then there exists another feasible solution (x', y', z') satisfying $\sum_{u,v} x'_{uv} > \sum_{u,v} x_{uv}$.

Proof. Starting from initial feasible solution x let $x_{ik} + x_{kj} = 1 - \delta$ for some $\delta > 0$ and choose $0 < \varepsilon \leq \delta$ with $0 < \varepsilon \leq 1 - x_{ik} - x_{kj}$ to consider increasing x_{ik} by ε . By Equation (32), increasing x_{ik} decreases

$$\sum_{t=1}^n y_{iti} - \sum_{t=1}^n x_{it}$$

by ε . Therefore, to maintain Equation (32), either

$\square$

Theorem 1. *Let x be an optimal solution to the linear relaxation defined by Equations (29)–(43). If $g_{ij} = 0$, then for every $k \in \{1, \dots, n\}$,*

$$x_{ik} + x_{kj} = 1.$$

Proof. Using Lemma 2 $x_{ik} + x_{kj} \leq 1$ and using Proposition 1 $x_{ik} + x_{kj} \geq 1$ such that $x_{ik} + x_{kj} = 1$ as required. $\square$