

HOLY CROSS COLLEGE

SEMESTER 2, 2017

Question/Answer Booklet

SOLUTIONS

## 12 PHYSICS

Please place your student identification label in this box

Student Name \_\_\_\_\_

Student's Teacher \_\_\_\_\_

### Time allowed for this paper

Reading time before commencing work: 10 minutes

Working time for paper: 3 hours

### Materials required/recommended for this paper

#### *To be provided by the supervisor*

This Question/Answer Booklet

Multiple-choice Answer Sheet

Data Sheet

#### *To be provided by the candidate*

Standard items: pens, pencils, eraser, correction fluid, ruler, highlighters

Special items: non-programmable calculators satisfying the conditions set by the School Curriculum and Standards Authority for this course

### Important note to candidates

No other items may be taken into the examination room. It is **your** responsibility to ensure that you do not have any unauthorised notes or other items of a non-personal nature in the examination room. If you have any unauthorised material with you, hand it to the supervisor **before** reading any further.

## Structure of this paper

Section	Number of questions available	Number of questions to be answered	Suggested working time (minutes)	Marks available	Percentage of exam
Section One: Short Answers	11	11	50	53	30
Section Two: Problem-solving	7	7	90	91	50
Section Three: Comprehension	2	2	40	36	20
				180	100

## Instructions to candidates

1. The rules for the conduct of examinations at Holy Cross College are detailed in the College Examination Policy. Sitting this examination implies that you agree to abide by these rules.
2. Write your answers in this Question/Answer Booklet.
3. Working or reasoning should be clearly shown when calculating or estimating answers.
4. You must be careful to confine your responses to the specific questions asked and to follow any instructions that are specific to a particular question.
5. Spare pages are included at the end of this booklet. They can be used for planning your responses and/or as additional space if required to continue an answer.
  - Planning: If you use the spare pages for planning, indicate this clearly at the top of the page.
  - Continuing an answer: If you need to use the space to continue an answer, indicate in the original answer space where the answer is continued, i.e. give the page number. Fill in the number of the question(s) that you are continuing to answer at the top of the page.
6. Answers to questions involving calculations should be **evaluated and given in decimal form**. It is suggested that you quote all answers to **three significant figures**, with the exception of questions for which estimates are required. Despite an incorrect final result, credit may be obtained for method and working, providing these are **clearly and legibly set out**.
7. Questions containing the instruction "**estimate**" may give insufficient numerical data for their solution. Students should provide appropriate figures to enable an approximate solution to be obtained. Give final answers to a maximum of **two significant figures** and include appropriate units where applicable.
8. Note that when an answer is a vector quantity, it must be given with magnitude and direction.
9. In all calculations, units must be consistent throughout your working.

**Additional Data**

Table of common mesons

Particle	Structure	Charge	Baryon Number	Strangeness
$\pi^0$	$u\bar{u}$ or $d\bar{d}$	0	0	0
$\pi^+$	$u\bar{d}$	+1	0	0
$\pi^-$	$\bar{u}d$	-1	0	0
$K^0$	$d\bar{s}$	0	0	+1
$K^+$	$u\bar{s}$	+1	0	+1
$K^-$	$\bar{u}s$	-1	0	-1

**Properties of quarks**  
*antiquarks have opposite signs*

type	charge	baryon number	strangeness
u	$+\frac{2}{3}e$	$+\frac{1}{3}$	0
d	$-\frac{1}{3}e$	$+\frac{1}{3}$	0
s	$-\frac{1}{3}e$	$+\frac{1}{3}$	-1

**Properties of leptons**

	lepton number
particles: $e^-$ , $\nu_e$ ; $\mu^-$ , $\nu_\mu$	+1
antiparticles: $e^+$ , $\overline{\nu}_e$ ; $\mu^+$ , $\overline{\nu}_\mu$	-1

**Redshift and recessional velocity**

$$z = \frac{\Delta\lambda}{\lambda} \quad \text{It can also be shown that:} \quad z = \frac{v}{c_0}$$

where:  $z$  = redshift  
 $\Delta\lambda$  = change in wavelength (moving source) (nm)  
 $\lambda$  = wavelength of stationary source (nm)  
 $v$  = recessional speed of galaxy ( $\text{ms}^{-1}$ )  
 $c_0$  = speed of light in a vacuum ( $\text{ms}^{-1}$ )

**Hubble's Law**

$$v_{\text{galaxy}} = H_0 d$$

$V_{\text{galaxy}}$  = recessional speed of galaxy ( $\text{kms}^{-1}$ )  
 $d$  = distance to galaxy (Mpc)  
 $H_0$  = Hubble's constant ( $\text{kms}^{-1}\text{Mpc}^{-1}$ )


**Astronomical distances**

$$1.00 \text{ Mpc} = 3.26 \times 10^6 \text{ ly}$$

**SECTION ONE: Short Answers****Marks Allotted: 53 marks out of 180 total.**Attempt **ALL 11** questions in this section. Answers are to be written in the space below or next to each question.


1. The distance between the proton and the electron in the ground state of the hydrogen atom is defined as the Bohr radius. Given that the Bohr radius can be measured as  $r = 5.29 \times 10^{-11}$  metres, what is the speed of the electron as it orbits in the atom?

[5 marks]



$$\begin{aligned}
 F_{\text{Electric}} &= F_{\text{Centripetal}} \quad (1) \\
 \Rightarrow \frac{1}{4\pi\epsilon_0} \cdot \frac{q_1 q_2}{r^2} &= \frac{mv^2}{r} \quad (1) \\
 \Rightarrow v &= \sqrt{\frac{q_1 q_2}{4\pi\epsilon_0 mr}} \quad (1) \\
 &= \sqrt{\frac{(1.60 \times 10^{-19})^2}{4\pi(8.85 \times 10^{-12})(9.11 \times 10^{-31})(5.29 \times 10^{-11})}} \quad (1) \\
 &= 2.19 \times 10^6 \text{ ms}^{-1} \quad (1)
 \end{aligned}$$

2. Consider the decay equation shown here showing the hadronic process of lambda decay.



**Note:**  $\Lambda = uds$ ,  $p = uud$ ,  $\pi^- = d\underline{u}$

- (a) Show that charge and baryon number are conserved.

[2 marks]

$$\begin{aligned}
 \text{Charge: } \frac{2}{3} + \left(-\frac{1}{3}\right) + \left(-\frac{1}{3}\right) &\rightarrow \frac{2}{3} + \frac{2}{3} + \left(-\frac{1}{3}\right) + \left(-\frac{2}{3}\right) \quad (1) \\
 \Rightarrow 0 &= 0
 \end{aligned}$$

$$\begin{aligned}
 \text{Baryon number: } \frac{1}{3} + \frac{1}{3} + \frac{1}{3} &\rightarrow \frac{1}{3} + \frac{1}{3} + \frac{1}{3} + \frac{1}{3} + \left(-\frac{1}{3}\right) \quad (1) \\
 \Rightarrow 1 &= 1
 \end{aligned}$$

- (b) Is strangeness conserved? All working must be shown to explain your answer.


[3 marks]

$$\begin{aligned}
 0 + 0 - 1 &\rightarrow 0 + 0 + 0 + 0 + 0 \quad (1) \\
 \Rightarrow -1 &\neq 0 \quad (1)
 \end{aligned}$$

$\therefore \underline{\text{No.}} \quad (1)$

3. A wire is conducting a DC current of 1.50 A. At point X, a magnetic flux density of  $3.86 \times 10^{-6}$  T is detected. Calculate the distance between the current carrying wire and point X. (Ignore the effects of the Earth's magnetic field in this question.)

[2 marks]



$$B = \frac{\mu_0 I}{2\pi r}$$

$$\begin{aligned} \Rightarrow r &= \frac{\mu_0 I}{2\pi B} \\ &= \frac{(4\pi \times 10^{-7})(1.50)}{2\pi (3.86 \times 10^{-6})} \quad (1) \\ &= \underline{7.77 \times 10^{-2} \text{ m}} \quad (1) \end{aligned}$$


4. A proton has been accelerated to 95.0 % of the speed of light in the Large Hadron Collider (LHC). Calculate its relativistic energy.

[3 marks]

$$\begin{aligned} E &= \frac{mc^2}{\sqrt{1 - \frac{v^2}{c^2}}} \\ &= \frac{(1.67 \times 10^{-27})(3.00 \times 10^8)^2}{\sqrt{1 - \frac{(0.950c)^2}{c^2}}} \quad (1) \\ &= \underline{4.81 \times 10^{-10} \text{ J}} \quad (1) \end{aligned}$$

5. The diagram shows a uniform ladder of mass 20.0 kg and length 5.00 m resting on firm ground and against a frictionless wall. Friction acts at the base of the ladder from the ground (as shown) to stop the ladder collapsing. The force from the wall ( $F_{\text{wall}}$ ) and friction both act in the horizontal and are in equilibrium. A person of mass 80.0 kg is standing on the ladder 3.50 m from the base. The ladder makes an angle of  $65.0^\circ$  with the ground.

- (a) Calculate the force of friction acting on the ladder in the position shown. [4 marks]



$$\sum \text{CM} = \sum \text{ACM} \quad (1)$$

$$\Rightarrow (196 \cos 65.0^\circ)(2.50) + (784 \cos 65.0^\circ)(3.50) = (F_{\text{w}} \cos 25.0^\circ)(5.00) \quad (2)$$

$$\Rightarrow F_{\text{w}} = \frac{302 \text{ N}}{= F_{\text{FRICTION}}} \quad (1)$$

- (b) If the angle that the ladder makes to the horizontal is changed to  $45.0^\circ$ , how would this change the magnitude of friction required to maintain equilibrium? The friction would:

increase (1) stay the same decrease insufficient data to determine

Circle a response and explain your choice.

[3 marks]

- $\Sigma \text{CM}$  increases. (1)
- $\Sigma \text{ACM}$  must increase to maintain equilibrium.  
∴  $F_{\text{w}}$  must increase. (1)

6. A subatomic particle, consisting of a quark-antiquark pair, has a rest-life measured to be  $2.34 \times 10^{-8}$  s in the laboratory. What is the speed of the same particle, if its average lifetime is measured to be  $3.95 \times 10^{-8}$  s in a cosmic ray experiment? [6 marks]

$$t = \frac{t_0}{\sqrt{1 - \frac{v^2}{c^2}}}$$

$$\Rightarrow 3.95 \times 10^{-8} = \frac{2.34 \times 10^{-8}}{\sqrt{1 - \frac{v^2}{c^2}}} \quad (2)$$

$$\Rightarrow 1.56 \times 10^{-15} \left(1 - \frac{v^2}{c^2}\right) = 5.48 \times 10^{-16} \quad (2)$$

$$\Rightarrow 1.56 \times 10^{-15} - 1.73 \times 10^{-32} v^2 = 5.48 \times 10^{-16} \quad (1)$$

$$\Rightarrow \underline{v = 2.42 \times 10^8 \text{ ms}^{-1}} \quad (1)$$

7. The light received from many distant galaxies is red-shifted.

- (a) State the cause of this red-shift, using a diagram to explain your answer. [3 marks]

- Expansion of space-time - distant galaxies are moving away from us. (1)
- Wavelength of light appears "stretched". ( $\frac{1}{2}$ )
- Absorption spectrum appears moved towards the red end of the spectrum. ( $\frac{1}{2}$ )



- (b) A spectral line of light from a laboratory source has a wavelength of 658 nm. The wavelength of this line of light from a distant galaxy (NGC5128) is 670 nm. Calculate the recessional speed of NGC5128. [2 marks]

$$V = \frac{\Delta\lambda}{\lambda} c$$

$$= \frac{(670 \times 10^{-9} - 658 \times 10^{-9})}{(658 \times 10^{-9})} (3 \times 10^8) \quad (1)$$


$$= \underline{5.47 \times 10^6 \text{ ms}^{-1}} \quad (1)$$

- (c) Given that a current value for Hubble's constant is  $72.0 \text{ kms}^{-1}\text{Mpc}^{-1}$ , how far is the observer from NGC5128? [2 marks]

$$\begin{aligned}
 V &= H_0 d \\
 \Rightarrow d &= \frac{V}{H_0} \\
 &= \frac{5.47 \times 10^3}{72.0} \quad (1) \\
 &= 76.0 \text{ Mpc} \quad (1)
 \end{aligned}$$

8. Consider the table of elementary particles shown below.


- (a) Write the missing headings in the spaces provided. [2 marks]



- (b) Briefly explain the difference between mesons and baryons. [2 marks]

- MESON - made up of 2 quarks. (1)
- BARYONS - made up of 3 quarks. (1)

9. A current-carrying straight conductor is placed in a magnetic field and experiences a magnetic force equal to 75.0 % of the maximum value this force could be in this field. Calculate the size of the angle  $\theta$  between the conductor and the magnetic field. Show working. [3 marks]



$$\text{For max: } F = IlB.$$

$$\text{At angle } \theta: 0.750 F = IlBs \sin \theta \quad (1)$$

$$\Rightarrow \sin \theta = 0.750 \quad (1)$$


$$\Rightarrow \theta = 48.6^\circ \quad (1)$$

10. An AC Generator has an armature with an area of  $4.70 \times 10^{-2} \text{ m}^2$ . The coil rotates from vertical to horizontal in 25.0 ms through a magnetic field strength of 0.430 T.

- (a) Identify the parts 'A' and 'B' shown in the diagram. [1 mark]

Part A: slip rings  $(\frac{1}{2})$

Part B: brushes  $(\frac{1}{2})$



- (b) How many loops must the coil contain in order to generate a potential difference ( $V_{\max}$ ) of 175 volts? [3 marks]

$$f = \frac{1}{T}$$

$$= \frac{1}{4(25.0 \times 10^{-3})}$$

$$= 10.0 \text{ Hz.} \quad (1)$$

$$\text{EMF} = 2\pi N B A f$$

$$\Rightarrow N = \frac{\text{EMF}}{2\pi B A f}$$

$$= \frac{(175)}{2\pi (0.430)(4.70 \times 10^{-2})(10.0)} \quad (1)$$

$$= 138 \text{ loops} \quad (1)$$

11. A photoelectric effect experiment was performed in which a monochromatic light beam was shone onto a clean metal surface. The wavelength of the incident beam was varied and the maximum kinetic energy of the emitted photoelectrons was recorded in the table below.

Wavelength (nm)	Frequency (Hz)	$KE_{max}$ (eV)	$KE_{max}$ (J)
750	$4.00 \times 10^{14}$	0.22	$3.52 \times 10^{-20}$
587	$5.11 \times 10^{14}$	0.67	$1.07 \times 10^{-19}$
506	$5.93 \times 10^{14}$	(1) 0.98	$1.57 \times 10^{-19}$
444	$6.76 \times 10^{14}$	1.35	$2.16 \times 10^{-19}$
400	$7.50 \times 10^{14}$	1.63	$2.61 \times 10^{-19}$

- (a) Briefly explain the significance of the photoelectric effect to modern physics.

[2 marks]

- Demonstrated the quantisation of energy.
- Showed the particle nature of light.

- (b) Complete the table for the missing values.

[2 marks]

- (c) By making use of the data from the table, calculate the work function of the clean metal surface used during this experiment.

[3 marks]

$$\begin{aligned}
 E &= hf = W + E_K \\
 \Rightarrow W &= hf - E_K \quad (1) \\
 &= (6.63 \times 10^{-34})(4.00 \times 10^{14}) - 3.52 \times 10^{-20} \quad (1) \\
 &= 2.30 \times 10^{-19} \text{ J} \\
 &= 1.44 \text{ eV.} \quad (1)
 \end{aligned}$$



**This page has been deliberately left blank.**

**SECTION TWO: Problem Solving****Marks allotted: 91 marks out of 180 marks total.**


Attempt **ALL** 7 questions in this section. The marks allocated to each question are given and the answers should be written in the spaces provided.

**12. [7 marks]**

A car of mass  $2.20 \times 10^3$  kg is in horizontal circular motion on a banked track. The car has a speed of  $14.0 \text{ ms}^{-1}$  and is relying on friction to stay at a fixed height on the banked track. The radius of the circle is 32.0 m. The track is banked at an angle of  $20.0^\circ$  to the horizontal. Friction acts from the track onto the car parallel to the track as shown.



- (a) Construct a vector diagram to the right of the diagram above. Show the forces acting on the car and the nett force. [2 marks]
- (b) Calculate the magnitude of friction acting on the car from the banked track. [5 marks]



$$\sum F_v = 0 \Rightarrow R \cos 20.0^\circ = F_W + F_{FR} \cos 70.0^\circ \quad (1)$$

$$\Rightarrow R = \frac{(2.20 \times 10^3)(9.80) + F_{FR} \cos 70.0^\circ}{\cos 20.0^\circ}$$

$$\Rightarrow R = 2.294 \times 10^4 + 0.364 F_{FR} - \textcircled{1} \quad (1)$$

HORizontally:  $F_h = F_c$

$$\Rightarrow R \cos 70.0^\circ + F_{FR} \cos 20.0^\circ = \frac{mv^2}{r} \quad (1)$$

Sub \textcircled{1} for R

$$\therefore (2.294 \times 10^4 + 0.364 F_{FR})(\cos 70.0^\circ) + F_{FR} \cos 20.0^\circ = \frac{(2.20 \times 10^3)(14.0)^2}{(32.0)}$$

$$\Rightarrow 7.846 \times 10^3 + 0.1245 F_{FR} + 0.9397 F_{FR} = 1.348 \times 10^4 \quad (1)$$

$$\Rightarrow F_{FR} = \underline{5.29 \times 10^3 \text{ N}} \quad (1)$$

## 13. [15 marks]

Our Sun is a medium sized star that is part of a spiral galaxy called the Milky Way. Like all spiral galaxies, the stars in the Milky Way rotate around a galactic centre.

Our Sun's orbit is virtually circular with a radius of  $2.50 \times 10^{20}$  m (about 26 000 light years); its average orbital speed is about  $2.20 \times 10^5$  ms $^{-1}$ .

- (a) Calculate the orbital period of the Sun around the galactic centre of the Milky Way  
(in years). [4 marks]

$$\begin{aligned} V &= \frac{2\pi r}{T} && (1) \\ &= \frac{2\pi (2.50 \times 10^{20})}{(2.20 \times 10^5)} && (1) \\ &= 7.14 \times 10^{15} \text{ s} && (1) \\ &= \underline{2.26 \times 10^8 \text{ years}} && (1) \quad (\text{using } 1 \text{ year} = 365.25 \text{ days}) \end{aligned}$$

- (b) Calculate the gravitational field strength due to the Milky Way galaxy at the Sun's distance from the galactic centre. [3 marks]

$$\begin{aligned} g = a_c &= \frac{V^2}{r} && (1) \\ &= \frac{(2.20 \times 10^5)^2}{2.50 \times 10^{20}} && (1) \\ &= \underline{1.94 \times 10^{-10} \text{ N kg}^{-1} (\text{ms}^{-2})} && (1) \end{aligned}$$

- (c) The circular orbit of the Sun around the galactic centre of the Milky Way is due to the gravitational force of attraction between the Sun's and Milky Way's centres of mass.

Use the data provided and answers calculated thus far to show that the mass of our galaxy inside our Sun's orbit must be about  $1.80 \times 10^{41}$  kg.

[If you could not calculate an answer to part (a), use  $7.00 \times 10^{15}$  s; if you could not calculate an answer to part (b), use  $1.90 \times 10^{-10}$  N kg $^{-1}$ .]

[3 marks]

$$\begin{aligned} g &= \frac{GM}{r^2} \\ \Rightarrow M &= \frac{gr^2}{G} && (1) \\ &= \frac{(1.94 \times 10^{-10})(2.50 \times 10^{20})^2}{(6.67 \times 10^{-11})} && (1) \\ &= \underline{1.81 \times 10^{41} \text{ kg}} && (1) \end{aligned}$$

[Alternate answer =  $1.78 \times 10^{41}$  kg]

- (d) If the mass of our Sun can be considered to be an average mass for the stars in our galaxy, estimate how many stars there must be inside our Sun's orbit in the Milky Way. Show your working. [2 marks]

$$\text{Number of stars} = \frac{1.81 \times 10^{41}}{1.99 \times 10^{30}} \quad (1)$$

$$= 9.09 \times 10^{10} \quad (1)$$

- (e) The mass of the Milky Way inside our Sun's orbit is about  $1.80 \times 10^{41}$  kg, which is about  $10^{11}$  times the mass of our Sun. However, when scientists estimate the mass of the **visible matter** inside the Sun's orbit, it only comes to about  $10^{10}$  times the mass of our Sun.

- (i) What does this imply about the types of matter in our Galaxy? [2 marks]


- A significant amount of matter is not visible. (1)
- The invisible matter still exerts a gravitational force on other matter. (1)

- (ii) If the mass of our galaxy was only  $10^{10}$  times the mass of our Sun, describe one (1) effect this would have on our Sun's motion. (1 mark)

- A decrease in orbital velocity.
  - An increase in orbital period
- } Either OK - 1 mark.

## 14. [13 marks]

A stone of mass 52.0 g is thrown from a building of height 30.0 m. The stone is launched with an angle of elevation of  $38.0^\circ$  above the horizontal. It takes a time of 3.15 s for the stone to reach ground level. You can ignore air resistance for this question.



- (a) Calculate the initial launch speed  $u$  of the stone.

[4 marks]

VERTICALLY  $\downarrow$   $+ve$

$$v = ? \quad \text{Correct sign convention (1)} \quad s = ut + \frac{1}{2}at^2$$

$$u = -(u \cos 52.0^\circ) \text{ ms}^{-1} \quad (1) \quad \Rightarrow 30.0 = -(u \cos 52.0^\circ)(3.15) + \frac{1}{2}(9.80)(3.15)^2 \quad (1)$$

$$a = 9.80 \text{ ms}^{-2}$$

$$t = 3.15 \text{ s} \quad \Rightarrow \quad u = \underline{9.60 \text{ ms}^{-1}} \quad (1)$$

$$s = 30.0 \text{ m}$$

[For the following calculations, use a numerical value of  $9.50 \text{ ms}^{-1}$  for the initial launch speed of stone if you could not calculate an answer for part (a).]

- (b) Calculate the horizontal range of the stone.

[2 marks]

$$V_h = \frac{s_h}{t}$$

$$\Rightarrow s_h = (9.60 \cos 38.0^\circ)(3.15) \quad (1)$$

$$= \underline{23.8 \text{ m}} \quad (1)$$

- (c) Calculate the velocity of the stone after 2.50 s of flight. You must give a magnitude and direction. [5 marks]

VERTICALLY

$$V_v = ?$$

$$u = -(9.60 \cos 52.0^\circ) \text{ ms}^{-1}$$

$$a = 9.80 \text{ ms}^{-2}$$

$$t = 2.50 \text{ s}$$

$$s = ?$$

$$v = u + at$$

$$= (-9.60 \cos 52.0^\circ) + (9.80)(2.50) \quad (1)$$

$$= 18.6 \text{ ms}^{-1} \text{ down} \quad (1)$$

$$V_h = 9.60 \cos 38.0^\circ$$

$$= 7.56 \text{ ms}^{-1} \quad (1)$$

$$\vec{R} = \sqrt{(18.6)^2 + (7.56)^2}$$

$$= 20.1 \text{ ms}^{-1} \quad (1)$$

$$\tan \theta = \frac{18.6}{7.56}$$

$$\Rightarrow \theta = 67.9^\circ \quad (1)$$

$\therefore$  velocity =  $20.1 \text{ ms}^{-1}$  at  $67.9^\circ$  to the horizontal

- (d) Calculate the work done on the stone by the Earth's gravitational field in the motion from launch to reaching ground level. [2 marks]


$$W = \Delta E_p = mg \Delta h$$

$$= (0.0520)(9.80)(30.0) \quad (1)$$

$$= \underline{15.3 \text{ J}} \quad (1)$$

## 15. [16 marks]

A nature lookout consists of an elevated concrete walkway high above the ground. A uniform platform has been constructed so people can walk out over a gorge and view it. The entire platform structure is shown in the figure below.




The platform is designed to support a load of 8.50 tonnes and is 15.0 m long. A single steel cable supports the platform, attached 10.0 m from the end at  $55.0^\circ$  as shown in the figure. The platform has a mass of 0.700 tonnes.

The platform is uniform and it can be assumed that, when it is acting, the 8.50 tonne maximum load acts half-way along its length.

The steel cable shown has a maximum tensile strength of  $1.50 \times 10^5$  N.

- On the diagram above, draw all the forces acting on the platform when in it is an unloaded state as drawn above. Label the forces appropriately. [3 marks]
- Show that with the maximum load acting through the platform's midpoint, the cable will be able to support the platform. Support your answer with calculations. [4 marks]




$$\begin{aligned} \sum CM &= \sum ACM \\ \Rightarrow (9.02 \times 10^4)(7.50) &= (T \cos 35.0^\circ)(10.0) \quad (1) \\ \Rightarrow T &= 8.26 \times 10^4 \text{ N} \quad (1) \\ \text{As } T < 1.50 \times 10^5 \text{ N, cable does not break.} \quad (1) \end{aligned}$$

- (c) Hence, calculate the force that the wall exerts on the platform.

[Hint - if you could not calculate an answer for part (a), use a value of  $9.00 \times 10^4 \text{ N}$  for the tension in the cable.]

[4 marks]



$$\begin{aligned}\vec{R} &= \sqrt{(9.02 \times 10^4)^2 + (8.26 \times 10^4)^2 - 2(9.02 \times 10^4)(8.26 \times 10^4) \cos 35.0^\circ} \\ &= 5.25 \times 10^4 \text{ N} \quad (1)\end{aligned}$$

$$\frac{8.26 \times 10^4}{\sin \theta} = \frac{5.25 \times 10^4}{\sin 35.0^\circ}$$

$$\Rightarrow \theta = 64.5^\circ \quad (1)$$

$$\therefore \vec{R} = 5.25 \times 10^4 \text{ N at } 64.5^\circ \text{ to the vertical} \quad (1)$$


- (d) If the maximum load of 8.50 tonnes is gradually moved towards the end of the platform, describe what happens to the magnitude and direction of the force you calculated in part (c). No calculation is required.

[2 marks]

By inspection:  $\vec{R}$  gets bigger (1)  
 $\theta$  gets bigger (1)

- (e) If the maximum load continues to move towards the end of the platform, the cable will eventually exceed its load limit and snap. Calculate how far towards the edge of the platform the load can move until the load limit on the wire is exceeded.


[3 marks]



$$\begin{aligned}\sum \text{CM} &= \sum \text{ACM} \\ \Rightarrow (6.86 \times 10^3)(7.50) + (8.50 \times 10^3)(9.30) r &= (1.50 \times 10^5 \cos 35.0^\circ)(10.0) \quad (2) \\ \Rightarrow r &= 14.1 \text{ m} \quad (1)\end{aligned}$$

## 16. [13 marks]

The diagram shows the side view of a DC electric motor. A square coil is placed flat in the uniform magnetic field between the North and South magnetic poles. Current direction in the coil is shown on the sides adjacent to the magnetic poles. The commutator and carbon brushes are also shown.



- (a) In which direction will the coil turn from this start position? [1 mark]

\* Anti-clockwise. (1)

- (b) Explain the function of the **brushes** and the **commutator**. [3 marks]

BRUSHES - transfer current from an external source into the coil. (1)

COMMUTATOR - switches the direction of the current in the coil every 180° of rotation. (1)

- ensures force (torque) is in a constant direction. (1)

- (c) On the diagram above, use the symbols  $\odot$  and  $\otimes$  to sketch the location of the coil sides adjacent to the magnetic poles after 30.0° of rotation from this start position. Put arrows on your symbols to indicate the direction of magnetic force acting on them. [2 marks]

- (d) At this new position after  $30.0^\circ$  of rotation from the start position, determine the torque value of the motor as a percentage of maximum torque. [2 marks]

$$\tau_{(\text{max})} = Fr \sin 90^\circ \quad \therefore \quad \frac{\tau}{\tau_{(\text{max})}} = \frac{\sin 60^\circ}{\sin 90^\circ} \times \frac{100}{1} \quad (1)$$

$$\tau = Fr \sin 60^\circ \quad = \underline{86.6\%} \quad (1)$$

- (e) A single 0.120 m length of wire, adjacent to one of the magnetic poles, experiences a 0.0280 N magnitude of force when a current of 5.30 A is present. Calculate the magnetic flux density between the poles.

[2 marks]

$$F = IlB$$

$$\Rightarrow B = \frac{F}{Il}$$

$$= \frac{0.0280}{(5.30)(0.120)} \quad (1)$$

$$= \underline{4.40 \times 10^{-2} \text{ T}} \quad (1)$$

- (f) After the motor is switched on, its rate of rotation increases. As this happens, the nett current in the coil decreases. Clearly explain why this happens.

[3 marks]

- As the wire cuts through the magnetic field, an EMF is induced in the coil (called a back-EMF). (1)
- This moves the charges in the opposite direction in the coil to produce an opposing force. (1)
- The current will decrease due to the reduction of the EMF. (1)

## 17. [12 marks]

The trains on the Perth to Fremantle rail line are powered by four 600.0 V DC motors. The current is delivered to the motors from the sub-station overhead power lines, which are at a potential of  $2.50 \times 10^4$  V AC. The AC voltage needs to be converted to 600.0 V DC by a transformer.

The overhead lines have a resistance of  $2.10 \Omega \text{km}^{-1}$  and the motors each have a resistance of  $2.00 \Omega$ . When the train is close to the Perth sub-station and operating at full power, the train draws 1.00 MW of electrical power.

- (a) Why do overhead transmission lines operate at 25.0 kV AC and not at 600.0 V DC?

[2 marks]

- Transformers only work on AC - they need a continual change in flux. (1)
- Power is transmitted at high voltage (25.0 kV) to minimise energy losses in the transmission lines. (1)

- (b) What is the current in the overhead lines when the train is close to the Perth sub-station and operating at full power?

[2 marks]

$$\begin{aligned} P &= VI \\ \Rightarrow I &= \frac{1.00 \times 10^6}{2.50 \times 10^4} \quad (1) \\ &= \underline{400 \text{ A}} \quad (1) \end{aligned}$$

- (c) What is the starting current in **ONE** motor?

[2 marks]

$$\begin{aligned} V &= IR \\ \Rightarrow I &= \frac{V}{R} \\ &= \frac{6.00 \times 10^2}{2.00} \quad (1) \\ &= \underline{3.00 \times 10^2 \text{ A}} \quad (1) \end{aligned}$$

- (d) If the train is 20.0 km from the sub-station, the power developed by the train will be less than when it is close to the substation. If the train is now drawing 0.700 MW and the current drawn from the power lines is 28.0 A, what is the voltage available to the motors?

[6 marks]

$$R_{wires} = (20.0)(2 \times 2.10) \quad (1)$$

$$= 84.0 \Omega \quad (1)$$

$$V_{drop} = I R_{wires}$$

$$= (28.0)(84.0) \quad (1)$$

$$= 2.35 \times 10^3 V \quad (1)$$


$$V_{train} = V_{wires} - V_{drop}$$

$$= 2.50 \times 10^4 - 2.35 \times 10^3 \quad (1)$$

$$= \underline{2.26 \times 10^4 V \quad (22.6 kV)} \quad (1)$$

## 18. [15 marks]

Consider the following diagram that shows the same collection of minerals in (a) daylight and (b) "black light".



- (a) Complete the following sentence: The correct terminology for "black light" is

ultraviolet light (1)

and the phenomenon is called

fluorescence (1).

[2 marks]

- (b) The first 4 energy levels for a potassium mineral are shown (not to scale) as follows.

$E_4 = -1.10 \text{ eV}$	3.29 eV
$E_3 = -1.78 \text{ eV}$	2.61 eV
$E_2 = -2.87 \text{ eV}$	1.52 eV
$E_1 = -4.39 \text{ eV}$	0.00 eV

Could a sample of this potassium mineral display the phenomenon as shown in (a) above? Justify your answer, showing the necessary calculations. [4 marks]

$$\begin{aligned} \text{Max. energy} &= E_4 - E_1 = hf \\ \Rightarrow \left[ (-1.10) - (-4.39) \right] (1.60 \times 10^{-19}) &= (6.63 \times 10^{-34}) f \quad (1) \\ \Rightarrow f &= 7.94 \times 10^{14} \text{ Hz.} \quad (1) \end{aligned}$$

From the data sheet, this is in the visible region of the spectrum.  
- Fluorescence won't occur. (1)

- (c) Consider again the first 4 energy levels for the potassium mineral in (b). What would be detected if particles of the sample were bombarded by:

[4 marks]

- (i) photons of energy 2.65 eV?

- Nothing (1)
- Doesn't match an energy level above ground state.

Mention scattered electrons. (1)

- (ii) electrons of energy 2.65 eV?

- Transition  $E_1 \rightarrow E_3 \Rightarrow$  scattered electrons would have  $E_k = 0.04\text{eV}$ . (1)
- Transition  $E_1 \rightarrow E_2 \Rightarrow$  scattered electrons would have  $E_k = 1.13\text{eV}$  (1)

- (d) If an electron was excited from the ground state to the - 1.78 eV level:

- (i) when it returned to the ground state, what would be the wavelength of the photon emitted? [3 marks]

$$E_3 - E_1 = hf = \frac{hc}{\lambda} \quad (1)$$

$$\Rightarrow [( -1.78 ) - (-4.39)] (1.60 \times 10^{-19}) = \frac{(6.63 \times 10^{-34})(3.00 \times 10^8)}{\lambda} \quad (1)$$

$$\Rightarrow \lambda = 4.76 \times 10^{-7} \text{ m} \quad (1)$$

- (ii) To which region of the electromagnetic spectrum would the photon belong? Support your answer by referring to its wavelength or frequency. [2 marks]

- $\lambda \approx 5 \times 10^{-7} \text{ m}$  (1)

- From the data sheet, this is in the blue/violet region of the visible spectrum. (1)

This page has been deliberately left blank.

**SECTION C: Comprehension and Interpretation****Marks Allotted: 36 marks out of 180 marks total.**


Read the passage carefully and answer all of the questions at the end. Candidates are reminded of the need for correct English and clear and concise presentation of answers. Diagrams (sketches), equations and/or numerical results should be included where appropriate.

**Question 19 [18 marks]****Particle Physics — Basic Principles and Techniques**

Particle physics is the modern version of the age-old quest — to find the smallest particles that cannot be broken down. Particle accelerators are the 'laboratory equipment' in this area of study.

Charged particles can be accelerated in two senses - by their change of direction in circular paths or by increasing their speed. Studies can be made on the radiation that they emit whilst being accelerated or the after effects of collisions between high speed particles.

The cyclotron first came into use in 1928 using a combination of magnetic and electric fields to accelerate particles in a spiral path. Development of this technology led to the synchrotron, which uses an evacuated circular tube with many magnets placed around its circumference. As particles are accelerated, the electric field is adjusted and the strength of the magnets is increased to maintain a constant radius and compensate for relativistic effects that become important at high particle energies.



Any charged particle that accelerates will radiate electromagnetic energy. This is true even at a constant speed in a circular path. So, a continual supply of energy is required in synchrotrons to just maintain a constant particle speed, let alone increase their speed. The emitted radiation is known as synchrotron radiation and can cover the entire electromagnetic spectrum.


Linear accelerators (LINACs) use a straight path and a series of accelerating voltages as the particles move along the line. LINACs are often used to provide the early stages of acceleration before particles are fed into large synchrotrons.

Collider experiments take two beams of particles that have been separately accelerated in opposite directions and smash them into each other. This is difficult to achieve but if successful, it is an efficient use of energy.


When two particles with an equal magnitude of momentum collide head on, the total momentum is zero before and after the collision. If particles are stationary after the collision, then their kinetic energy is zero. By the conservation of energy and mass principle, the energy before the collision is transformed into the mass of new particles formed in the collision. The particles that are present after a collision reaction can be different to those that went in. This is exactly what particle physicists aim to achieve, and the discovery and study of these new particles underpins their work.

Every collision is governed by one of the fundamental forces (except the force of gravity, which has no significant influence on such tiny particles in this context).


- The **electromagnetic force** leads to simple collisions between charged particles. No new particles are formed when this force is at work.



- The **strong force** dominates reactions between hadrons (which contain quarks).



- The **weak force** is likely to be involved in lepton reactions, especially if one of the leptons is a neutrino.



Einstein's theory of special relativity has led us to the idea that the mass of a moving object is not the same as its rest mass ( $m_0$ ).

The mass of a moving object cannot be measured directly; it must be calculated from a measurement of momentum and velocity. The relativistic equations for momentum and total energy can be found in the Formulae and Data Booklet. (Note: These equations are only applicable for non-zero mass.)

Relativity has also given us the idea of mass-energy equivalence. In Newton's version of mechanics, a lone particle not influenced by gravity or electromagnetism, but moving at a given speed, could only have a single form of energy, which is kinetic. At rest, it had no energy at all. This is not the case in relativity.

The relationship is described by the equation (EQ1) below.

$$E^2 - p^2 c^2 = m_0^2 c^4 \quad (\text{EQ1})$$

Photons are packets of energy travelling at the speed of light.

Surprisingly, it has been proved that although photons have zero mass, they do have momentum. Particle physics has also proven to be vital in understanding the nature of the universe a few fractions of a second after the Big Bang. The conditions created in the mightiest accelerators are very similar to those that existed when the universe was  $10^{12}$  seconds old.

- (a) In what sense can a particle be accelerated if its speed remains constant? Explain.  
[2 marks]

- If a particle undergoes circular motion or a change in direction. (1)
- A change in direction leads to a change in velocity and  $a = \frac{\Delta v}{t}$ . (1)

- (b) Once a charged particle has been accelerated to a given speed in a circular path, is further energy required to maintain a constant speed? Explain. [2 marks]

- Yes (1)
- It radiates energy as it moves (electromagnetic). (1)

- (c) Can electrons and neutrinos be subject to the strong force? Explain. [2 marks]

- No (1)
- Strong nuclear force only acts between hadrons, nucleons and quarks, not leptons. (1)

- (d) If neutrinos are involved in a collision reaction, why is it unlikely that this was governed by the electromagnetic force? [1 mark]

- Either acceptable (1)*
- Neutrinos have no charge.
  - Don't generate magnetic or electric fields so they are not influenced by electromagnetic force.

- (e) If you hit a ping pong ball with a table tennis bat, which of the three fundamental forces described governs this collision? Justify your answer. [2 marks]

- Electromagnetic force acts within a nucleus (strong force)  $\left(\frac{1}{2}\right)$
- Weak force is involved between the charges in the bat and ball.  $\left(\frac{1}{2}\right)$

- (f) Calculate the momentum of a proton travelling at 95.0 % of the speed of light. The rest mass of a proton is given in the formula and constant sheet. [3 marks]

$$\begin{aligned}
 p &= \frac{mv}{\sqrt{1 - \frac{v^2}{c^2}}} \\
 &= \frac{(1.67 \times 10^{-27})(0.950 \times 3.00 \times 10^8)}{\sqrt{1 - \frac{(0.950c)^2}{c^2}}} = 1.52 \times 10^{-18} \text{ kg ms}^{-1}
 \end{aligned}$$

↑ (1)

- (g) Given the equation (EQ1) for Einstein's mass-energy equivalence, show that for a **particle at rest** the equation simplifies to  $E = kc^2$ , where  $k$  is a constant. State clearly the physical quantity that  $k$  represents. [2 marks]

$$E^2 - p^2 c^2 = m_0^2 c^4$$

If particle is at rest,  $p=0$ .

$$\therefore E^2 - 0 = m_0^2 c^4$$

$$\Rightarrow E = m_0 c^2 \text{ where } k = m_0 \text{ (rest mass)} \quad (1)$$

$$= kc^2 \quad (1)$$

- (h) From the equation EQ1, express the momentum of a photon with zero mass in terms of  $wE$ , where  $w$  is a constant. State clearly the physical quantity that  $w$  represents.

[2 marks]

$$E^2 - p^2 c^2 = m_0^2 c^4$$

$$\text{If } m_0 = 0: \quad E^2 = p^2 c^2$$

$$\Rightarrow p = \frac{E}{c}$$

$$= wE \quad (1) \text{ where } w = \frac{1}{c} \quad (1)$$

- (i) Calculate the momentum of a photon of 550 nm yellow light.

[2 marks]

$$\begin{aligned} p &= \frac{h}{\lambda} \\ &= \frac{6.63 \times 10^{-34}}{550 \times 10^{-9}} \quad (1) \\ &= 1.20 \times 10^{-27} \text{ kg ms}^{-1} \quad (1) \end{aligned}$$

## Question 20 [18 marks]

The Physics behind the AFP Crime Scene Investigation

Australian Federal Police have isolated ***an element*** found in a specimen at a crime scene. They suspect that the element may be sodium or potassium or unknown element **X** or **Y**, and have asked the forensic laboratory to run tests on the crime scene specimen to identify which one element it contains.

***The laboratory is able to ionise the element to give it a single positive charge.*** They then accelerate the ions through a potential difference **V** (measured in volts) and, by use of a velocity filter, are able to send ions that have reached their maximum kinetic energy into a mass spectrometer. When the ions enter the mass spectrometer, they are acted on by a uniform magnetic field and follow a semi-circular path.

Technicians conduct a series of tests and measure the radius **r** of circular motion for different values of potential difference used to accelerate the charged ions.




Table 1 (next page) shows the results obtained when the **magnetic flux density *B*** in the mass spectrometer was **fixed at  $3.50 \times 10^{-2} T$** .

Potential Difference V (V)	Radius of circular path r (m)	Radius squared $r^2$ (m <sup>2</sup> )
200	0.270 ± 0.014	0.0730 ± 0.007
400	0.370 ± 0.019	0.137 ± 0.014
600	0.490 ± 0.025	0.240 ± 0.024 (1)
800	0.530 ± 0.027	0.281 ± 0.028 (1)
1000	0.620 ± 0.031	0.384 ± 0.038 (1)
1200	0.670 ± 0.034	0.449 ± 0.045

Table 1

It can be shown by an equation on the Formulae and Data Booklet that the radius  $r$  of circular motion for an ion of mass  $m$  and charge  $q$  enters the mass spectrometer at speed  $v$  and being deflected by a magnetic field of flux density  $B$ .

- (a) Use the equation and other equations on the Formulae and Data Booklet that link the kinetic energy (in Joules) attained by a mass of charge  $q$  in a potential difference  $V$  to **derive or show** that:

$$r^2 = gV \quad \text{where } g \text{ is a constant.}$$


State clearly the physical interpretation (or quantities) that  $g$  represents.

[3 marks]

$$\begin{aligned} t &= \frac{mv}{qB} \\ \Rightarrow v^2 &= \frac{r^2 q^2 B^2}{m^2} \quad (1) \end{aligned}$$

$$\begin{aligned} W &= Vq = \frac{1}{2}mv^2 \\ \Rightarrow Vq &= \frac{mr^2 q^2 B^2}{2m^2} \quad (1) \\ \Rightarrow r^2 &= \frac{2mV}{q^2 B^2} \\ &= gV \quad \text{where } g = \frac{2m}{q^2 B^2} \quad (1) \end{aligned}$$

- (b) Complete Table 1 by filling in the values of radius squared  $r^2$  with the appropriate uncertainty range. Three values have been done for you. [3 marks]
- (c) Plot the data from Table 1 on the grid provided, demonstrating the relationship between radius squared  $r^2$  (on the vertical axis) and potential difference  $V$  (on the horizontal axis). Draw the line of best fit including error bars. [5 marks]



- (d) Calculate the gradient of your line of best fit from your graph, showing all working.

[3 marks]

$$\text{gradient} = \frac{\Delta r^2}{\Delta V}$$

$$= \frac{0.520 - 0.00}{1400 - 0.00} \quad (1)$$

$$= \frac{3.7 \times 10^{-4} \text{ m}^2 \text{ V}^{-1}}{(1)} \quad (1)$$

- (e) Use the value of the gradient that you obtained to determine the identity of the charged ion.  
[4 marks]

*[If you could not obtain a gradient from part (d), use the numerical value of  $8.00 \times 10^{-4}$  for gradient.]*

You may use the following data:

Mass of a potassium K<sup>+</sup> ion =  $6.49 \times 10^{-28}$  kg

Mass of sodium Na<sup>+</sup> ion =  $3.68 \times 10^{-26}$  kg

Mass of X<sup>+</sup> ion =  $7.84 \times 10^{-26}$  kg

Mass of Y<sup>+</sup> ion = not available

$$\text{gradient} = \frac{\Delta r^2}{\Delta V} = \frac{2m}{qB^2}$$

$$\Rightarrow m = \frac{\text{gradient} \times qB^2}{2} \quad (1)$$

$$= \frac{(3.7 \times 10^{-4})(1.60 \times 10^{-19})(3.50 \times 10^{-2})^2}{2} \quad (1)$$


$$= 3.6 \times 10^{-26} \text{ kg.} \quad (1)$$

∴ Particle is Na<sup>+</sup> (1)

[If gradient =  $8.0 \times 10^{-4}$ , m =  $7.8 \times 10^{-26}$  kg, ion is X<sup>+</sup>]

**ADDITIONAL WORKING SPACE**

**SPARE GRID**



**12 Physics ATAR 2017 Semester 2**  
**ADDITIONAL WORKING SPACE**

**12 Physics ATAR 2017 Semester 2**

**ADDITIONAL WORKING SPACE**