


Copyright for test papers and marking guides remains with *West Australian Test Papers*.
Test papers may only be reproduced within the purchasing school according to the advertised Conditions of Sale.
Test papers should be withdrawn after use and stored securely in the school until Friday 1st December 2017.



SEMESTER TWO

**MATHEMATICS
SPECIALIST
UNITS 1 & 2**

2017

SOLUTIONS

Calculator-free Solutions


1. (a) $1000 = 310 + 650 + 440 - 170 - 150 - 180 + x$ ✓

$x = 100$ ✓

(b) $150 - 100 = 50$ ✓

(c) (i) 44 ✓

(ii)



$\therefore n(B) = 34$ ✓✓

[6]

2. (a) (i) Substitute $z = 2i$ to get $(2i)^4 - 2(2i)^3 + 7(2i)^2 - 8(2i) + 12$
which reduces to 0 ✓

(ii) $z = -2i$ (the conjugate) is the other root. ✓

(b) $2x^2 + 10 = 3 - 5x$ reduces to $2x^2 + 5x + 7 = 0$ ✓

$\therefore x = \frac{-5 \pm i\sqrt{31}}{4}$ from quadratic formula ✓✓

[5]

3. (a) (i) ${}^5C_2 = {}^5C_3 = 10$ This statement is true. ✓✓

(ii) ${}^5C_1 \neq 2 \times {}^5C_0$ This statement is false ✓✓


(b) (i) ${}^5C_3 = 10$ ✓✓

(ii) $2 \times 4! = 48$ ✓✓

[8]

4. (a) $p = 4, q = 0.2$ ✓✓
 (b) $y = 1 - x$ becomes $y = 4[1 - 0.2x]$ ✓
 i.e. $y = 4 - 0.8x$
 or, if (c) is done before (b), gradient is -0.8 and intercept is 4
 $\therefore y = 4 - 0.8x$
 (c) $A' = (5, 0)$ and $B' = (0, 4)$ ✓✓
 $\begin{bmatrix} 5 & 0 \\ 0 & 4 \end{bmatrix}$
 (d) ✓✓
 $\begin{bmatrix} 0.2 & 0 \\ 0 & 0.25 \end{bmatrix}$
 (e) ✓✓
 (f) Reflection across y axis
 i.e. $g(x)$ becomes $-g(x)$ ✓✓
 (g) $A = 0.5(ms - rn)$ ✓ [12]

5. (a)



- (b) $x \approx \frac{11\pi}{12}$ line ✓ ✓✓
 of intersection and accuracy ✓✓
 (c) $\sin x$ ✓✓
 (d) $2\cos x \cdot \sin x = \sin 2x$ ✓✓ [8]

6. (a) Let the numbers be $2k - 1, 2k + 1, 2k + 3, 2k + 5, 2k + 7$ ✓
 $2k - 1 + 2k + 1 + 2k + 3 + 2k + 5 + 2k + 7$
 $= 10k + 15$ ✓
 Since $10k + 15 = 5(2k + 3)$ then divisible by 5. ✓

- (b) Assume that $-\pi$ is rational, hence $-\pi = \frac{a}{b}$ ✓

$$\therefore \pi = -\frac{a}{b} = \frac{-a}{b}$$

But $-a$ and b are integers, so $-\pi$ is rational. ✓

This contradicts the supposition, and

therefore by contradiction $-\pi$ must be irrational. ✓ [7]

7. For $n = 1$:

$$\frac{1 - x^1}{(1 - x)} = 1 \quad \therefore \text{True for } n = 1$$

✓

Assume true for $n = k$:

$$\text{ie. } 1 + x + x^2 + \dots x^{k-1} = \frac{1 - x^k}{(1 - x)}$$

✓

Prove true for $n = k + 1$:

$$\text{Proof: } 1 + x + x^2 + \dots x^{(k+1)-1} = \frac{1 - x^k}{(1 - x)} + x^{(k+1)-1}$$

$$1 + x + x^2 + \dots x^k = \frac{(1 - x^k) + x^k(1 - x)}{(1 - x)}$$

✓

$$1 + x + x^2 + \dots x^k = \frac{1 - (x^{k+1})}{(1 - x)} \quad \text{as required}$$

✓

Therefore, True for $n = k + 1$, and since true for $n = 1$,
true for all whole numbers.

✓

[5]

Calculator-assumed Solutions

8. $wz = (2 + ai)(3b + i) = 4$

$$\therefore 6b + 2i + 3abi - a = 4$$

✓

$$\therefore 6b - a = 4 \text{ and } 2 + 3ab = 0$$

✓

$$\therefore 2 + 3(6b - 4)(b) = 0$$

✓

$$\therefore 9b^2 - 6b + 1 = 0$$

✓

$$\therefore b = \frac{1}{3} \text{ and } a = -2$$

✓

[5]

$$\begin{aligned}
 & \frac{1 + \frac{\sin x}{\cos x} \cdot \frac{\sin y}{\cos y}}{\frac{\sin x}{\cos x} + \frac{\sin y}{\cos y}} \\
 9. \quad (a) \quad \text{RHS} &= \frac{\cos x \cos y + \sin x \sin y}{\cos x \cos y} \div \frac{\sin x \cos y + \sin y \cos x}{\cos x \cos y} \\
 &= \frac{\cos(x-y)}{\sin(x+y)} = \text{LHS}
 \end{aligned}$$

$$(b) \quad (i) \quad \mathbf{p} \cdot \mathbf{q} = |\mathbf{p}||\mathbf{q}|\cos(A-B)$$

$$\text{and } \mathbf{p} \cdot \mathbf{q} = (|\mathbf{p}|\cos A \mathbf{i} + |\mathbf{p}|\sin A \mathbf{j}) \cdot (|\mathbf{q}|\cos B \mathbf{i} + |\mathbf{q}|\sin B \mathbf{j})$$

$$= |\mathbf{p}||\mathbf{q}|(\cos A \mathbf{i} + \sin A \mathbf{j}) \cdot (\cos B \mathbf{i} + \sin B \mathbf{j})$$

$$= |\mathbf{p}||\mathbf{q}|[\cos A \cos B + \sin A \sin B]$$

$$\therefore \cos(A-B) = \cos A \cos B + \sin A \sin B$$

$$(ii) \quad \cos(A+B) = \cos(A-(-B)) = \cos A \cos(-B) + \sin A \sin(-B)$$

$$= \cos A \cos B - \sin A \sin B$$

$$(iii) \quad \cos^2 A + [\cos(120^\circ + A)]^2 + [\cos(120^\circ - A)]^2$$

$$= \cos^2 A + [\cos 120^\circ \cos A - \sin 120^\circ \sin A]^2 + [\cos 120^\circ \cos A + \sin 120^\circ \sin A]^2$$

$A]^2$

$$= \cos^2 A + \left[-\frac{\cos A}{2} - \left(\frac{\sqrt{3}}{2}\right) \sin A \right]^2 + \left[-\frac{\cos A}{2} + \left(\frac{\sqrt{3}}{2}\right) \sin A \right]^2$$

$$= 1.5 \cos^2 A + 1.5 \sin^2 A$$

$$= 1.5 \quad [13]$$

$$10. \quad (a) \quad (3\mathbf{a} - \mathbf{b}) \cdot (\mathbf{a} + 3\mathbf{b}) = 3\mathbf{a} \cdot \mathbf{a} - \mathbf{b} \cdot \mathbf{a} + 9\mathbf{a} \cdot \mathbf{b} - 3\mathbf{b} \cdot \mathbf{b}$$

$$= 3|\mathbf{a}|^2 + 8\mathbf{a} \cdot \mathbf{b} - 3|\mathbf{b}|^2$$

$$= 3 + 8|\mathbf{a}||\mathbf{b}|\cos \theta - 3$$

$$= 8\cos \theta \text{ as required}$$

$$(b) \quad \overrightarrow{PQ} = (4, 1)$$

$$\text{Unit vector on } x \text{ axis} = (1, 0)$$

$$\text{Length of projection} = |\overrightarrow{PQ} \cdot \hat{x}| = |4| = 4 \quad [7]$$

$$11. \quad (a) \quad \text{Let the cost of a bottle of orange concentrate cost } x$$

$$\text{Let the cost of a bottle of banana concentrate cost } y$$

$$5x + 1y = 19$$

$$2x + 3y = 18$$

$$(b) \quad \begin{bmatrix} 5 & 1 \\ 2 & 3 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 19 \\ 18 \end{bmatrix}$$

$$\therefore \begin{pmatrix} x \\ y \end{pmatrix} = \begin{bmatrix} 5 & 1 \\ 2 & 3 \end{bmatrix}^{-1} \begin{bmatrix} 19 \\ 18 \end{bmatrix}$$


✓

$$= \frac{1}{13} \begin{bmatrix} 3 & -1 \\ -2 & 5 \end{bmatrix} \begin{pmatrix} 19 \\ 18 \end{pmatrix} = \begin{pmatrix} 3 \\ 4 \end{pmatrix}$$

✓

[4]

12. (a)



✓✓

$$(b) \begin{bmatrix} 0 \\ -30 \end{bmatrix} + \begin{bmatrix} 60\cos\theta \\ 60\sin\theta \end{bmatrix} = \begin{bmatrix} x \\ 0 \end{bmatrix}$$

✓✓

$$\therefore 60\sin\theta = 30$$

✓

$$\therefore \sin\theta = \frac{1}{2}$$

$$\therefore \theta = 30^\circ \rightarrow \text{Bearing is } 060^\circ \text{ T}$$

✓

$$(c) \text{ Speed in Easterly direction is } 60\cos 30^\circ = 60 \times \frac{\sqrt{3}}{2}$$

✓

$$\text{Time taken is } \frac{8}{60 \times \frac{\sqrt{3}}{2}} = 9.23 \text{ minutes}$$

✓

[8]

$$13. (a) (i) \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix} \therefore \text{Rotation of } 180^\circ$$

✓✓

$$(ii) \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \therefore \text{Rotation of } 270^\circ \text{ clockwise}$$

✓✓

$$(b) P = 6(B - 2A) \times B^{-1}$$

✓

$$\therefore P = \frac{3}{11} \begin{bmatrix} 24 & -12 \\ 8 & 18 \end{bmatrix}$$

✓✓

$$(c) BA = \begin{bmatrix} -2 & 6 \\ -4 & 1 \end{bmatrix}$$

$$BAX = \begin{bmatrix} -2 & 6 \\ -4 & 1 \end{bmatrix} \begin{bmatrix} 2 \\ 3 \end{bmatrix} = \begin{bmatrix} 14 \\ -5 \end{bmatrix}$$

✓

Co-ordinates are (14, -5)

✓

$$(d) \text{Det } B = 22 \therefore \text{Area} = 25 \times 22 = 550$$

✓✓

$$(e) \text{Singular matrix has det} = 0$$

✓

$$\therefore \text{Area} = 0 \text{ i.e. A line}$$

✓

[13]

14. (a) $3(2\mathbf{i} + 3\mathbf{j}) - (m\mathbf{i} - 5\mathbf{j}) = 8\mathbf{i} + 14\mathbf{j}$ ✓

$$\therefore (6 - m)\mathbf{i} + 14\mathbf{j} = 8\mathbf{i} + 14\mathbf{j}$$

$$\therefore 6 - m = 8$$

$$\therefore m = -2$$
 ✓

(b) $2\mathbf{i} + 3\mathbf{j} = k(m\mathbf{i} - 5\mathbf{j})$ ✓

$$\therefore 2 = km \text{ and } 3 = -5k$$

$$\therefore k = -0.6 \text{ and by substitution, } m = -\frac{10}{3}$$
 ✓

(c) $\begin{pmatrix} 2 \\ 3 \end{pmatrix} \cdot \begin{pmatrix} m \\ -5 \end{pmatrix} = 0$ ✓

$$\therefore 2m - 15 = 0$$

$$\therefore m = 7.5$$
 ✓

[6]

15. (a) $R \cos(A - \theta) = R \cos(A) \cos(\theta) + R \sin(A) \sin(\theta)$

$$= -3 \cos(A) + 3\sqrt{3} \sin(A)$$

$$\therefore R \sin(\theta) = 3\sqrt{3} \text{ and } R \cos(\theta) = -3$$

hence, $R^2 = (-3)^2 + (3\sqrt{3})^2 = 36 \therefore R = 6$ ✓✓

and $\cos(\theta) = \frac{-3}{6} = -\frac{1}{2} \therefore \theta = \frac{2\pi}{3}$ ✓

therefore, $R \cos(A - \theta) = 6 \cos\left(A - \frac{2\pi}{3}\right)$ ✓


(b) (i) $g(x)_{\min} = -6$ ✓

(ii) for $\cos\left(A - \frac{2\pi}{3}\right) = -1$

hence $A - \frac{2\pi}{3} = \pi \therefore \theta = \frac{5\pi}{3}$ ✓✓

[7]


16. (a)



$\theta = 60^\circ$ (equilateral triangle) ✓

$\therefore \alpha = 30^\circ$ (central angle theorem) ✓

(b) (i)



$AOB = 60^\circ$ (proved in (a))

$\therefore ACB = 30^\circ$ (theorem)

Similarly, $DBC = 30^\circ$

$\therefore BEC = 120^\circ$

$\therefore AEB = 60^\circ$ ✓✓✓

(ii) Assume E is the centre.

All angles of $\triangle ABE = 60^\circ$

and all angles of $\triangle BEC = 60^\circ$

But $\angle AEB = 2\angle ACB$ which is impossible if they are both 60°

\therefore E is not the centre.

✓✓✓

[8]

17.


$$z = 2 + 3i$$

(i) $2 - 3i$ ✓✓

(ii) $-3 + 2i$ ✓✓

(iii) $-5 + 12i$ ✓✓

(iv) 13 ✓✓



18. (a) 2m

(b) $\cos \frac{\pi t}{15} = 1 \rightarrow \frac{\pi t}{15} = 2\pi$

$\therefore t = 30$ secs

(c) 138m

(d) $-68 \cos \frac{\pi t}{15} + 70 = 100$

$\therefore t = 9.68, 20.32$

$\therefore 10.64$ minutes

✓

✓

✓

✓

✓

✓

✓

[8]

[7]

19. (a) It is given that $(A + B)^2 = A^2 + BA + AB + B^2$

Since $AB \neq BA$,

$\therefore (A + B)^2 \neq A^2 + 2AB + B^2$

(b) $AB = BC$

$\therefore AAB = ABC$

$\therefore A^2B = BC^2$

$\therefore AA^2B = ABC^2$

$\therefore A^3B = BC^3$

$\therefore A^3BB^{-1} = BC^3B^{-1}$

✓

✓

✓

✓

✓

$\therefore A^3 = BC^3B^{-1}$ as required

✓

[6]

20. (a) $\overrightarrow{OB} = 4i + 4j$ ✓
 $\overrightarrow{CA} = 4i - 4j$ ✓
- (b) $\overrightarrow{CA} \cdot \overrightarrow{OB} = (4i + 4j) \cdot (4i - 4j) = 0$ ✓
 $\therefore \overrightarrow{CA} \perp \overrightarrow{OB}$ ✓
- (c) Let k be the midpoint of \overrightarrow{OB} .
Then $K = (2, 2)$
 $\therefore \overrightarrow{OK} = 2i + 2j$ ✓
So $\overrightarrow{CK} = \overrightarrow{KO} + \overrightarrow{OC} = -(2i + 2j) + 4j = -2i + 2j$ ✓
But $\overrightarrow{CA} = -4i + 4j = 2\overrightarrow{CK}$
 $\therefore K$ is the midpoint of \overrightarrow{CA} ✓
 \therefore Diagonals bisect each other. [7]