

# ATAR Course Examination, 2019

## Question/Answer Booklet

### PHYSICS

Name: SOLUTIONS

Teacher: \_\_\_\_\_

#### Time allowed for this paper

Reading time before commencing work: ten minutes  
Working time for paper: three hours

#### Material required/recommended for this paper

##### *To be provided by the supervisor*

This Question/Answer booklet  
Formula and Data booklet

##### *To be provided by the candidate*

Standard Items: pens (blue/black preferred), pencils (including coloured), sharpener, correction fluid/tape, eraser, ruler, highlighters

Special Items: non-programmable calculators satisfying the conditions set by the School Curriculum and Standards Authority for this course, drawing templates, drawing compass and a protractor

#### Important note to candidates

No other items may be taken into the examination room. It is **your** responsibility to ensure that you do not have any unauthorised notes or other items of a non-personal nature in the examination room. If you have any unauthorised material with you, hand it to the supervisor **before** reading any further.

**Structure of this paper**

Section	Number of questions available	Number of questions to be answered	Suggested working time (minutes)	Marks available	Percentage of examination
Section One Short response	11	11	50	49	27
Section Two Problem-solving	7	7	90	99	54
Section Three Comprehension	2	2	40	36	19
<b>Total</b>					<b>100</b>

**Instructions to candidates**

1. The rules for the conduct of Western Australian external examinations are detailed in the *Year 12 Information Handbook 2019*. Sitting this examination implies that you agree to abide by these rules.
2. Write answers in this Question/Answer Booklet.
3. When calculating or estimating answers, show all your working clearly. Your working should be in sufficient detail to allow your answers to be checked readily and for marks to be awarded for reasoning.

In calculations, give final answers to three significant figures and include appropriate units where applicable.

In estimates, give final answers to a maximum of two significant figures and include appropriate units where applicable.

4. You must be careful to confine your responses to the specific questions asked and to follow any instructions that are specific to a particular question.
5. Supplementary pages for the use of planning/continuing your answer to a question have been provided at the end of this Question/Answer booklet. If you use these pages to continue an answer, indicate at the original answer where the answer is continued, i.e. give the page number.
6. The Formulae and Data booklet is not to be handed in with your Question/Answer booklet.

**SECTION ONE: Short Response****30% (49 marks)**

This section has **11 questions**. Answer **all** questions. Write your answers in the spaces provided.

When calculating numerical answers, show your working or reasoning clearly. Give final answers to **three** significant figures and include appropriate units where applicable.

When estimating numerical answers, show your working or reasoning clearly. Give final answers to a maximum of **two** significant figures and include appropriate units where applicable.

Supplementary pages for the use of planning/continuing your answer to a question have been provided at the end of this Question/Answer booklet. If you use these pages to continue an answer, indicate at the original answer where the answer is continued, i.e. give the page number.

Suggested working time: 50 minutes.

**Question 1****(4 marks)**

Mars' mass is  $6.39 \times 10^{23}$  kg and has an orbital radius around the Sun of 228 million kilometres. Calculate the weakest gravitation force that can act between Earth and Mars, assuming both Earth and Mars have circular orbits.

$$\textcircled{M} \quad \textcircled{S} \quad \textcircled{E} \\ 2.28 \times 10^9 \text{ m} \rightarrow 2.150 \times 10^{12} \text{ m} \rightarrow$$

*Earth & Mars on opposite sides*

$$\begin{aligned} G &= \frac{GM_E M_m}{r^2} \quad (1) \\ &= \frac{(6.67 \times 10^{-11})(5.97 \times 10^{24})(6.39 \times 10^{23})}{(2.28 \times 10^9 + 1.50 \times 10^{12})^2} \quad (1) \\ &= \underline{\underline{1.78 \times 10^{15} \text{ N}}} \quad (1) \end{aligned}$$

Answer: 1.78  $\times 10^{15}$  N

**Question 2**

(4 marks)

An electron with 2.80 eV of kinetic energy bombards an atom with a single ground state electron. The atom's electron is excited and later transitions back to the ground state, emitting a single 518 nm photon. Calculate the kinetic energy of the bombarding electron after it scattered off the atom.


$$\begin{aligned} E_{\text{photon}} &= \frac{hc}{\lambda} \\ &= \frac{(6.63 \times 10^{-34})(3.00 \times 10^8)}{(518 \times 10^{-9})} \quad (1) \\ &= 3.84 \times 10^{-19} \text{ J} \quad (1) \\ &= 2.40 \text{ eV.} \quad (1) \end{aligned}$$

$$\begin{aligned} E(\text{scattered } e^-) &= 2.80 - 2.40 \\ &= 0.40 \text{ eV} \quad (1) \end{aligned}$$

Answer: 0.40 eV**Question 3**

(4 marks)

A 30.0 g golf ball at a mini-golf course approaches a small vertical loop obstacle at  $3.45 \text{ ms}^{-1}$ . The ball follows the track, completing the vertical loop.



Calculate the magnitude of the reaction force applied to the ball by the track when the ball is at the top of the loop.

$$\begin{aligned} E_p + E_k(\text{top}) &= E_k(\text{bottom}) \\ \Rightarrow gh + \frac{1}{2}mv^2 &= \frac{1}{2}mv^2 \quad (1) \\ \Rightarrow v &= \sqrt{2\left(\frac{v^2}{2} - gh\right)} \\ &= \sqrt{2\left[\frac{(3.45)^2}{2} - (9.80)(0.400)\right]} \\ &= 2.02 \text{ ms}^{-1} \quad (1) \end{aligned}$$


$$\begin{aligned} \sum F &= F_c = F_w + R \\ \Rightarrow R &= F_c - F_w \\ &= \frac{mv^2}{r} - mg \\ &= \frac{(0.0300)(2.02)^2}{(0.200)} - (0.0300)(9.80) \\ &= 0.318 \text{ N down} \quad (1) \end{aligned}$$

Answer: 0.318 N

**Question 4**

(4 marks)

The diagram below shows an electron entering a uniform  $2.00 \text{ NC}^{-1}$  electric field. There is also a magnetic field in this region (not shown on the diagram).



The electron has a constant velocity of  $8540 \text{ ms}^{-1}$  while in the presence of the two fields. State the direction of the magnetic field and calculate its strength.

- $B$  is into the page.

$$F_B = F_E \quad (1)$$

$$\Rightarrow qvB = Eq \quad (1)$$

$$B = \frac{E}{v} \quad (1)$$

$$= \frac{2.00}{8540}$$


$$= 2.34 \times 10^{-4} \text{ T} \quad (1)$$

Direction: into page (1) Strength:  $2.34 \times 10^{-4}$  T

**Question 5**

(4 marks)

The following particle reaction is proposed by a PhD student while studying new, exotic particles of the standard model.



Justify whether this reaction is possible based on baryon number and electric charge.

	LHS	RHS	
charge	$\frac{2}{3} + \left(-\frac{1}{3}\right) + \left(-\frac{1}{3}\right) \\ = 0 \quad (1)$	$\frac{2}{3} - \frac{2}{3} + \left(-\frac{1}{3}\right) + \left(-\frac{2}{3}\right) + \frac{2}{3} + \frac{2}{3} + \left(-\frac{1}{3}\right) \\ = 0 - 1 + 1 = 0 \quad (1)$	$\therefore$ Possible for both charge and baryon number.
baryon	$\frac{1}{3} + \frac{1}{3} + \frac{1}{3} \\ = +1 \quad (1)$	$\frac{1}{3} - \frac{1}{3} + \frac{1}{3} - \frac{1}{3} + \frac{1}{3} + \frac{1}{3} + \frac{1}{3} \\ = +1 \quad (1)$	


**Question 6**

(4 marks)

A square coil moves into a uniform 0.260 T magnetic field which is aligned perpendicular to the area of the coil. The coil is induced with a 0.650 V emf as it enters the field at 4.75 ms<sup>-1</sup>. For what amount of time does the coil have an induced emf?

*Consider the leading edge.*

$$\begin{aligned} \mathcal{E} &= Blv \\ \Rightarrow l &= \frac{\mathcal{E}}{Bl} \quad (1) \\ &= \frac{0.650}{(0.260)(4.75)} \\ &= 0.526 \text{ m} \quad (1) \end{aligned}$$



X X X  
X X X  
X X X  
X X X

*Calculate time t to enter the field.*

$$\begin{aligned} v &= \frac{s}{t} \\ \Rightarrow t &= \frac{s}{v} = \frac{0.526}{4.75} = 0.111 \text{ s} \quad (1) \end{aligned}$$

Answer: 0.111 s

**Question 7**

(4 marks)

The red-shift of light from galaxies not our own is supporting evidence of the Big Bang Theory. Describe what causes the increasing amount of red-shift of light from galaxies further away and why **only** nearby galaxies may have blue-shifted light.

- Red-shift is caused by the expansion of space, causing the wavelength to stretch. [Could also mention recessional velocities as measured from Earth.]
- Light from distant galaxies is stretched more due to the increased time spent in expanding space. (1)
- The space closest to us is not expanding greatly. (1)
- Close galaxies may have been moving towards us. (1)  
[Could also mention localised gravitational effects are pulling neighbouring galaxies towards us.]


**Question 8**

(5 marks)

Laptop chargers are traditional transformers that create an alternating current in the secondary coil.



The circuit diagram depicting the above situation is shown below.




- (a) Explain briefly how the transformer creates an alternating current in the secondary coil.
- An alternating current is applied to the primary coil. (1)
  - This creates a changing magnetic field in the primary coil. (1)
  - The secondary coil experiences a changing magnetic field, inducing a current in the secondary coil. (1)
- (b) Using information from the diagram, calculate the potential difference that the charger provides to the laptop. (2 marks)

$$\begin{aligned} \frac{V_p}{V_s} &= \frac{N_p}{N_s} \\ \Rightarrow V_s &= \frac{V_p N_s}{N_p} \quad (1) \\ &= \frac{(240)(65)}{600} \\ &= \underline{\underline{26.0 \text{ V}}} \quad (1) \end{aligned}$$

Answer 26.0 V

**Question 9****(5 marks)**

The photoelectric effect can be demonstrated by illuminating a negatively-charged electroscope, with ultraviolet (UV) light. A particular electroscope has a zinc metal plate. Once the electroscope was charged, a student switched on an ultraviolet light. After a few seconds, the electroscope was discharged as electrons were emitted from the surface of the zinc.



- (a) Would a positively charged electroscope have been discharged in the same way? Explain. (2 marks)

- No (1)
- Only electrons (negative particles) are emitted in the photoelectric effect. (1)

- (b) Calculate the velocity of the emitted electrons, if the UV light has a frequency of  $1.20 \times 10^{15}$  Hz. The work function of zinc is 4.30 eV. (3 marks)



$$\begin{aligned} E &= hf = \phi + E_k(\text{max}) \quad (1) \\ \Rightarrow (6.63 \times 10^{-34})(1.20 \times 10^{15}) &= (4.30)(1.60 \times 10^{-19}) + \frac{1}{2}(9.1 \times 10^{-31})v^2 \quad (1) \\ \Rightarrow v &= 4.86 \times 10^5 \text{ ms}^{-1} \quad (1) \end{aligned}$$

Answer  $4.86 \times 10^5$   $\text{ms}^{-1}$

**Question 10**

(6 marks)

The figure on the left below shows the side view of an electron diffraction tube used to demonstrate the wave properties of an electron. An electron beam is incident on a thin graphite target that behaves like the slits in a diffraction grating experiment. After passing through the graphite target the electrons strike a fluorescent screen. The figure on the right below shows the appearance of the fluorescent screen when the electrons are incident on it.



- (a) Explain how the pattern produced on the screen supports the idea that the electron beam is behaving as a wave rather than as a stream of particles. (2 marks)

- Interference patterns are seen on the fluorescent screen - this shows wave behaviour. (1)
- If electrons were acting as particles, they would clump together on the screen and not interfere. (1)

- (b) When the electrons strike the graphite target, they have a speed of  $2.20 \times 10^7 \text{ ms}^{-1}$ .

- (i) Calculate the potential difference required to accelerate the electrons to this speed. (2 marks)

$$\begin{aligned} W &= Vq = \frac{1}{2}mv^2 \\ \Rightarrow V &= \frac{mv^2}{2q} = \frac{(9.11 \times 10^{-31})(2.20 \times 10^7)^2}{2(1.60 \times 10^{-19})} \quad (1) \\ &= 1.38 \times 10^3 \text{ V} \quad (1) \end{aligned}$$

Answer  $1.38 \times 10^3$  V

- (ii) Calculate the de Broglie wavelength of the electrons as they strike the graphite. (2 marks)


$$\begin{aligned} \lambda &= \frac{h}{mv} \\ &= \frac{(6.63 \times 10^{-34})}{(9.11 \times 10^{-31})(2.20 \times 10^7)} \quad (1) \\ &= \underline{3.31 \times 10^{-11} \text{ m}} \quad (1) \end{aligned}$$

Answer  $3.31 \times 10^{-11}$  m

## Question 11

(5 marks)

During a lightning strike, there is a **negative** discharge from a cloud to the ground. This discharge produces a 325 kA current that falls to 0.00 A in 50.0  $\mu$ s. There is a 55 turn coil of radius 0.800 m placed 225 m from the strike as shown below.



\* diagram not to scale

- (a) Calculate the average emf induced in the coil during this strike. (4 marks)

$$\begin{aligned} B &= \frac{\mu_0 I}{2\pi r} \\ &= \frac{(4\pi \times 10^{-7})(325 \times 10^3)}{2\pi (225)} \quad (1) \\ &= 2.89 \times 10^{-4} \text{ T} \quad (1) \end{aligned}$$

$$\begin{aligned} \mathcal{E} &= -\frac{N \Delta \Phi}{\Delta t} = -\frac{N \Delta B A}{\Delta t} \\ &= -\frac{(55)(0.289 \times 10^{-4}) \pi (0.800)^2}{50.0 \times 10^{-6}} \quad (1) \\ &= \underline{639 \text{ V}} \quad (1) \end{aligned}$$

DO NOT WRITE IN THIS AREA AS IT WILL BE CUT OFF

Answer 639 V

- (b) On the diagram indicate the direction of the induced emf in the coil. (1 mark)

• Anticlockwise (1)

**End of Section One**

See next page

**SECTION TWO: Problem-solving****50% (99 marks)**

This section has **six** questions. Answer **all** questions. Write your answers in the spaces provided.

When calculating numerical answers, show your working or reasoning clearly. Give final answers to **three** significant figures and include appropriate units where applicable.

When estimating numerical answers, show your working or reasoning clearly. Give final answers to a maximum of **two** significant figures and include appropriate units where applicable.


Supplementary pages for the use of planning/continuing your answer to a question have been provided at the end of this Question/Answer booklet. If you use these pages to continue an answer, indicate at the original answer where the answer is continued, i.e. give the page number.

Suggested working time: 90 minutes.

---

**Question B-1****(14 marks)**

A gantry crane is being used to lift a 2.50 tonne crate as shown in the diagram below. The gantry consists of a 4.32 tonne, 21.0 m uniform beam that is supported by two 10.0 m support columns A and B. The 2.50 tonne crate is 4.50 m from the center of column A and 16.50 m from column B. There is a 750 kg winch whose center of mass is directly above the center of mass the crate.




- (a) In the space below, draw a free body diagram of the beam, clearly labelling all forces acting on it. (4 marks)

- (b) Calculate the reaction force provided by each column on the beam. (4 marks)

Reaction force from column A: \_\_\_\_\_ N

Reaction force from column B: \_\_\_\_\_ N

In a variation of the gantry crane, the beam is supported by one column, pivoted at its base and at its point of attachment with the beam, which is held in place by a steel cable attached to a large concrete block. The beam is supported by a 250 kg strut CD which is pivoted at both ends. CD makes an angle of  $60^\circ$  with the beam and is attached 11.0 m from the right hand end of the beam. The cable makes an angle of  $40^\circ$  with the support column.



- (c) Calculate the tension in the steel cable. (3 marks)

DO NOT WRITE IN THIS AREA AS IT WILL BE CUT OFF

Answer: \_\_\_\_\_ N

- (d) Calculate the force of compression in strut CD.

(3 marks)

Answer: \_\_\_\_\_ N

DO NOT WRITE IN THIS AREA AS IT WILL BE CUT OFF

**Question B-2****(12 marks)**

Claire is standing on Earth. She observes Jim passing by in a spaceship at  $0.60 c$ . Jim observes the spaceship to be 18.0 m long. Jim is playing hyperspace pong where he hits a ball towards the front of the spaceship from the back at  $0.40 c$  (according to Jim). The ball has a rest mass of 0.500 kg.

- (a) What time does Jim observe the ball take to reach the front of the spaceship?

(2

marks)

Answer: \_\_\_\_\_ s

- (b) As the ball completes the journey towards the front of the spaceship, does Jim observe the proper length of the ball's journey or the proper time for the ball's journey or both? Justify your choice.

(2 marks)

- (c) How long is the spaceship as measured by Claire?

(2 marks)

DO NOT WRITE IN THIS AREA AS IT WILL BE CUT OFF

Answer: \_\_\_\_\_ m

- (d) What is the velocity of the ball as measured by Claire? Give your answer as a fraction of the speed of light. (2 marks)

Answer: \_\_\_\_\_ c

- (e) Calculate the energy of the ball as measured by Jim. (2 marks)


Answer: \_\_\_\_\_ J

- (f) Calculate the momentum of the ball as measured by Claire. (2 marks)

Answer:  $\text{kg m s}^{-1}$   
**(12 marks)**

**Question B-3**

A trebuchet is a siege weapon that flings boulders from a great distance. Consider the arrangement of a trebuchet and a castle shown below.



- (a) The boulder lands at the same height it was launched from, was fired at  $45.0^\circ$  above the horizontal and was airborne for 8.16 s. Complete the following questions:

- i. Calculate the launch velocity of the boulder. (3 marks)

Answer: \_\_\_\_\_  $\text{m s}^{-1}$


See next page

- ii. Calculate the maximum height the boulder achieved above its launch point.  
(3 marks)

Answer: \_\_\_\_\_  $\text{m s}^{-1}$

DO NOT WRITE IN THIS AREA AS IT WILL BE CUT OFF

Medieval castles were often built at higher elevations to give an advantage to those under siege.




- (b) A launched boulder is in the air for 4.80 s. The distances,  $s_h$  and  $s_v$  above indicate how far the boulder travelled to hit the castle. Determine both the speed and angle above the horizon the boulder was launched at. You may make use of the trigonometric identity  $\tan\theta = \frac{\sin\theta}{\cos\theta}$  and air resistance can be ignored. (6 marks)

DO NOT WRITE IN THIS AREA AS IT WILL BE CUT OFF

Speed: \_\_\_\_\_ m s<sup>-1</sup> Angle: \_\_\_\_\_ °**Question B-4****(17 marks)**

A small rectangular coil ABCD contains 140 turns of wire. The sides AB and BC of the coil are of lengths 4.5 cm and 2.8 cm respectively, as shown in the figure below.



The coil is held between the poles of a large magnet so that the coil can rotate about an axis through its centre. When the current in the coil is 170 mA and the coil is stationary, the maximum torque produced in the coil is  $2.1 \times 10^{-3}$  N m.

- (a) Is the position shown, one of minimum or maximum torque? Explain. (2 marks)

---



---



---



---

- (b) For the coil in the position shown calculate the magnitude of the force on

- (i) side AB of the coil (2 marks)

Answer \_\_\_\_\_ N


- (ii) side BC of the coil (2 marks)

Answer \_\_\_\_\_ N


- (c) Calculate the strength of the magnetic field experienced by the sides of the coil. (4 marks)

Answer: \_\_\_\_\_

- (d) The above diagram does not show how the coil is connected to a potential difference. Of the two mechanisms shown below, which mechanism should be used for the coil to rotate as a DC motor. Name the mechanism and explain your choice. (3 marks)



Mechanism 1



Mechanism 2

- (e) Once the coil has started rotating as a DC motor, does the maximum torque increase, decrease or remain the same. Explain. (4 marks)

---


---

---

---

**Question B-5** (16 marks)

During an experiment, a pendulum is set up, as shown in the diagram. The length of the cord attached to the bob is 1.55 m. The bob has a mass of 1.80 kg and is released from rest from the position shown. At the lowest point of its path, the bob is 7.80 cm beneath its starting point.




- (a) By considering conservation of energy, calculate the velocity of the bob at its lowest point.  
(3 marks)

Answer = \_\_\_\_\_  $\text{m s}^{-1}$

- (b) Calculate the tension in the cord at the lowest point of its path.  
(3 marks)

Answer = \_\_\_\_\_ N

Later, the experimental setup is modified so that the bob swings in a horizontal circular path, with radius 0.290 m, as a conical pendulum.



- (c) On the above diagram, indicate all forces acting on the bob as clearly labelled arrows **and** indicate the direction of the net force on the bob as a dashed arrow (---→). (3 marks)
- (d) Show that the tension in the cord is now 18.0 N. (4 marks)
- (e) Calculate the magnitude of the velocity of the bob at the position shown. (3 marks)

DO NOT WRITE IN THIS AREA AS IT WILL BE CUT OFF

Answer \_\_\_\_\_ m s<sup>-1</sup>


DO NOT WRITE IN THIS AREA AS IT WILL BE CUT OFF

---

**Question B-6****(16 marks)**

Digital television in Australia can be accessed by using a satellite dish pointed at a satellite in space. The satellite used to transmit the signals appears to stay still relative to the Earth. The satellite, with a mass of 300 kg, is actually travelling around the Earth in a geostationary orbit.

The picture below show the three main types of satellite orbits. Low Earth Orbits (LEO), Medium Earth Orbits (MEO) and Geostationary Earth Orbits (GEO).



- (a) In the picture, there is an error with the indicated orbit a GEO satellite. Indicate this error and explain why the orbit shown must be an error. (3 marks)

---

---

---

- (b) Which of these satellites experiences the greatest gravitational force from the Earth? Circle the correct answer from the choices below. Explain your answer in the space provided. (3 marks)

LEO      MEO      GEO      All satellites experience the same force

Explanation

---

---

---

- (c) Which of these satellites is travelling at the greatest speed relative to the Earth? Circle the correct answer from the choices below. Explain your answer in the space provided.  
(3 marks)

LEO

MEO

GEO

All satellites have the same speed

Explanation

---

---

---

- (d) Kepler's Third Law is given on your data sheet. By using relevant equations, in the space below, derive Kepler's Third Law.  
(3 marks)

- (e) Using the information in the picture, calculate the minimum period of a LEO satellite. Provide your answer in minutes and seconds.  
(4 marks)

Answer: \_\_\_\_\_ minutes and \_\_\_\_\_ seconds

**Question B-7=1****(12 marks)**

The Large Hadron Collider (LHC) uses magnetic fields to confine fast-moving charged particles travelling repeatedly around a circular path. The LHC is installed in an underground circular tunnel of circumference 27 km.

- (a) In the presence of a suitably directed uniform magnetic field, charged particles move at constant speed in a circular path of constant radius. By reference to the force acting on the particles, explain how this is achieved and why it happens. (4 marks)

---

---

---

---

---

---

DO NOT WRITE IN THIS AREA AS IT WILL BE CUT OFF

- (b) (i) In one particular experiment, the charged particles travelling around the LHC were protons. Calculate the centripetal force acting on a proton when travelling in a circular path of circumference 27.0 km at one-tenth of the speed of light. Ignore relativistic effects. (3 marks)

Force: \_\_\_\_\_ N

- (ii) Calculate the flux density of the uniform magnetic field that would be required to produce this force. State an appropriate unit. (3 marks)

Answer: \_\_\_\_\_ Units: \_\_\_\_\_

- (c) The Large Hadron Collider is an example of a high energy particle accelerator. Explain the importance of high energy particle accelerators to Physicists. (2 marks)

---

---

---

---

**Question B-7-2** (14 marks)

The Large Hadron Collider is the largest synchrotron in the world, with a total circumference of 26.7 km. While capable of accelerating protons up to 6.50 TeV, first operations in 2013 were run at the

relatively lower 3.50 TeV. The Large Hadron Collider has multiple stages of particle accelerators, starting with a simple linear accelerator and eventually confining the proton beam in the main ring. Very powerful, expensive magnets, powered and cooled to near absolute zero are required to confine the beam.

- (a) Describe how a magnetic field can help keep protons confined within the ring of a synchrotron. (3 marks)

- (b) Explain why the protons in the Large Hadron Collider must first be accelerated in a straight line, with a linear accelerator, rather than starting in a ring like in the synchrotron. (2 marks)

(c) What percentage of the energy of the proton beam used in 2013 is due to the rest mass of the proton? (3 marks)

(d) Explain a benefit of making the confinement ring so large. (3 marks)

(e) Describe how the Large Hadron Collider is used to make new scientific discoveries. (3 marks)

---

**End of Section Two**

**SECTION THREE: Comprehension****20% (36 marks)**

This section has **two (2) questions**. You must answer **both** questions. Write your answers in the spaces provided.

When calculating numerical answers, show your working or reasoning clearly. Give final answers to **three** significant figures and include appropriate units where applicable.

When estimating numerical answers, show your working or reasoning clearly. Give final answers to a maximum of **two** significant figures and include appropriate units where applicable.

Additional working space pages at the end of this Question/Answer booklet are for planning or continuing an answer. If you use these pages, indicate at the original answer, the page number it is planned/continued on and write the question number being planned/continued on the additional working space page.

Suggested working time: 40 minutes.

---

**Question 18****(18 marks)*****The Universe Has A Speed Limit, And It Isn't The Speed Of Light***

When it comes to speed limits, the ultimate one set by the laws of physics themselves is the speed of light. Moreover, anything that's made of matter can only approach, but never reach, the speed of light. If you don't have mass, you must move at the speed of light; if you do have mass, you can never reach it. But practically, in our Universe, there's an even more restrictive speed limit for matter, and it's lower than the speed of light. Here's the scientific story of the real cosmic speed limit.

When scientists talk about the speed of light -  $299,792,458 \text{ ms}^{-1}$  - we implicitly mean "the speed of light in a vacuum." Only in the absence of particles, fields, or a medium to travel through can we achieve this ultimate cosmic speed. Even at that, it's only the truly massless particles and waves that can achieve this speed. This includes photons, gluons, and gravitational waves, but not anything else we know of. But there's no such thing, practically, as a perfect vacuum. Even in the deepest abyss of intergalactic space, there are three things you absolutely cannot get rid of.


1. The WHIM: the warm-hot intergalactic medium. This tenuous, sparse plasma are the leftovers from the cosmic web. While some matter clumps into stars and galaxies, other matter remains in the great voids of the Universe. Starlight ionizes these atoms, creating a plasma that may make up about 50% of the total normal matter in the Universe.
2. The CMB: the cosmic microwave background. This leftover bath of photons originates from the Big Bang, where it was at extremely high energies. Even today, at temperatures just 2.7 degrees above absolute zero, there are over 400 CMB photons per cubic centimetre of space at an average energy of 0.00023 eV per photon.
3. The CNB: the cosmic neutrino background. The Big Bang, in addition to photons, creates a bath of neutrinos. Outnumbering protons by perhaps a billion to one, many of these now-slow-moving particles fall into galaxies and clusters, but many remain in intergalactic space as well.

Any particle traveling through the Universe will pass through particles from the WHIM, neutrinos from the CNB, and photons from the CMB.

DO NOT WRITE IN THIS AREA AS IT WILL BE CUT OFF

The Large Hadron Collider accelerates particles here on Earth up to a maximum velocity of  $299,792,455 \text{ ms}^{-1}$ , or 99.999999% the speed of light. The highest-energy cosmic rays have approximately 36 million times the energy of the fastest protons ever created at the Large Hadron Collider. Assuming that these cosmic rays are also made of protons gives a speed of  $299,792,457.999999999992 \text{ ms}^{-1}$ , which is extremely close to, but still below, the speed of light in a vacuum. There's a very good reason that, by the time we receive them, these cosmic rays aren't more energetic than this.

If there is a particle with energies in excess of  $5.00 \times 10^{19} \text{ eV}$ , they can only travel a few million light years - max - before a photon from the CMB, interacts with it. When that interaction occurs, there will be enough energy to produce a neutral pion, which steals energy away from the original particle, following from  $E = mc^2$ .



The more energetic your particle is, the more likely you are to produce pions, which you'll continue to do until you fall below this theoretical cosmic energy limit of  $5.00 \times 10^{19} \text{ eV}$ , known as the GZK cutoff. *cut-off*.

We believe that every charged particle in the cosmos - every cosmic ray, every proton, every atomic nucleus - should be limited by this speed. Not just the speed of light, but a little bit lower, thanks to the leftover glow from the Big Bang and the particles in the intergalactic medium. If we see anything that's at a higher energy, then it either means:

1. particles at high energies might be playing by different rules than the ones we presently think they do,
2. they are being produced much closer than we think they are: within our own Local Group or Milky Way, rather than these distant, extragalactic black holes,
3. or they're not protons at all, but composite nuclei.

The few particles we've seen that break the GZK barrier are indeed in excess of  $5 \times 10^{19} \text{ eV}$ , in terms of energy, but do not exceed  $3 \times 10^{21} \text{ eV}$ , which would be the corresponding energy value for an iron nucleus. Since many of the highest-energy cosmic rays have been confirmed to be heavy nuclei, rather than individual protons, this reigns as the most likely explanation for the extreme ultra-high-energy cosmic rays.

There is a speed limit to the particles that travel through the Universe, and it isn't the speed of light. Instead, it's a value that's very slightly lower, dictated by the amount of energy in the leftover glow from the Big Bang. As the Universe continues to expand and cool, that speed limit will slowly rise over cosmic timescales, getting ever-closer to the speed of light. But remember, as you travel through the Universe, if you go too fast, even the radiation left over from the Big Bang can fry you. So long as you're made of matter, there's a cosmic speed limit that you simply cannot overcome.

- (a) Name three things in the Universe that can move at the cosmic speed limit and describe the requirements to reach the cosmic speed limit. (3 marks)

- *Photons, gluons and gravity waves.* (1)
- *Must be massless.* (1)
- *Move at the cosmic speed limit in the absence of any field, particles or medium.* (1)

- (b) A neutral pion has a rest mass of  $135 \text{ MeV}/c^2$ . Calculate the minimum number of neutral pions that need to be produced by a  $5.03 \times 10^{19} \text{ eV}$  particle to reduce to the cosmic energy limit. (3 marks)

$$\begin{aligned} E_{\text{lost}} &= 5.03 \times 10^{19} - 5.00 \times 10^{19} \\ &= 3.00 \times 10^7 \text{ eV} \quad (1) \end{aligned}$$

$$E_{\text{rest}} = 135 \times 10^6 \text{ eV}. \quad (1)$$

$$\begin{aligned} \therefore \# \text{ particles} &= \frac{3.00 \times 10^7}{135 \times 10^6} \\ &= \underline{\underline{2.22 \times 10^9}} \quad (1) \end{aligned}$$

- (c) The article compares the Large Hadron Collider and cosmic rays. Protons in the Large Hadron Collider are accelerated up to 6.50 TeV. What is the energy of the most energetic cosmic rays in eV? (3 marks)

$$\begin{aligned} E(\text{cosmic rays}) &= (36 \times 10^6)(6.50) \quad (1) \\ &= 2.34 \times 10^8 \text{ TeV} \quad (1) \\ &= \underline{\underline{2.34 \times 10^{20} \text{ eV}}} \quad (1) \end{aligned}$$

Answer:  $2.34 \times 10^{20}$  eV

- (d) Calculate the average frequency of a photon from the CMB.

(3 marks)

$$\begin{aligned} E(\text{average}) &= 0.00023 \text{ eV} \\ &= 3.68 \times 10^{-23} \text{ J} \quad (1) \end{aligned}$$

$$\begin{aligned} E &= hf \\ \Rightarrow f &= \frac{E}{h} \\ &= \frac{3.68 \times 10^{-23}}{6.63 \times 10^{-34}} \quad (1) \\ &= 5.55 \times 10^{10} \text{ Hz} \quad (1) \end{aligned}$$

Answer:  $5.55 \times 10^{10}$  Hz

- (e) Explain why the interaction of a very energetic particle with a photon can cause the particle to slow down.

(3 marks)

- When a photon meets a high-energy particle, it can create a new particle. (1)
- Creating a new particle requires energy (according to  $E=mc^2$ ). (1)
- This energy is taken from the  $E_k$  of the original particle, slowing it down. (1)

- (f) Describe which of the three reasons given in the article is the most likely cause when we have detected particles more energetic than the GZK barrier.

(3 marks)

- Reason 3 - They are composite nuclei. (1)
- The energy has not exceeded the energy of the iron nucleus. (1)
- High-energy cosmic rays have been confirmed as heavy nuclei. (1)

**Question 19****(18 marks)*****Escape velocity and The Black Hole***

Article adapted from Fundamentals of Modern Physics by Peter J Nolan. 2014



The simplest way to describe the black hole is to start with a classical analogue. Suppose we wished to launch a rocket from the earth to a far distant place in outer space. How fast must the rocket travel to escape the gravitational pull of the earth? This value is known as the *Escape Velocity*. When we launch the rocket it has a velocity  $v$ , and hence, a kinetic energy. As the rocket proceeds into space, its velocity decreases but its potential energy increases. The absolute potential energy of an object when it is a distance  $r$  away from the center of the earth is found from:

$$PE = -\frac{GM_e m}{r}$$

where  $G$  is the universal gravitational constant,  $M_e$  is the mass of the Earth, and  $m$  is the mass of the object. Let us now apply this potential energy term to a rocket that is trying to escape from the gravitational pull of the earth. The total energy of the rocket at any time is equal to the sum of its potential energy and its kinetic energy, that is:

$$E = KE + PE = \frac{1}{2}mv^2 - \left[ GM_e m \left( \frac{1}{r} \right) \right]$$

When the rocket is fired from the surface of the earth,  $r = R$ , at an escape velocity  $v_e$ , its total energy will be:

$$E = \frac{1}{2}mv_e^2 - \left[ GM_e m \left( \frac{1}{R} \right) \right]$$

By the law of conservation of energy, the total energy of the rocket remains a constant. Hence, we can equate the total energy at the surface of the earth to the total energy when the rocket is far removed from the earth. That is:

$$\frac{1}{2}mv_e^2 - \left[ GM_e m \left( \frac{1}{R} \right) \right] = \frac{1}{2}mv^2 - \left[ GM_e m \left( \frac{1}{\infty} \right) \right]$$

When the rocket escapes the pull of the earth, it has effectively traveled to infinity, that is,  $r = \infty$ , and its velocity at that time is reduced to zero, that is,  $v = 0$ . Hence, the equation reduces to:

$$\begin{aligned} \frac{1}{2}mv_e^2 - \left[ GM_e m \left( \frac{1}{R} \right) \right] &= 0 - \left[ GM_e m \left( \frac{1}{\infty} \right) \right] = 0 \\ \frac{1}{2}mv_e^2 &= \frac{GM_e m}{R} \\ v_e^2 &= \frac{2GM_e}{R} \\ v_e &= \sqrt{\frac{2GM_e}{R}} \end{aligned}$$

If we substitute  $v_e$  for the  $c$ , the speed of light and rearrange, we get a formula that tells us the maximum radius of any object with mass, for light to be able to escape from it.

$$R_s = \frac{2GM_e}{c^2}$$

This value is called the *Schwarzchild Radius* and any distance to an object closer than this value is said to be within the *Event Horizon*, a region from which nothing, not even light can escape!

The reason for the name, black hole, comes from the idea that if we look at an object in space, such as a star, we see light coming from that star. If the star became a black hole, no light could come from that star. Hence, when we look into space we would no longer see a bright star at that location, but rather nothing but the blackness of space. There seems to be a hole in space where the star used to be and therefore we say that there is a black hole there.

Solving the Schwarzschild radius of the sun, by replacing the mass of the earth by the mass of the sun, we get  $2.95 \times 10^3$  m. Thus, if the sun were to contract to a radius below  $2.95 \times 10^3$  m the gravitational force would become so great that no light could escape from the sun, and the sun would become a black hole.

- (a) Explain the relationship between the concept of escape velocity and the concept of a black hole. (2 marks)

Escape velocity - minimum velocity an object needs to escape the gravitational attraction of an object. This is determined by the object's mass. (1)

Black hole - so massive they have an extreme gravitational attraction. Escape velocity is greater than the speed of light so light can't escape. (1)

- (b) The article states that:

"When the rocket escapes the pull of the earth it has effectively travelled to infinity, that is,  $r = \infty$ , and its velocity at that time is reduced to zero, that is,  $v = 0$ ."

- (i) Use Newton's Law of Universal Gravitation to support the argument that at  $r = \infty$ , the rocket has escaped the pull of the Earth. (2 marks)

- $F = \frac{GM_E m}{r^2}$  (1)

- As  $r \rightarrow \infty$ ,  $F \rightarrow 0$ . so the object can leave the Earth as there is no gravitational attraction. (1)

- (ii) If an object left the Earth with the minimum *escape velocity*, why would this value be zero, when  $r$  is equal to infinity? (2 marks)

- When leaving at escape velocity,  $E_K = \text{the absolute E.p. (1)}$

- As  $E_p$  becomes less negative as  $r \rightarrow 0$ ,  $E_k$  reduces as well } (1)
- As  $E_p \rightarrow 0$ ,  $E_k \rightarrow 0 \Rightarrow v \rightarrow 0$ .

- (c) (i) Show that the escape velocity for an object to leave the Earth's gravitational pull is equal to  $1.12 \times 10^4 \text{ ms}^{-1}$ . \_\_\_\_\_ (2 marks)

$$\begin{aligned}
 \checkmark v_{\text{escape}} &= \sqrt{\frac{2GM_E}{r}} \\
 &= \sqrt{\frac{2(6.67 \times 10^{-11})(5.97 \times 10^{24})}{(6.37 \times 10^6)}} \quad (1) \\
 &= 1.12 \times 10^4 \text{ ms}^{-1} \quad (1)
 \end{aligned}$$

- (ii) Show that the total energy of 12.0 tonne space craft launched at its escape velocity from the surface of the Earth is zero. (3 marks)

$$\begin{aligned}
 E &= \frac{1}{2}mv_{\text{escape}}^2 - \frac{GM_em}{R} \quad (1) \\
 &= \frac{1}{2} (12.0 \times 10^3) (1.12 \times 10^4)^2 - \frac{(6.67 \times 10^{-11})(5.97 \times 10^{24})(12.0 \times 10^3)}{6.37 \times 10^6} \quad (1) \\
 &= 7.53 \times 10^{11} - 7.50 \times 10^{11} \quad (1) \\
 &\approx 0
 \end{aligned}$$

- (iii) At what speed would this space craft be travelling in deep space at a distance of 325 million kilometres from the Earth? (3 marks)

$$\begin{aligned} E_p &= -\frac{GM_{EM}}{r} \\ &= -\frac{(6.67 \times 10^{-11})(5.97 \times 10^{24})}{(6.37 \times 10^6 + 325 \times 10^9)} (1) \\ &= -1.47 \times 10^7 \text{ J} \end{aligned}$$

$$\therefore E_k = 1.47 \times 10^7 \text{ J} \quad ((1))$$

See next page

$$E_k = \frac{1}{2}mv^2$$

$$\Rightarrow v = \sqrt{\frac{2E_k}{m}}$$

$$= \sqrt{\frac{2(1.47 \times 10^7)}{(12.0 \times 10^{-3})}}$$

Speed \_\_\_\_\_ ms<sup>-1</sup>

$$= 49.5 \text{ ms}^{-1} \quad (1)$$

- (d) One of the largest stars in our galaxy is Betelgeuse. This star has a radius 887 times that of our Sun and a mass 11.6 times that of our Sun. Explain what is meant by the Schwarzschild radius for this star and calculate its value. (4 marks)

• Schwarzschild radius - minimum distance from a black hole to be beyond the event horizon. (1)

$$R_s = \frac{2GM}{c^2} \quad (1)$$

$$= \frac{2(6.67 \times 10^{-11})(11.6)(1.99 \times 10^{30})}{(3.00 \times 10^8)^2} \quad (1)$$

$$= \underline{3.42 \times 10^4 \text{ m}} \quad (1)$$

Schwarzschild radius:  $3.42 \times 10^4 \text{ m}$

End of Questions

## Additional Working Space

Question number: \_\_\_\_\_

**DO NOT WRITE IN THIS AREA AS IT WILL BE CUT OFF**

## Additional Working Space

Question number: \_\_\_\_\_

DO NOT WRITE IN THIS AREA AS IT WILL BE CUT OFF

**Acknowledgements:**

Picture of Earth Satellites:

<https://assets.pcmag.com/media/images/600043-leo-meo-geo-satellite-orbital-heights.jpg?thumb=y>

Antihydrogen Antics

Article adapted from: <https://www.physicscentral.com/explore/action/antics.cfm>

Escape velocity and The Black Hole

Article adapted from Fundamentals of Modern Physics by Peter J Nolan. 2014

DO NOT WRITE IN THIS AREA AS IT WILL BE CUT OFF