

Drug	Form	Use	Effects	Risks
Ketamine	Liquid, pills, white powder	Injected, snorted, swallowed	This depressant drug is an anaesthetic that causes users to experience feelings of disconnectedness and relaxation, as well as hallucinations and paralysis.	There is a risk of abdominal pain, bladder problems, increased heart rate, liver damage, and infection as a result of injecting. It is linked to depression with frequent use.
LSD (acid)	Small absorbant squares of paper known as "tabs", liquid	Swallowed	This hallucinogenic drug is very powerful. It causes a "trip," a period of time in which a user's perception of sounds, color, objects, time, and movement is altered.	Bad trips can occur, with frightening or unpredictable hallucinations. LSD should not be taken if the user is in a bad mood or feeling anxious or depressed. It can exacerbate existing mental health problems. There is a risk of flashbacks weeks, months, and years after taking LSD.
Magic mushrooms	Dried mushrooms of different colors	Brewed as a tea, eaten	This hallucinogen causes altered perception of colors, sounds, objects, and time, and feelings of relaxation or fright.	Users may eat poisonous mushrooms by mistake. Some people experience sickness and confusion, flashbacks, or feelings of terror afterward.
MDMA/Ecstasy	White powder (MDMA), pills (ecstasy); MDMA is the active ingredient in ecstasy pills	Powder is rubbed on gums; ecstasy pills are swallowed	MDMA is a stimulant that makes users feel chatty, energized, elated, and loving. Users experience colors, feelings, and sounds more intensely.	Users can dehydrate and overheat, or drink too much water, which can also be dangerous. It can cause confusion, anxiety, and panic. Ecstasy pills often contain other unknown drugs, which can lead to unexpected reactions. Taking MDMA or ecstasy has been linked to kidney, liver, and heart problems, and death.
Poppers	Liquid	Inhaled, sniffed	Poppers can be both a stimulant and a depressant. They cause "head rushes" and feelings of faintness, sickness, or lowered sexual inhibition.	Users might feel disorientated or sick, or become unconsciousness, or have reduced blood pressure. Users may engage in risky sexual behavior. The liquid can burn the skin.
Solvents	Aerosols, gases, glues	Inhaled, sniffed	Solvents can work as depressants and hallucinogens, leading users to experience feelings of drunkenness, slow reactions, and hallucinations.	Users might experience confusion, disorientation, unconsciousness, mood swings, poor judgment, or vomiting, and higher risk of death.
Tranquilizers	Injections, pills	Injected, swallowed	Tranquilizers can be depressants and hallucinogens. They slow the user's reactions and cause hallucinations and feelings of calmness and relaxation. They are highly addictive.	Users are at risk of anxiety, depression, headaches, memory loss, seizures, and vomiting.