





Two toes point forward and two point backward to give a strong grip.

Toucans live in groups and often play games. They wrestle with their bills or toss fruit to each other.



Powerful, hooked bill for tearing flesh from food.

Massive sharp talons enable the eagle to grab and crush prey.

The harpy eagle pursues capuchin monkeys and other small animals, such as opossums and coatis, through the rainforest.

Harpy eagle
(*Harpia harpyja*)
Length: 3 ft 4 in (1 m)

Harpy eagle

The huge and fearsome harpy eagle swoops through the treetops seeking its prey. It can fly at speeds of up to 50 mph (80 kph). Its gray, mottled coloring helps camouflage it among the leaves so the animals it hunts do not see it coming. Harpy eagles build nests made of sticks high up in the trees.

White-plumed antbird

Despite its name, the white-plumed antbird does not eat ants. It follows army ants and feeds on the spiders and insects that are trying to escape them. Usually the antbird darts down to the ground or hangs from a branch to snatch up a meal. But sometimes it will hop among the ants, holding its tail up out of the way. Its long legs help protect its body from the ants' stings.




White-plumed antbird
(*Pithys albifrons*)
Length: 5 in (12.5 cm)

The antbird often perches on a small tree and waits for ants to appear.

Columns of army ants swarming over the forest floor.

Rufous-breasted hermit

The hermit is a kind of hummingbird. The hermit's thin, curved bill probes flowers for nectar, which flows into the grooves of its long tongue as it flicks in and out. Nectar is easy to digest so it provides instant energy to power the hummingbird's whirring wings, and also helps the bird generate body warmth.



Rufous-breasted hermit
(*Glaucis hirsutus*)
Length: 5 in (12 cm)

Hoatzin

The hoatzin has weak wings and it flaps clumsily through flooded parts of the rainforest along quiet riverbanks. It uses its wings and tail to help it clamber through the trees as its feet are not very strong. Hoatzins move about and nest in groups, building untidy twig nests by the water. Young hoatzins leave the nest soon after hatching and may leap into the water to escape danger. Hoatzins feed mostly on leaves and have multichambered stomachs, like cows, to digest them.

Hoatzin
(*Opisthocomus hoazin*)
Length: 2 ft 3 in (70 cm)

A baby hoatzin has two tiny claws on the bend of each wing to help it grip and climb.

