


MONSTER CLAWS

People have always feared unknown monsters of the deep. The creature in this old engraving looks like a lobster with two giant-sized pincers. Real lobsters can grow to 35 in (90 cm) long and weigh up to 45 lb (20 kg).



Surface is rough and hard to the touch

Very long antenna, or feeler


Small pincer to tear food

SMALL LOBSTERLIKE CRUSTACEAN

Squat lobsters are small enough to hide under stones, emerging under cover of darkness to find food. They usually crawl along the bottom, but in danger they can swim for short distances by flapping their abdomen.

Body length up to 5 in (13 cm)

Abdomen tucks under body



Mouth in center of star, beneath the body

Diameter can be as much as 12 in (30 cm)



ROCK STAR


Normally purple, this North Atlantic starfish is sometimes yellow or red. It hangs onto rocks with rows of suckered tube-feet underneath its arms. While most starfish have five arms or multiples of five, the purple sunstar has between seven and 13 arms.

FINE FEATHERS

Feather stars are echinoderms (creatures with body parts radiating from the center), along with starfish, sea urchins, and sea lilies. Like most echinoderms, their body pattern is arranged in fives or sets of fives. This feather star has 10 feathery arms for capturing food particles in the water. Rows of pointed tube-feet pass food particles down a slimy channel and into the mouth in the middle of its upper side. The arms are used to crawl along the seabed and for swimming. It also has leglike cirri for hanging onto rocks.

Arms span body and vary in size from 3 to 6 in (7 to 15 cm) long

Tube-feet line the fine branches of the feathered arms



Long dorsal fin

