



Zantedeschia aethiopica
'Crowborough'

Early to mid-summer-flowering tuber with arrow-shaped, semierect, basal, deep green leaves. Produces a succession of arum-like, white spathes, each with a yellow spadix. **H** 1½–3ft (45–100cm), **S** 1–1½ft (35–45cm).

☀️💧 Z8–10 ⓘ



Rhynchospora colorata

Slow-growing, deciduous, spreading, perennial, marginal water plant or bog plant with pointed, green-tipped, white bracts surrounding an inconspicuous flower head in late summer. Has linear, slightly hairy, mid-green leaves. **H** 18in (45cm), **S** indefinite.


☀️💧 Z7–10



Rodgersia aesculifolia

Clump-forming, rhizomatous perennial that is excellent for a bog garden or pond margin. In midsummer, plumes of fragrant, pinkish-white flowers rise above crinkled, bronze foliage like that of a horse-chestnut tree. **H** and **S** 3ft (1m).

☀️💧 Z5–8



Sanguisorba canadensis
(Canadian burnet)

Clump-forming perennial. In late summer, bears slightly pendent spikes of bottlebrush-like, white flowers on stems that arise from toothed, divided, mid-green leaves. **H** 4–6ft (1.2–2m), **S** 2ft (60cm).


☀️💧 Z3–8



Rodgersia sambucifolia

Deciduous, clump-forming, perennial, bog plant with emerald-green leaves, sometimes bronze-tinged, divided into large, lobed leaflets. Bears broad spires of creamy-white flowers in summer. **H** 5ft (1.5m), **S** 3ft (1m).

☀️💧 Z5–8



Lysimachia clethroides

Vigorous, clump-forming, spreading perennial bearing spikes of small, white flowers above mid-green foliage in late summer. **H** 3ft (1m), **S** 2–3ft (60–100cm).

☀️💧 Z4–9



Anemone rivularis

Perennial with stiff, free-branching stems bearing delicate, cup-shaped, white flowers in summer above deeply divided, dark green leaves. **H** 24in (60cm), **S** 12in (30cm).

☀️💧 Z6–8 ⓘ