

ADROMISCHUS

white-margined, pink or purple lobes, on a stem 10in (25cm) or more long. Z14-15 (min. 45°F/7°C)
A. maculatus illus. p.492.

AECHMEA

BROMELIACEAE

Genus of evergreen, rosette-forming, mostly epiphytic perennials, cultivated for their foliage, flowers and fruits. May be grown in full light or a semishaded site. Provide a rooting medium of equal parts humus-rich soil and either sphagnum moss or bark or plastic chips used for orchid culture. Using soft water, water moderately in summer, sparingly at all other times, and keep cuplike, rosette centers filled with water from spring through to fall. Propagate by offsets in late spring.
A. distichantha. Evergreen, basal-rosetted, epiphytic perennial. H and S to 3ft (1m). Forms dense rosettes of narrowly oblong, round-tipped, arching leaves, dull green above, gray and scaly beneath. Has panicles of small, tubular, purple or blue flowers among white-felted, pink bracts, usually in summer. Z15 (min. 50°F/10°C)
A. fasciata, syn. *Billbergia rhodocyanea* (Silver vase plant, Urn plant; illus. p.471). Evergreen, tubular-rosetted, epiphytic perennial. H 16-24in (40-60cm), S 12-20in (30-50cm). Has loose rosettes of broadly oblong, round-tipped, incurved, arching leaves with dense, gray scales and silver cross-banding. Bears dense, pyramidal panicles of tubular, blue-purple flowers among pink bracts, above foliage, from spring to fall. Z11-12 (min. 50°F/10°C)
A. Foster's Favorite Group (Lacquered wine-cup). Evergreen, basal-rosetted, epiphytic perennial. H and S 1-2ft (30-60cm). Has loose rosettes of strap-shaped, arching, lustrous, wine-red leaves. Drooping spikes of small, tubular, deep purple-blue flowers are borne in summer, followed later by pear-shaped, red fruits. Z15 (min. 50°F/10°C)
A. fulgens (Coral berry). Evergreen, basal-rosetted, epiphytic perennial. H to 20in (50cm), S 16in (40cm). Forms loose rosettes of broadly oblong, arching, glossy, mid-green leaves with gray scales beneath and rounded or pointed tips. In summer, produces, above foliage, erect panicles of small, tubular, violet-purple flowers that turn red with age. These are succeeded by small, rounded to ovoid, red fruits on red stalks. Z15 (min. 50°F/10°C)
A. nudicaulis. Evergreen, basal-rosetted, epiphytic perennial. H to 28in (70cm), S 10in (25cm). Produces loose rosettes of a few broadly strap-shaped, arching, olive-green leaves with spiny edges and usually banded with gray scales beneath. Spikes of small, tubular, yellow flowers open above large, red bracts in summer. Z9-11 (min. 50°F/10°C)
A. recurvata (illus. p.471). Evergreen, basal-rosetted, epiphytic perennial. H to 8in (20cm), S to 20in (50cm). Narrowly triangular, tapered, spiny-edged, arching, red-flushed, mid-green leaves are produced in dense rosettes. In summer, bears a short, dense spike of tubular, red-and-white flowers, with red bracts, just above leaves. Z9-11 (min. 50°F/10°C)

AEGOPODIUM

Bishop's weed, Goutweed, Ground elder

APIACEAE/UMBELLIFERAE

Genus of invasive, rhizomatous perennials, most of which are weeds, although *A. podagraria* 'Variegatum' is excellent as groundcover. Tolerates sun or shade and any well-drained soil. Propagate by division of rhizomes in spring or fall.
A. podagraria 'Variegatum'. Vigorous, spreading perennial. H 1-2ft (30-60cm), S indefinite. Has lobed, creamy-white-variegated leaves. Insignificant, white flowers, borne in summer, are best removed. Z4-9

AEONIUM

CRASSULACEAE

Genus of evergreen, perennial, rarely biennial succulents, often subshrubby, grown for their rosettes of bright green or blue-green, occasionally purple, leaves. Prefers partial shade and very well-drained soil. Most species grow from fall to spring and are semidormant in midsummer. Propagate by seed in summer or, for branching species, by stem cuttings in spring or summer.
A. arboreum. Erect, succulent subshrub. H and S to 6ft (2m). Branched stems are each crowned by a rosette, up to 6in (15cm) across, of broadly lance-shaped, glossy, bright green leaves. In spring, produces cones of small, star-shaped, golden flowers on 2-3-year-old stems, which then die back. Z9-11. (min. 41°F/5°C).
'Schwarzkopf' see *A. 'Zwartkop'*.
A. haworthii (Pinwheel) illus. p.489.
A. tabuliforme illus. p.491.
A. 'Zwartkop', syn. *A. arboreum* 'Schwarzkopf', illus. p.310.

AESCHYNANTHUS

GESNERIACEAE

Genus of evergreen subshrubs, climbers and trailing and semitrailing perennials, some of which are epiphytic, useful for growing in hanging baskets. Needs a fairly humid atmosphere and a position out of direct sun. Water sparingly in low temperatures. Propagate by tip cuttings in spring or summer.
A. 'Black Pagoda'. Semitrailing perennial. H 2ft (60cm), S to 18in (45cm). Elliptic leaves are pale green with dark brown marbling above, purple beneath. Produces terminal clusters of burnt-orange flowers, with green calyces, from summer to winter. Z15 (min. 59-64°F/15-18°C)
A. longicaulis, syn. *A. marmoratus*, *A. zebrinus*. Evergreen, semitrailing perennial. H 2ft (60cm), S to 36in (90cm). Ovate, waxy leaves are dark green, veined yellowish-green above, purplish below. Bears tubular, greenish flowers, with dark brown markings, in terminal clusters in summer. Z15 (min. 59-64°F/15-18°C)
A. marmoratus. See *A. longicaulis*.
A. pulcher (Lipstick plant, Royal red bugler). Evergreen, epiphytic climber. H 30in (75cm). Produces thick, ovate leaves and small, tubular, hooded,

bright red flowers, with yellow throats, borne in terminal clusters from summer to winter. Z14-15 (min. 59-64°F/15-18°C)
A. speciosus, syn. *A. splendens*, illus. p.478.
A. splendens. See *A. speciosus*.
A. zebrinus. See *A. longicaulis*.

AESCULUS

Buckeye, Horse chestnut

SAPINDACEAE/HIPPOCASTANACEAE

Genus of deciduous trees and shrubs, grown for their bold, divided leaves and conspicuous, upright panicles or clusters of flowers, followed by fruits (horse chestnuts) sometimes with spiny outer casings. Requires sun or semishade and fertile, well-drained soil. Propagate species by sowing seed in fall, cultivars by budding in late summer or by grafting in late winter. Leaf spot may affect young foliage, and coral spot fungus may attack damaged wood. ☉ All parts of these plants may cause mild stomach upset if ingested.
A. californica (California buckeye) illus. p.81.
A. x carnea (Red horse chestnut). **'Briotii'** illus. p.60.
A. chinensis (Chinese horse chestnut) illus. p.60.
A. flava, syn. *A. octandra* (Sweet buckeye, Yellow buckeye), illus. p.76.
A. glabra (Ohio buckeye). Deciduous, broadly conical, sometimes shrubby tree. H 50ft (15m), S 30ft (10m). Leaves, usually composed of 5 narrowly ovate leaflets, are dark green. Bears 4-petaled, greenish-yellow flowers in upright clusters in late spring and early summer. Z3-7. **'October Red'** has bronze-red fall leaves.
A. hippocastanum. Vigorous, deciduous, spreading, rounded tree. H 80ft (25m), S 70ft (20m). Has large leaves with 5 or 7 leaflets and spires of white flowers, flushed pink and yellow in centers in spring. Spiny fruits contain glossy, brown nuts in fall. Z3-8. **'Baumannii'** has dark green leaves turning to yellow in fall and double, yellow- or red-marked, white flowers.
'Hampton Court Gold', H 50ft (15m), S 25ft (8m), is slower-growing, and bears soft-yellow leaves in spring, greener in summer; foliage may scorch in hot sun.
A. indica (Indian horse chestnut). Deciduous, spreading, rounded tree. H and S 50ft (15m). Glossy, dark green leaves with usually 7 narrowly ovate leaflets are bronze when young, orange or yellow in fall. Upright panicles of 4-petaled, pink-tinged, white flowers, marked with red or yellow, are borne in midsummer. Z7-8.
'Sydney Pearce' illus. p.73.
A. x neglecta (Sunrise horse chestnut). **'Erythroblastos'** illus. p.72.
A. octandra. See *A. flava*.
A. parviflora (Bottlebrush buckeye) illus. p.113.
A. pavia (Red buckeye). Deciduous, conical shrub or small tree. H 15ft (5m), S 10ft (3m). Glossy, dark green leaves consist of 5 narrowly ovate leaflets. Has panicles of 4-petaled, red flowers in early summer. Z5-9. **'Atrosanguinea'** illus. p.86. **var. discolor 'Koehnei'**, H and S 6ft (2m), is very slow-growing and compact, with deep bronze young leaves and soft orange-red and yellow flowers.

A. turbinata (Japanese horse chestnut). Deciduous, spreading, sturdy-branched tree. H 70ft (20m), S 40ft (12m). Large, dark green leaves have 5 or 7, narrowly ovate leaflets. Bears panicles of creamy-white flowers in late spring and early summer. Z6-8

AETHIONEMA

BRASSICACEAE/CRUCIFERAE

Genus of evergreen or semievergreen, dwarf subshrubs, woody-based perennials and annuals, grown for their prolific flowers. Needs sun and well-drained soil. Propagate by softwood cuttings in spring or by seed in fall. Most species self-seed readily.
A. armenum illus. p.364.
A. grandiflorum, syn. *A. pulchellum* (Persian stone cress), illus. p.338.
A. iberideum. Evergreen or semievergreen, rounded, compact shrub. H and S 6in (15cm). Bears small, lance-shaped, gray-green leaves and, in summer, ¾in (2cm) stems each bear a raceme of small, saucer-shaped, white flowers. Z6-8
A. pulchellum. See *A. grandiflorum*.
A. 'Warley Rose' illus. p.362.
A. 'Warley Ruber'. Evergreen or semievergreen, rounded, compact subshrub. H and S 6in (15cm). Has tiny, linear, bluish-green leaves. Racemes of small, deep rose-pink flowers are produced on ¾-1¼in (2-3cm) stems in spring-summer. Z6-8

AGAPANTHUS

AGAPANTHACEAE

Genus of clump-forming perennials, some evergreen, with erect stems that carry large umbels of bell- to tubular-bell-shaped or trumpet-shaped flowers, usually blue and often fading to purple with age. Leaves are strap-shaped. Grow in full sun and in moist, well-drained soil. Protect crowns with mulch in winter. Plants increase slowly but may be propagated by division in spring; may also be raised from seed in fall or spring. Named cultivars will not come true from seed.
A. africanus (African lily). Evergreen, clump-forming perennial. H 3ft (1m), S 20in (50cm). In late summer, has rounded umbels of deep blue flowers on upright stems, above broad, dark green leaves. Z9-10
A. 'Alice Gloucester'. Clump-forming perennial. H 3ft (1m), S 20in (50cm). Produces large, dense, rounded umbels of white flowers in summer, above narrow, mid-green leaves. Z7-11
A. 'Ben Hope'. Clump-forming perennial. H to 4ft (1.2m), S 2ft (60cm). Erect stems support dense, rounded umbels of deep blue flowers in late summer and early fall, borne over narrow, grayish-green leaves. Z7-10
A. 'Blue Giant'. Clump-forming perennial. H 4ft (1.2m), S 2ft (60cm). Has rounded heads of open, bell-shaped rich blue flowers in mid- to late summer. Z8-11
A. campanulatus. Clump-forming perennial. H 2-4ft (60cm-1.2m), S 18in (45cm). Rounded umbels of blue flowers are borne on strong stems in summer, above narrow, grayish-green leaves. Z7-11