

semiripe cuttings taken in late summer or by seed sown in spring.

A. polifolia. Evergreen, open, twiggy shrub. H to 16in (40cm), S to 2ft (60cm). Has narrow, leathery, glossy, mid-green leaves. Bears terminal clusters of pitcher-shaped, pink flowers in spring and early summer. Z2–6. ‘**Alba**’ illus. p.332.

‘**Compacta**’ illus. p.333. ‘**Macrophylla**’, H 2–6in (5–15cm), S 10in (25cm), is low-growing, with broadly ovate, dark green leaves and numerous deep pink and white flowers, slightly larger and more rounded than the species.

ANDROSACE

PRIMULACEAE

Genus of annuals, biennials and predominantly evergreen perennials, usually compact cushion-forming and often with soft, hairy leaves. Many species are suitable for cold greenhouses and troughs with winter cover. Needs sun and very well-drained soil; some species prefer acidic soil. Resents wet foliage in winter. Propagate by tip cuttings in summer or by seed in fall. Prone to botrytis and attack by aphids.

A. carnea illus. p.350. **subsp. laggeri** illus. p.353.

A. chamaejasme. Evergreen, basal-rosetted, variable perennial with easily rooted stolons. H 1¼–2½in (3–6cm), S 6–8in (15–20cm). Produces ovate, hairy leaves and, in spring, clusters of 2–8 flattish, white flowers, each with a yellow eye that sometimes turns red with age. Z4–7.

A. cylindrica. Evergreen, basal-rosetted perennial. H ½–¾in (1–2cm), S 4–6in (10–15cm). Leaves are linear and glossy. Flower stems each carry up to 10 small, flattish, white flowers, each with a yellow-green eye, in early spring. Suitable for a cold greenhouse. Z6–7.

A. hedraeantha. Evergreen, cushion-forming perennial. H ½–¾in (1–2cm), S to 4in (10cm). Has loose rosettes of narrowly ovate, glossy leaves, and bears umbels of 5–10 flattish, yellow-throated, pink flowers in spring. Best in a cold greenhouse. Z6–7.

A. hirtella. Evergreen, tight cushion-forming perennial. H ½in (1cm), S to 4in (10cm). Has rosettes of small, thick, linear to oblong, hairy leaves. Almond-scented, flattish, white flowers are borne in spring on very short stems, 1 or 2 per rosette. Z6–8.

A. imbricata. See *A. vandellii*.

A. lanuginosa illus. p.363.

A. pyrenaica illus. p.350.

A. sarmentosa. Evergreen, mat-forming perennial, spreading by runners. H 2–4in (5–10cm), S to 12in (30cm). Has open rosettes of small, narrowly elliptic, hairy leaves. Large clusters of flattish, yellow-eyed, bright pink flowers are produced in spring. A good rock plant in all but very wet areas. Z5–7.

A. sempervivoides. Evergreen, mat-forming, rosetted perennial with stolons. H 1–2in (2.5–5cm), S 6–8in (15–20cm). Has leathery, oblong or spoon-shaped leaves. In spring, produces small heads of 4–10 flattish, pink flowers, with yellow, then red, eyes. A good rock plant. Z5–7.

A. vandellii, syn. *A. imbricata*, illus. p.347.

A. villosa illus. p.349. **var. jacquemontii** illus. p.366.

ANEMANTHELE

POACEAE/GRAMINEAE

See also GRASSES, BAMBOOS, RUSHES and SEDGES.

A. lessoniana, syn. *Stipa arundinacea* (Pheasant’s-tail grass), illus. p.289.

ANEMONE

Windflower

RANUNCULACEAE

Genus of spring-, summer- and fall-flowering perennials, sometimes tuberous or rhizomatous, with mainly rounded, shallowly cup-shaped flowers. Leaves are rounded to ovate, often divided into 3–15 leaflets. Most thrive in full light or semishade in humus-rich, well-drained soil. Propagate by division in spring, by seed sown in late summer, when fresh, or by root cuttings in winter. ☉Contact with the sap may irritate skin.

A. apennina (Apennine anemone). Spreading, spring-flowering, rhizomatous perennial. H 8in (20cm), S 12in (30cm). Fernlike leaves have 3 deeply toothed lobes. Each stem bears a large, upright, flattish, blue, white or pink flower, with 10–20 narrow petals. Z5–9.

A. blanda. Spreading, early spring-flowering perennial with a knobbly tuber. H and S 6in (15cm). Leaves are broadly ovate and semierect, with 3 deeply toothed lobes. Each stem bears an upright, flattish, blue, white or pink flower, 1½–2in (4–5cm) across, with 9–14 narrow petals. Z4–8.

‘**Atrocaerulea**’ illus. p.419. ‘**Ingramii**’ bears purple-backed, deep blue flowers.

var. rosea ‘Radar’, syn. *A.b. ‘Radar’*, illus. p.418. ‘**Violet Star**’ illus. p.418.

‘**White Splendour**’ illus. p.414.

A. coronaria. Spring-flowering perennial with a misshapen tuber. H 12–18in (30–45cm), S 6in (15cm). Produces parsleylike, divided, semierect leaves. Each stiff stem bears a large, 5–8-petaled, shallowly cup-shaped flower in shades of red, pink, blue or purple. Z8–11. Garden groups include **De Caen Group** and **Saint Bridgid Group**, which have larger flowers varying in color from white through red to blue.

‘**Mister Fokker**’ illus. p.403.

A. x fulgens illus. p.418.

A. hepatica. See *Hepatica nobilis*.

A. hupehensis. Erect perennial with a creeping, woody-based rootstock. H 24–36in (60–90cm), S 16in (40cm) or more. Large, rounded, dark green leaves have 3 toothed lobes. In late summer, flower stems produce smaller leaves and 5-petaled, slightly cupped, pale pink or white flowers, 2–2½in (5–6cm) across, with yellow stamens. Z5–7. ‘**Bowles’s Pink**’ (illus. p.222) has rich purple-pink flowers. ‘**Hadspen Abundance**’ illus. p.247. **var. japonica ‘Bressingham Glow’** (syn. *A. x hybrida* ‘Bressingham Glow’; illus. p.222), H 4–5ft (1.2–1.5m), S 2ft (60cm), bears semidouble, rose-purple flowers on wiry stems. **var. japonica ‘Pamina’** (illus. p.222), H 32in (80cm), produces abundant, rather small, semidouble, reddish-purple flowers. **var. japonica ‘Prinz Heinrich’** (syn. *A. x hybrida* ‘Prince Henry’) bears single, deep pink flowers on slender stems. ‘**Praecox**

(illus. p.222), H 32in (80cm), bears single, pink flowers in summer. ‘**Rotkäppchen**’, H 32–48in (80–120cm), is compact, with deep reddish-pink flowers. ‘**September Charm**’ see *A. x hybrida* ‘September Charm’.

A. x hybrida, syn. *A. japonica* of gardens (Japanese anemone). Group of vigorous, branching, perennials. H 4–5ft (1.2–1.5m), S indefinite. Bears shallowly cup-shaped, single, semidouble or double flowers in late summer and early fall. Leaves are deeply divided and dark green. Z4–8.

‘**Bressingham Glow**’ see *A. hupehensis* var. *japonica* ‘Bressingham Glow’.

‘**Elegans**’ (syn. *A. x h. ‘Max Vogel’*) has semidouble, pinkish-mauve flowers on wiry stems. ‘**Honorine Jobert**’ (single) illus. p.220. ‘**Königin Charlotte**’ (illus. p.222), H 5ft (1.5m), has large, semidouble, bright pink flowers. ‘**Lady Gilmour**’, H 3ft (1m), produces crested leaves and pink flowers with often uneven, twisted petals. ‘**Lady Gilmour**’ of gardens, see *A. x h. ‘Montrose*’. ‘**Margarete**’ of gardens, see *A. x h. ‘Montrose*’. ‘**Max Vogel**’ see *A. x h. ‘Elegans*’. ‘**Montrose**’ (syn. *A. x h. ‘Lady Gilmour*’ of gardens; *A. x h. ‘Margarete*’ of gardens)H 32in–4ft (80cm–1.2m), has large, double, soft reddish-pink flowers with rather twisted petals. ‘**Prince Henry**’ see *A. hupehensis* var. *japonica* ‘Prinz Heinrich’. ‘**Richard Ahrens**’, H 32–39in (80–100cm), bears large, single, soft pink flowers in midsummer. ‘**Robustissima**’ (illus. p.222), H 3–5ft (1–1.5m), S 4–20in (10–50cm), has single, dark pink flowers borne on reddish stems. ‘**September Charm**’ (syn. *A. hupehensis* ‘September Charm’; illus. p.222), H 30in (75cm), S 20in (50cm), has single, clear pink flowers on wiry stems. ‘**Whirlwind**’ (syn. *A. x h. Wirbelwind*; illus. p.222), H 32–39in (80–100cm), has irregularly formed double, white flowers, with some petals green flushed.

Wirbelwind see *A. x h. ‘Whirlwind*’.

A. x intermedia. See *A. x lipsiensis*.

A. japonica of gardens. See *A. x hybrida*.

A. x lipsiensis, syn. *A. x intermedia*, *A. x seemannii*, illus. p.262. ‘**Pallida**’ is a prostrate, carpeting perennial. H 2–6in (5–15cm), S to 18in (45cm). Produces single, pale cream flowers above deeply cut, mid-green foliage in spring. Z5–8.

A. narcissiflora illus. p.255.

A. nemorosa (Wood anemone). Vigorous, carpeting, rhizomatous perennial. H 3–6in (8–15cm), S 12in (30cm). Produces abundant, star-shaped, single, white flowers, with prominent, yellow stamens, in spring and early summer, above deeply cut, mid-green leaves. Likes woodland conditions. Z4–8. ‘**Allenii**’ produces many large, cup-shaped, single, rich lavender-blue flowers in spring. ‘**Bowles’ Purple**’ is late-flowering, sometimes with purple-edged leaves and purple flowers.

‘**Lychette**’ is early-flowering, with white flowers. ‘**Robinsoniana**’ has lavender-blue flowers, pale creamy-gray beneath, on maroon stems. ‘**Vestal**’ has anemone-centered, double, white flowers.

‘**Virescens**’ is infertile and flowers are replaced by rosettes of leaflike green tepals. ‘**Wilks’ Giant**’ (syn. *A.n. ‘Wilk’s Giant*’) has larger, single, white flowers.

A. pavonina illus. p.402.

ANEMOPAEGMA

A. ranunculooides illus. p.263.

‘**Pleniflora**’ (syn. *A.r. ‘Flore Pleno*’) is a spreading, rhizomatous perennial. H 2–4in (5–10cm), S to 18in (45cm) or more. Bears buttercup-like, double, yellow flowers in spring. Leaves are divided. Likes damp, woodland conditions. Z4–8.

A. rivularis illus. p.437.

A. x seemannii. See *A. x lipsiensis*.

A. sylvestris (Snowdrop windflower) illus. p.255.

A. tomentosa, syn. *A. vitifolia* of gardens. Vigorous, clump-forming perennial with a creeping rootstock. H 3–5ft (1–1.5m), S 2ft (60cm). Large, 3–7 lobed mid-green leaves with toothed margins have white-hairy undersides. In summer, flower stems bear 5-petaled, slightly cupped, pale pink flowers, 2–2½in (5–6cm) across, with yellow stamens. Z5–9.

A. trullifolia illus. p.350.

A. tschaernjaewii illus. p.416.

A. vitifolia. Branching, clump-forming perennial. H 3ft (1m), S indefinite. Has vinelike leaves, woolly beneath, and open cup-shaped, occasionally pink-flushed, white flowers with yellow stamens in summer. Z5–8.

A. vitifolia of gardens. See *A. tomentosa*.

ANEMONELLA

RANUNCULACEAE

Genus of one species of tuberous perennial, grown for its flowers. Needs shade and humus-rich, moist soil. Propagate by seed when fresh or by division every 3–5 years in fall.

A. thalictroides illus. p.350. ‘**Oscar Schoaf**’ (syn. *A.t. ‘Schoaf’s Double*’) is a slow-growing, tuberous perennial. H 4in (10cm), S 12in (30cm). Has delicate, fernlike leaves. From spring to early summer bears small, cup-shaped, double, strawberry-pink flowers, singly on finely branched, slender stems. Z4–7.

ANEMONOPSIS

False anemone

RANUNCULACEAE

Genus of one species of clump-forming perennial, related to *Anemone*. Likes a sheltered, semishaded position and humus-rich, moist but well-drained soil. Propagate by division in spring or by seed sown in late summer, when fresh.

A. macrophylla (False anemone) illus. p.270.

ANEMOPAEGMA

BIGNONIACEAE

Genus of evergreen, tendril climbers, grown for their flowers. Needs partial shade in summer and humus-rich, well-drained soil. Water regularly and freely when in full growth, less at other times. Provide support, and in summer thin out stems at intervals; shorten all growths by half in spring. Propagate by softwood or semiripe cuttings in spring or summer.

A. chamberlaynei. Fast-growing, evergreen, tendril climber. H 12–20ft (4–6m). Leaves have 2 pointed, ovate leaflets and a 3-hooked tendril. Foxglovelike, primrose-yellow flowers are borne in upper leaf axils in summer. Z12–15. (min. 45°F/7°C)