

‘Filifera Aurea’ (illus. p.105), H 40ft (12m), S 10–15ft (3–5m), also has whiplike shoots, but with golden-yellow leaves. **‘Filifera Nana’**, H 2ft (60cm), S 3ft (1m), is a dwarf form with whiplike branches. **‘Nana’**, H and S 20in (50cm), is also dwarf, with dark bluish-green foliage. **‘Plumosa’** is broadly conical to columnar, with yellowish, gray-green leaves. **‘Plumosa Rogersii’**, H 6ft (2m), S 3ft (1m), has yellow foliage. Slow-growing **‘Squarrosa’**, H to 70ft (20m), has a broad crown and soft, blue-gray foliage. **C. thyoides**, syn. *Cupressus thyoides* (White cypress), illus. p.101. **‘Andelyensis’** is a slow-growing, conical, dwarf conifer. H 10ft (3m), S 3ft (1m). Has wedge-shaped tufts of scalelike, aromatic, blue-green leaves. Female cones are glaucous blue-gray, later purple-black to red-brown. Z4–9

CHAMAECYTISUS
FABIACEAE

Genus of evergreen or deciduous, small trees, shrubs, and subshrubs, occasionally spiny, grown for their pealike flowers. Best in full sun and moderately fertile, well-drained soil. Propagate by seed in fall or spring, or by semiripe cuttings in summer. **C. albus**. See *Cytisus albus*. **C. hirsutus**, syn. *Cytisus hirsutus*. Deciduous, bushy, rounded shrub. H and S 3ft (1m). Dense, terminal heads of large, yellow flowers are borne from midsummer to fall, amid gray-green leaves with 3-palmate, oblong to elliptic leaflets. Z5–7 **C. purpureus**, syn. *Cytisus purpureus* (Purple broom). Deciduous, arching shrub. H 18in (45cm), S 2ft (60cm). Semierect stems are clothed with leaves of 3-palmate, obovate leaflets. Clusters of 2 or 3 pale lilac to purple flowers open in early summer on previous year’s wood. Suits a bank or sunny border. Z5–7. **f. albus** illus. p.337.

CHAMAEDAPHNE
SYN. CASSANDRA
ERICACEAE

Genus of one species of evergreen shrub, grown for its white flowers. Needs sun or semishade and moist, peaty, acidic soil. Propagate by semiripe cuttings in summer. **C. calyculata** (Leatherleaf). Evergreen, arching, open shrub. H 2½ft (75cm), S 3ft (1m). Leaves are small, oblong, leathery and dark green. Leafy racemes of small, urn-shaped, white flowers are borne on slender branches in mid- to late spring. Z6–9

CHAMAEDOREA
ARECACEAE/PALMAE

Genus of evergreen palms, grown for their overall appearance. Needs shade or semishade and humus-rich, well-drained soil. Water containerized plants moderately, less when temperatures are low. Propagate by seed in spring at not less than 25°C (77°F). Red spider mite may be troublesome. **C. elegans**, syn. *Neanthe bella* (Dwarf mountain palm, Parlour palm), illus. p.458.

CHAMAEMELUM
Chamomile
ASTERACEAE/COMPOSITAE

Genus of aromatic annuals and perennials, suitable as groundcover or for a lawn. Flowers may be used to make tea. Needs sun and well-drained soil. Propagate by division in spring or by seed in fall. **C. nobile**, syn. *Anthemis nobilis* (Lawn chamomile, Roman chamomile). Evergreen, mat-forming, invasive perennial. H 4in (10cm), S 18in (45cm). Has finely divided, aromatic leaves and daisylike heads of white flowers, with yellow centers, borne in late spring or summer. ☉ Contact with foliage may aggravate skin allergies. Z4–9. **‘Treneague’** is a non-flowering, less invasive cultivar that, requiring less mowing, is better for a lawn.

CHAMAENERION
SYN. CHAMERION
ONAGRACEAE

Genus of woody-based, herbaceous perennials, grown for their pink or white flowers that are borne for long periods from summer to fall. Some species are invasive, but *C. angustifolium* is good for a rock garden or border. Tolerates sun or shade and prefers moist, but well-drained soil. Propagate by seed in spring, or by softwood cuttings from sideshoots in spring. **C. angustifolium f. album**, syn. *Epilobium angustifolium f. album* (White rosebay), illus. p.216.

Chamaepericlymenum canadense.
See Cornus canadensis.

CHAMAEROPS
Dwarf fan palm
ARECACEAE/PALMAE

Genus of one species of shrubby palm, cultivated for its overall appearance. Needs full light and fertile, well-drained soil. Water containerized plants moderately, less when not in full growth. Propagate by seed in spring at not less than 22°C (72°F) or by suckers in late spring. Red spider mite may be a nuisance. **C. humilis** illus. p.165.

Chamaespartium sagittale. See Genista sagittalis.
Chamaespartium sagittale subsp. delphinense. See Genista delphinensis.

CHAMELAUCIUM
MYRTACEAE

Genus of evergreen shrubs, grown for their flowers and overall appearance. Needs full sun and well-drained, sandy, neutral to acidic soil. Water potted specimens moderately, sparingly when not in full growth. To maintain a compact habit, cut back flowered stems by half when the last bloom falls. Propagate by seed in spring or by semiripe cuttings in summer. **C. uncinatum** (Geraldton waxflower) [white] illus. p.453. **‘Bundara Excelsior’** illus. p.454.

Chamerion. See Chamaenerion.
CHASMANTHE
IRIDACEAE

Genus of cormous perennials, grown for their showy flowers. Needs full sun or partial shade and well-drained soil. Water well in late winter and early spring; reduce watering in summer and fall. Propagate by division in fall. **C. bicolor**. Spring-flowering corm. H 36in (90cm), S 20in (50cm). Has lance-shaped or linear leaves and spikelike, one-sided racemes of orange-red flowers, with green markings, maroon throats, and yellow-striped tubes. Z9–11 **C. floribunda**. Summer-flowering corm. H 20–60in (50cm–1.5m), S 6in (15cm). Bears lance-shaped leaves and numerous bright orange or yellow flowers, arranged in 2 ranks, on branched spikes. Z10–11. **var. duckittii** has yellow flowers.

CHASMANTHIUM
POACEAE/GRAMINEAE

See also GRASSES, BAMBOOS, RUSHES and SEDGES. **C. latifolium** illus. p.288.

CHEILANTHES
Lip fern
PTERIDACEAE

Genus of mainly evergreen ferns. Needs full light and humus-rich, well-drained soil. Do not overwater containerized plants or splash water on fronds. Remove fading foliage regularly. Propagate by spores in summer. **C. lanosa** of gardens. See *C. tomentosa*. **C. tomentosa**, syn. *C. lanosa* of gardens. Evergreen fern. H and S 6–9in (15–23cm). Leaves are triangular or lance-shaped and have much divided, soft green fronds on hairy, black stems. Z5–8

Cheiranthus. See Erysimum.

CHEIRIDOPSIS
AIZOACEAE

Genus of clump-forming, perennial succulents with pairs of semicylindrical leaves. Needs sun and well-drained soil. Water in fall to encourage flowers. Propagate by seed or stem cuttings in spring or summer. **C. candidissima**. See *C. denticulata*. **C. denticulata**, syn. *C. candidissima*. Clump-forming, perennial succulent. H 4in (10cm), S 8in (20cm). Has semicylindrical, slender, fleshy, blue-gray leaves, each 4in (10cm) long with a flat top, joined in pairs for almost half their length. Bears daisylike, shiny, white to pale yellow flowers, to 2½in (6cm) across, in spring. Z10–11 (min. 41°F/5°C) **C. purpurata**. See *C. purpurea*. **C. purpurea**, syn. *C. purpurata*. Carpeting, perennial succulent. H 4in (10cm), S 12in (30cm). Has semicylindrical, thick, short, glaucous green leaves, each with a flat top. In early spring, produces daisylike, purple-pink or yellow flowers, 1½in (4cm) across. Z10–11 (min. 41°F/5°C)

CHIONANTHUS
CHELIDONIUM
Celandine, Greater Celandine
PAPAVERACEAE

Genus of one species of variable biennial or short-lived perennial that rapidly forms groundcover. Grows in sun or shade and in any but very wet soil. Can be invasive; suitable for naturalizing in a wildflower garden or in light woodland. Propagate by seed or division in fall. ☉ Contact with the sap may cause skin blisters. **C. majus ‘Flore Pleno’** illus. p.227.

CHELONE
Turtle-head
PLANTAGINACEAE

Genus of summer- and fall-flowering perennials. Needs semishade and moist soil. Ideal for a late summer border. Propagate by soft-tip cuttings in summer or by division or seed in fall or spring. **C. barbata**. See *Penstemon barbatus*. **C. obliqua** (Turtle-head) illus. p.438.

Chiastophyllum oppositifolium. See Umbilicus oppositifolius.

CHIMONANTHUS
Wintersweet
CALYCANTHACEAE

Genus of deciduous or evergreen, winter-flowering shrubs, grown for their very fragrant flowers, borne in winter before the young leaves emerge. Reduce susceptibility of flowers to frost damage by training plants against a warm, sunny wall. Needs full sun and fertile, well-drained soil. Propagate species by seed when ripe, in late spring and early summer, cultivars by softwood cuttings in summer. **C. fragrans**. See *C. praecox*. **‘Luteus’** see *C. praecox* ‘Luteus’. **C. praecox**, syn. *C. fragrans* (Wintersweet), illus. p.144. **‘Concolor’** see *C.p.* ‘Luteus’. **‘Luteus’** (syn. *C. fragrans* ‘Luteus’, *C. praecox* ‘Concolor’) is a vigorous, broadly upright, deciduous shrub. H 12ft (4m), S 10ft (3m). Has long, lance-shaped, glossy, mid-green leaves, rough above, smooth beneath. Pendent, fragrant, wide-opening, pure yellow flowers are borne on bare stems in winter. Z7–9

CHIMONOBAMBUSA
POACEAE/GRAMINEAE

See also GRASSES, BAMBOOS, RUSHES, and SEDGES. **C. tumidissinoda**, syn. *Qiongzhuea tumidissinoda*, illus. p.287.

CHIONANTHUS
Fringe tree
OLEACEAE

Genus of evergreen and deciduous trees and shrubs, grown for their profuse, white flowers. Flowers more freely in areas with hot summers. Good specimen plants and also suitable for a shrub border. Prefers full sun and fertile,