

COTYLEDON

***C. tomentosa* subsp. *ladismithensis*.** Evergreen, freely branching, later prostrate, succulent subshrub. H and S 8in (20cm). Has fleshy, green leaves, swollen and blunt at tips and covered with short, golden-brown hairs. Clusters of tubular, brownish-red flowers are produced in fall. Z10–12 (min. 41°F/5°C)

***C. wallichii*.** See *Tylecodon wallichii*.

CRAMBE

BRASSICACEAE/CRUCIFERAE

Genus of annuals and, often woody-based, perennials, grown for their bold leaves and large sprays of white flowers in summer. Leaf shoots of *C. maritima* (sea kale) are eaten as a spring vegetable. Will grow in any well-drained soil; prefers an open position in full sun but tolerates some shade. Propagate by division in spring or by seed in fall or spring.

C. cordifolia illus. p.216.

C. maritima (Sea kale) illus. p.264.

CRASPEDIA

ASTERACEAE/COMPOSITAE

Genus of basal-rosetted, summer-flowering annuals and perennials, grown for their flowers and foliage. Flower heads are useful for dried arrangements. Requires a site in sun and well-drained soil. Suitable for a rock garden or scree bed. Propagate by seed sown when very fresh in summer.

***C. incana*.** Basal-rosetted perennial. H 8–12in (20–30cm), S 4in (10cm). Has narrowly ovate, basal leaves, with dense, woolly, white hairs beneath, and smaller leaves on flower stems. In summer, many domed heads of 3–10 tiny, tubular, yellow flowers are produced in large, terminal clusters. Z9–11

CRASSULA

CRASSULACEAE

Genus of annual and perennial succulents and evergreen, succulent shrubs and subshrubs, ranging from very succulent-leaved species, ¾in (2cm) high, to shrubby types, 15ft (5m) high. Most are easy to grow; in frost-prone areas, grow as houseplants. Most prefer full sun; others like partial shade. Needs very well-drained soil and a little water in winter. Propagate by seed or stem cuttings in spring or fall.

C. arborescens (Silver jade plant) illus. p.490.

C. argentea of gardens. See *C. ovata*.

***C. coccinea*,** syn. *Rochea coccinea*. Evergreen, erect, succulent shrub. H to 2ft (60cm), S 12in (30cm) or more. Alternate pairs of fleshy, ovate to oblong-, hairy-margined, dull green leaves, each united at the base, are arranged at right angles in 4 rows up the woody, green stems. Produces umbels of tubular, bright red flowers in summer or fall. Z10–11 (min. 41–45°F/5–7°C)

***C. deceptor*,** syn. *C. deceptrix*, illus. p.488.

***C. deceptrix*.** See *C. deceptor*.

***C. falcata*.** See *C. perfoliata* var. *minor*.

***C. lactea*.** Prostrate to semierect, perennial succulent. H 8in (20cm), S 3ft (1m). Leaves are triangular-ovate, glossy

and dark green. In winter, produces an abundance of small, 5-petaled, white flowers in terminal clusters. Likes partial shade. Z10–11 (min. 41–45°F/5–7°C)

***C. lycopodioides*.** See *C. muscosa*.

C. multicava illus. p.484.

***C. muscosa*,** syn. *C. lycopodioides*. Dense, bushy, woody-based, perennial succulent. H 6in (15cm), S 12in (30cm). Bears small, scalelike, neatly overlapping, mid-green leaves arranged in 4 rows around erect stems. In spring, produces tiny, 5-petaled, greenish-yellow flowers. Likes partial shade. Z10–11 (min. 41–45°F/5–7°C)

***C. ovata*,** syn. *C. argentea* of gardens, *C. portulacea* (Friendship tree, Jade tree, Money tree), illus. p.481.

***C. perfoliata* var. *minor*,** syn. *C. falcata* (Aeroplane propellor), illus. p.489.

***C. portulacea*.** See *C. ovata*.

C. sarcocaulis illus. p.339.

C. schmidtii illus. p.485.

C. socialis illus. p.480.

+ CRATAEGOMESPILUS

ROSACEAE

Group of grafted, hybrid, deciduous trees (*Crataegus* and *Mespilus*), grown for their flowers, foliage and fruits. Requires sun or semishade and fertile, well-drained soil. Propagate by grafting in late summer.

+ *C. dardarii* (Bronvaux medlar). ‘**Jules d’Asnières**’ is a deciduous, spreading tree. H and S 20ft (6m). Has drooping branches and spiny shoots. Variable, ovate or deeply lobed, dark green leaves, gray when young, turn orange and yellow in fall. Clusters of saucer-shaped, white, sometimes rose-tinted, flowers in late spring or early summer are followed by small, rounded, red-brown fruits. Z8–10

CRATAEGUS

Hawthorn, Thorn

ROSACEAE

Genus of deciduous, or more rarely semievergreen, spiny, often spreading trees and shrubs, grown for their clustered, 5-petaled, occasionally double flowers in spring–summer, ornamental fruits and, in some cases, fall color. Prefers full sun but is suitable for most sites and may be grown in any but very wet soil. Useful for growing in polluted urban areas, exposed sites and coastal gardens. Propagate species by seed in fall, cultivars by budding in late summer. Fireblight is sometimes a problem. ☉The seeds may cause mild stomach upset if ingested.

***C. coccinea*,** syn. *C. pedicellata*, illus. p.90.

***C. cordata*.** See *C. phaenopyrum*.

C. crus-galli (Cockspur thorn). Deciduous, flat-topped tree. H 25ft (8m), S 30ft (10m). Has shoots armed with long, curved thorns and ovate, glossy, dark green leaves that turn bright crimson in fall. Bears clusters of white flowers, with pink anthers, in late spring, followed by long-lasting, rounded, bright red fruits. Z3–7

C. crus-galli of gardens. See *C. persimilis* ‘Prunifolia’.

***C. elwangeriana*.** Deciduous, spreading tree. H and S 20ft (6m). Broadly ovate, dark green leaves are shallowly toothed

and lobed. Bears clusters of white flowers, with pink anthers, in late spring, followed by rounded, glossy, crimson fruits. Z3–7

***C. x grignonensis*.** Small, deciduous or semievergreen, spreading tree. H 15ft (5m), S 12ft (4m). Has almost thornless shoots with dark green leaves, glossy above, paler and more densely hairy beneath. Many small white flowers, which fade to pink, are produced in clusters in late spring to early summer, followed by showers of late-ripening, shiny, red fruits, which often persist into spring. Z7–10

C. laciniata of gardens. See *C. orientalis*.

***C. laevigata*,** syn. *C. oxyacantha* of gardens (Hawthorn, May). ‘**Paul’s Scarlet**’ illus. p.84. ‘**Punicea**’ is a deciduous, spreading tree. H and S 20ft (6m). In late spring and early summer, bears clusters of crimson flowers amid ovate, lobed, toothed, glossy, dark green leaves, followed by rounded, red fruits. ‘**Rosea Flore Pleno**’ bears double pink flowers. Z4–8

***C. x lavallei* ‘Carrierei’.** Vigorous, deciduous, spreading tree. H 22ft (7m), S 30ft (10m). Ovate, glossy, dark green leaves turn red in late fall. Bears clusters of white flowers in late spring, followed by long-lasting, rounded, orange-red fruits. Z4–8

***C. mollis*.** Deciduous, spreading tree. H 30ft (10m), S 40ft (12m). Large, broadly ovate, lobed, dark green leaves have white-haired undersides when young. Produces heads of large, white flowers in late spring, followed by short-lived, rounded, red fruits. Z4–8

C. monogyna (Common hawthorn). Deciduous, round-headed tree. H 30ft (10m), S 25ft (8m). Has broadly ovate, deeply lobed, glossy, dark green leaves. Clusters of fragrant, white flowers are produced from late spring to early summer, followed by rounded, red fruits. Will form a dense hedge. Z4–8. ‘**Biflora**’ (Glastonbury thorn) has flowers and leaves in mild winters as well as in spring.

***C. orientalis*,** syn. *C. laciniata* of gardens, illus. p.80.

C. oxyacantha of gardens. See *C. laevigata*.

***C. pedicellata*.** See *C. coccinea*.

***C. x persimilis* ‘Prunifolia’,** syn. *C. crus-galli* of gardens, *C. x prunifolia*. Deciduous, spreading, thorny tree. H 25ft (8m), S 30ft (10m). Ovate, glossy, dark green leaves turn red or orange in fall. Has clusters of white flowers, with pink anthers, in early summer, then rounded, dark red fruits. Z6–8

***C. phaenopyrum*,** syn. *C. cordata* (Washington thorn). Deciduous, round-headed tree. H and S 30ft (10m). Broadly ovate leaves are sharply lobed, glossy and dark green. Clusters of white flowers, with pink anthers, are produced from early to midsummer, followed by rounded, glossy, red fruits that last through winter. Z4–8

***C. x prunifolia*.** See *C. persimilis* ‘Prunifolia’.

C. tanacetifolia (Tansy-leaved thorn). Deciduous, upright, usually thornless tree. H 30ft (10m), S 25ft (8m). Has ovate to diamond-shaped, deeply cut, gray-green leaves, clusters of fragrant, white flowers, with red anthers, in midsummer, and small, apple-shaped, yellow fruits. Z6–8

CREMANTHODIUM

ASTERACEAE/COMPOSITAE

Genus of basal-rosetted perennials, grown for their pendent, half-closed, daisylike flower heads. Grow in woodland, a rock garden, or an alpine house. Dislikes winter wet. Needs shade and humus-rich, moist but well-drained soil. Propagate by seed when fresh.

***C. reniforme*.** Basal-rosetted perennial. H and S 8in (20cm). Leaves are large, kidney-shaped, toothed, glossy, mid-green. Sturdy stems each bear a large, daisylike, yellow flower head in summer. Z5–8

CREPIS

Hawk’s beard

ASTERACEAE/COMPOSITAE

Genus of summer-flowering annuals and perennials with long tap roots and leaves in flat rosettes. Many species are persistent weeds, but some are grown for their many-petaled, dandelionlike flower heads. Tolerates sun or shade and prefers well-drained soil. Propagate annuals and biennials by seed in fall, perennials by root cuttings (not from tap root) in late winter; most species self-seed freely.

C. aurea illus. p.345.

C. incana (Pink dandelion). Basal-rosetted perennial. H 8in (20cm), S 4in (10cm). Bears oblong, divided, hairy, grayish-green leaves. Uneven discs of ragged, pink flower heads are borne on stiff stems in summer. Good for a sunny rock garden or border. Z3–9

***C. rubra*.** Fairly fast-growing, rosette-forming annual. H 12in (30cm), S 6in (15cm). Has lance-shaped, serrated leaves. In summer, bears dandelionlike, pink, occasionally red or white flower heads. Z3–10

CRINODENDRON

ELAEOCARPACEAE

Genus of evergreen shrubs and trees, grown for their flowers and foliage. Requires shade or semishade, with plant base in cool shade, and fertile, moist but well-drained, acidic soil. Propagate by softwood cuttings in summer or by seed in fall.

***C. hookerianum*,** syn. *Tricuspidaria lanceolata* (Lantern tree), illus. p.202.

C. patagua illus. p.113.

x ***Crinodonna*.** See x *Amarcrinum*.

CRINUM

AMARYLLIDACEAE

Genus of robust bulbs, grown for their often fragrant, funnel-shaped flowers. Needs full sun, shelter and rich, well-drained soil. Propagate by offsets in spring or by seed when fresh or in spring. ☉All parts may cause severe discomfort if ingested; contact with the sap may irritate skin.

***C. bulbispermum*,** syn. *C. longifolium*. Summer-flowering bulb. H to 3ft (1m), S 2ft (60cm). Leafless flower stem bears a head of fragrant, long-tubed, white or pinkish-red flowers with darker red stripes. Long, strap-shaped, semierect leaves are grouped in a tuft on a short stalk. Z7–10