

C. longifolium. See *C. bulbisperrum*.
C. macowanii. Fall-flowering bulb. H and S 2ft (60cm) or more. Similar to *C. bulbisperrum*, but leaves are wavy-edged. Z9–11
C. moorei illus. p.383.
C. x powellii illus. p.385. ‘**Album**’ illus. p.383.

CROCOSMIA

Montbretia

IRIDACEAE

Genus of cormous perennials, grown for their brightly colored flowers, borne mainly in summer. Forms dense clumps of sword-shaped, erect leaves. Requires well-drained soil and an open, sunny site. In very cold areas, plant in a sheltered position or lift and store corms over winter. Propagate by division as growth starts in spring.
C. aurea. Tuft-forming, summer-flowering corm. H 20–30in (50–75cm), S 6–8in (15–20cm). Erect, basal leaves are long, narrow and sword-shaped. Bears a loosely branched spike of tubular, orange or yellow flowers, each 1–2in (3–5cm) long and with 6 spreading petals. Z8–11
C. ‘Bressingham Blaze’. Clump-forming, late summer-flowering corm. H 30in (75cm), S 6–8in (15–20cm). Produces sword-shaped, pleated, basal, erect leaves. Branched stem bears widely funnel-shaped, fiery-red flowers. Z8–11
C. x crocosmiiflora. Robust, sometimes invasive, variable, late summer-flowering corm. H 2ft (60cm), S 3in (8cm). Has erect, sword-shaped, pale green, basal leaves. Produces thin, slightly arching, sometimes branched spikes of funnel-shaped, orange or yellow flowers in summer. Z6–10.
‘Buttercup’ produces rich yellow flowers that open flat; Z6–10. **‘Coleton Fishacre’** (syn. *C. ‘Golden Fleece’*) illus. p.412.
‘Emily McKenzie’ (syn. *C. ‘Emily McKenzie’*), H to 2ft (60cm), S 6–8in (15–20cm), is compact, with a dense spike of widely funnel-shaped, deep orange flowers, each with a dark mahogany throat; Z6–10.
‘George Davison’ (syn. *C. ‘George Davison’*; illus. p.410), H 3–4ft (90–120cm), S 6in (15cm), produces branched stems bearing large, trumpet-shaped, pale orange-yellow flowers, tinted deeper orange externally; Z6–10. **‘Harlequin’** (syn. *C. ‘Harlequin’*), H 36in (90cm), S 3in (8cm), has well-branched, upright stems bearing funnel-shaped, bright yellow flowers with alternate, red and orange outer petals; Z6–10. **‘Honey Angels’** (syn. *C. ‘Honey Angels’*; illus. p.410), H 30in (75cm), S 6in (15cm), has sword-shaped, erect, pleated, bronzed-green leaves and trumpet-shaped, pale yellow flowers; Z6–10. **‘Jackanapes’** (syn. *C. ‘Jackanapes’*; illus. p.410), H 16–24in (40–60cm), S 6–8in (15–20cm), produces striking bicolored, yellow and orange-red flowers; Z6–10.
‘Lady Hamilton’ (syn. *C. ‘Lady Hamilton’*), H 24–30in (60–75cm), bears golden yellow flowers with apricot-yellow centers, on erect, branched spikes in late summer; Z6–10. **‘Solfatare’** (syn. *C. ‘Solfatare’*; illus. p.410), H 26–28in (65–70cm), S 6in (15cm), has trumpet-shaped pale to mid-yellow flowers; Z6–10. **‘Star of the East’** (syn. *C. ‘Star of the East’*; illus. p.410), H 28in (70cm), S 3in (8cm), bears horizontal-

facing, funnel-shaped, clear orange flowers, with a paler orange center, on branched stems; Z6–10.
C. ‘Emily McKenzie’. See *C. x crocosmiiflora* ‘Emily McKenzie’.
C. ‘George Davison’. See *C. x crocosmiiflora* ‘George Davison’.
C. ‘Golden Fleece’. See *C. x crocosmiiflora* ‘Coleton Fishacre’.
C. ‘Harlequin’. See *C. x crocosmiiflora* ‘Harlequin’.
C. ‘Honey Angels’. See *C. x crocosmiiflora* ‘Honey Angels’.
C. ‘Jackanapes’. See *C. x crocosmiiflora* ‘Jackanapes’.
C. ‘Lady Hamilton’. See *C. x crocosmiiflora* ‘Lady Hamilton’.
C. ‘Lucifer’ (illus. p.410). Robust, clump-forming corm. H to 3ft (1m), S 8–10in (20–25cm). Has sword-shaped, erect, basal, bright green leaves. In midsummer, bears funnel-shaped, deep rich red flowers in dense, branching spikes. Z6–9
C. masoniorum, syn. *C. masonorum* (illus. p.410). Robust, clump-forming corm. H to 5ft (1.5m), S 1–1½ft (30–45cm). Has erect, basal, deep green leaves, pleated lengthways. Erect, branched stem has a horizontal upper part, which bears reddish-orange flowers in summer-fall. Z6–10. **‘Rowallane Yellow’**, H 3ft (1m), S 3in (8cm), has upward-facing, funnel-shaped, yellow flowers.
C. masonorum. See *C. masoniorum*.
C. paniculata, syn. *Antholyza paniculata*, *Curtonus paniculatus*. Summer-flowering corm. H to 5ft (1.5m), S 1–1½ft (30–45cm). Has sword-shaped, erect, basal leaves, pleated lengthways. Produces long-tubed, orange flowers on branched stems, which are strongly zig-zag in shape. Z5–9
C. ‘Severn Sunrise’ (illus. p.410). Late summer-flowering corm. H 36in (90cm), S 3in (8cm). Has erect, sword-shaped, mid-green, basal leaves. Produces tightly clustered, funnel-shaped flowers in shades of salmon, apricot and yellow. Z5–9
C. ‘Solfatare’. See *C. x crocosmiiflora* ‘Solfatare’.
C. ‘Star of the East’. See *C. x crocosmiiflora* ‘Star of the East’.

CROCUS

IRIDACEAE

Genus of mainly spring- or fall-flowering cormous perennials, grown for their funnel-shaped to rounded, long-tubed flowers. Produces long, very narrow, semierect, basal leaves, each with a white line along center, usually 1–5 per corm. Some fall-flowering species have no leaves at flowering time, these appearing in winter or spring. Most species are less than 4in (10cm) tall when in flower and have a spread of 1–3in (2.5–8cm). Ideal for rock gardens, for planting in drifts at the front of a border and for forcing in bowls for an early indoor display. The most vigorous can be naturalized in short turf. Most require well-drained soil and a sunny situation; *C. banaticus* prefers moist soil and semishade. Plant 2–2½in (5–6cm) deep, in late summer or early fall. Propagate in early fall by seed or division of corm clumps. See also feature panel p.417.
C. ‘Advance’. Late winter- to midspring-flowering corm. Funnel-shaped flowers

are buttercup yellow inside and paler yellow outside, suffused violet-bronze. Z4–8
C. aerius of gardens. See *C. biflorus* subsp. *pulchricolor*.
C. ancypensis. Spring-flowering corm. Produces up to 7 fragrant, bright orange-yellow flowers, 2–2½in (5–6cm) long. Z4–8
C. angustifolius, syn. *C. susianus* (Cloth-of-gold crocus). Spring-flowering corm. Produces 5 or more fragrant flowers, which are bright golden-yellow, and striped or stained bronze outside. Z4–8
C. aureus. See *C. flavus*.
C. balansae. See *C. olivieri* subsp. *balansae*.
C. banaticus, syn. *C. iridiflorus*. Fall-flowering corm, with solitary flowers. Usually has one long-tubed, pale violet flower; outer 3 petals open wide and are much larger than 3 erect, paler, inner petals. Dark green leaves, without central stripes, are borne after the flowers. Z4–8
C. baytopiorum. Spring-flowering corm. Each bears 1 or 2 rounded, turquoise-blue, slightly darker-veined flowers. Z4–8
C. biflorus. Very variable, early spring-flowering corm. Bears fragrant, white or purplish-white flowers, with yellow throats, the outer petals sometimes vertically striped purple or brownish purple. Z4–8. **subsp. alexandri** has fragrant white flowers, the outer petals heavily marked deep purple. **‘Blue Pearl’** (syn. *C. ‘Blue Pearl’*) illus. p.421. **subsp. pulchricolor** (syn. *C. aerius* of gardens) has rich deep blue flowers, stained dark violet near the bases, with yellow throats. **subsp. weldenii ‘Fairy’** has white flowers, the outer tepals dusted violet.
C. ‘Blue Bird’ (illus. p.417). Late winter- to midspring-flowering corm. Funnel-shaped flowers are white inside with deep yellow throats and violet margined with white outside. Z4–8
C. ‘Blue Pearl’. See *C. biflorus* ‘Blue Pearl’.
C. boryi. Fall-flowering corm. Bears up to 4 well-rounded, creamy-white flowers, sometimes veined or flushed with mauve outside, with the leaves. Z6–10
C. cancellatus. Very variable, fall-flowering corm. Produces 1–3, slender pale blue flowers, striped violet outside. Gray-green leaves form after flowering, in spring. Z4–8
C. cartwrightianus. Fall- and early winter-flowering corm. Produces leaves with, or shortly after, 1–5, open goblet-shaped, fragrant, violet or white flowers, strongly veined dark purple, with 3 long, bright red stigmas, similar to those of *C. sativus*. Z4–8
C. chrysanthus. Late winter and early spring-flowering corm. Has dull green leaves and bears up to 4 rounded, scented flowers, varying from cream to deep golden yellow, often suffused or veined bronze-maroon outside and with orange-red stigmas. Z4–8. **‘Cream Beauty’** see *C. ‘Cream Beauty’*. **‘Zwanenberg Bronze’** see *C. ‘Zwanenberg Bronze’*.
C. ‘Cream Beauty’ (illus. p.417), syn. *C. chrysanthus* ‘Cream Beauty’. Compact, spring-flowering corm. Scented, rich cream flowers have deep yellow throats, and are stained purplish-brown at the base. Z4–8
C. cvijicii. Spring-flowering corm. Usually has one funnel-shaped, yellow

flower, sometimes white or cream. Leaves are only just visible at flowering time. Z4–8
C. dalmaticus. Late winter and early spring-flowering corm. Bears 1–3 rounded, purple-veined, pale violet flowers, with yellow centers, overlaid with silver or yellow outside. Z4–8
C. ‘Dutch Yellow’. See *C. x luteus* ‘Golden Yellow’.
C. ‘E.A. Bowles’ (illus. p.417). Early spring-flowering corm. Bears compact, scented, funnel-shaped, rich lemon yellow flowers, the outer petals with bronze-green bases and purple feathering. Increases well by offsets. Z4–8
C. etruscus. Late winter and early spring-flowering corm. Bears 1 or 2 long-tubed, funnel-shaped, pale purple-blue flowers, washed silver outside, with violet veining. Z4–8. **‘Zwanenburg’** (illus. p.417) has pale purple-blue flowers, washed with biscuit-brown and flecked violet outside.
C. ‘Eyecatcher’ (illus. p.417). Late winter- to midspring-flowering corm. Funnel-shaped, gray-white, yellow-throated flowers are heavily marked deep purple outside. Z4–8
C. flavus, syn. *C. aureus*. Spring-flowering corm. Produces 1–4 fragrant, bright yellow or orange-yellow flowers; often several flowers are produced together or in quick succession. Z4–8
C. gargaricus. Spring-flowering corm. Bears solitary, slender, bright orange-yellow flowers. Increases by stolons. Tolerates slightly damper conditions than most crocuses. Z4–8
C. ‘Golden Yellow’. See *C. x luteus* ‘Golden Yellow’.
C. goulimyi (illus. p.417). Fall-flowering corm. Produces 1–3 rounded, long-tubed, scented, pale lilac to pinkish-lilac flowers, with a white throat and 3 inner petals usually paler than the 3 outer ones. Leaves and flowers are produced together. Needs a warm site. Z4–8
C. hadriaticus (illus. p.417). Fall-flowering corm. Leaves are produced with 1–3 white flowers, which usually have yellow throats and may be lilac-feathered at the base. Z4–8
C. imperati. Strikingly bicolored, late winter or early spring-flowering corm. Bears 1 or 2 scented, long-tubed, purple flowers, fawn with purple striping outside and with yellow throats. Leaves are shiny and dark green. Z4–8. **‘De Jager’** bears flowers rich violet-purple inside, biscuit-colored with violet feathering outside.
C. iridiflorus. See *C. banaticus*.
C. korolkowii (Celandine crocus). Late winter and early spring-flowering corm. Has up to 20 narrow leaves and bears 3–5 fragrant, yellow flowers that are speckled or stained brown or purple outside. When open in sun, petals have glossy surfaces. Z4–8
C. kotschyanus, syn. *C. zonatus* (illus. p.417). Fall-flowering corm. Solitary, long-tubed, pinkish-lilac or purplish-blue flowers, with yellow centers and white anthers, are borne before the leaves. Each short-stemmed flower has a ring of yellow dots around the throat and creamy-white stamens. Z4–8. **var. leucopharynx** has pale lilac-blue flowers with white throats.
C. ‘Ladykiller’. Late winter- to midspring-flowering corm. Has funnel-shaped flowers, white or pale lilac within,

C