

(2m), is an irregularly rounded shrub; foliage that turns bronze in winter. **‘Cristata’** illus. p.104. **‘Elegans Compacta’**, H 6–15ft (2–5m), S 6ft (2m), is a dwarf form. **‘Pyramidata’** illus. p.104. **‘Sekkan-sugi’**, H 30ft (10m), S 10–12ft (3–4m), has semipendent branches and light golden-cream foliage. **‘Spiralis’**, H and S 6–10ft (2–3m), forms a tree or dense shrub with spirally twisted foliage and is very slow-growing. **‘Vilmoriniana’**, H and S 3ft (1m), forms a globular mound of yellow-green foliage that turns bronze in winter.

**Cryptostemma calendulaceum.** See *Arctotheca calendula*.

CTENANTHE  
MARANTACEAE

Genus of evergreen, bushy perennials, grown for their ornamental foliage. Requires a humid atmosphere, even temperature and partial shade. Prefers moist but well-drained soil and soft water; do not allow to dry completely. Propagate by division in spring. **C. lubbersiana.** Evergreen, clump-forming, bushy perennial. H and S to 30in (75cm) or more. Long-stalked, lance-shaped, sharply pointed leaves are 10in (25cm) long, green above, irregularly marked and striped with pale yellowish-green, and pale greenish-yellow below. Intermittently bears dense, one-sided spikes of small, 3-petaled, white flowers. Z11–12 (min. 55°F/13°C) **C. oppenheimiana**, syn. *Calathea oppenheimiana*. Robust, evergreen, bushy perennial. H and S 3ft (1m) or more. Lance-shaped, leathery leaves are over 1ft (30cm) long, red below, dark green above with pale green or white bands along veins on either side of midribs. Dense, one-sided spikes of many small, 3-petaled, white flowers are borne intermittently. Z11–12. (min. 55°F/13°C). **‘Tricolor’** illus. p.465.

**Cudrania tricuspidata.** See *Maclura tricuspidata*.

CUNNINGHAMIA  
CUPRESSACEAE

See also CONIFERS.

**C. lanceolata** (Chinese fir) illus. p.100.

CUNONIA  
CUNONIACEAE

Genus of evergreen, summer-flowering trees, grown for their foliage, flowers and overall appearance. Requires full light and well-drained soil. Water potted plants moderately, less in winter. Pruning is tolerated. Propagate by seed in spring or by semiripe cuttings in summer. **C. capensis** (African red alder). Moderately fast-growing, evergreen, rounded tree. H and S 30–50ft (10–15m), more in rich soil. Has lustrous, dark green leaves, divided into pairs of lance-shaped, serrated leaflets. In late summer, tiny, long-stamened, white flowers are borne in dense, bottlebrushlike spikes, each 4–5in (10–13cm) long. Z10–11 (min. 41°F/5°C)

CUPHEA  
LYTHRACEAE

Genus of annuals, short-lived, sometimes subshrubby perennials and evergreen shrubs, grown for their flowers. Prefers full sun and fertile, well-drained soil. Water freely when in full growth, moderately at other times. Remove flowered shoots after flowering to maintain a bushy habit. Propagate by seed in spring or greenwood cuttings in spring or summer. Susceptible to red spider mite. **C. cyanea** illus. p.162. **C. ‘Firecracker’**. See *C. x purpurea* ‘Firecracker’. **C. ignea**, syn. *C. platycentra* (Cigar flower) illus. p.308. **C. platycentra**. See *C. ignea*. **C. x purpurea ‘Firecracker’** illus. p.306.

x **Cupressocyparis.** See x *Cuprocyparis*

CUPRESSUS  
Cypress  
CUPRESSACEAE

See also CONIFERS.

**C. arizonica** var. **glabra**, syn. *C. glabra* (Arizona cypress, Smooth cypress). Conical conifer. H 30–50ft (10–15m), S 10–15ft (3–5m). Has smooth, flaking, reddish-purple bark and upright, spirally arranged sprays of scalelike, aromatic, glaucous blue-gray leaves that are flecked with white resin. Globular cones are chocolate-brown. Z7–9 **C. cashmeriana**, syn. *C. torulosa* ‘Cashmeriana’ (Kashmir cypress) illus. p.95. **C. glabra**. See *C. arizonica* var. *glabra*. **C. lawsoniana**. See *Chamaecyparis lawsoniana*. **C. lusitanica** (Cedar of Goa, Mexican cypress). Conical conifer. H 70ft (20m), S 15–25ft (5–8m). Has fissured bark and spreading, spirally arranged sprays of scalelike, aromatic, gray-green leaves. Bears small, globular cones that are glaucous blue when young, ripening to glossy brown. Z8–10 **C. macrocarpa** (Monterey cypress). Fast-growing, evergreen conifer, columnar when young, often wide-spreading with age. H 70ft (20m), S 20–80ft (6–25m). Bark is shallowly fissured. Scalelike, aromatic, bright to dark green leaves are borne in plumelike sprays. Globular cones are glossy and brown. Z8–10. **‘Goldcrest’** illus. p.104. **C. pisifera**. See *Chamaecyparis pisifera*. **C. sempervirens ‘Stricta’** (Italian cypress), illus. p.102. **C. thyoides**. See *Chamaecyparis thyoides*. **C. torulosa ‘Cashmeriana’**. See *C. cashmeriana*.

x **CUPROCYPARIS**  
SYN. x CUPRESSOCYPARIS  
CUPRESSACEAE

Ⓢ Contact with the foliage may aggravate skin allergies. See also CONIFERS.

x **C. leylandii ‘Castlewellan’** illus. p.99. **‘Haggerston Grey’** illus. p.95. **‘Harlequin’** is a very fast-growing, columnar conifer with a conical tip. H 80–120ft (25–35m), S 12–15ft (4–5m). Gray-green foliage, with patches of clear ivory-white, is held in plumelike sprays.

**‘Leighton Green’** bears flattened sprays of paired, scalelike, rich green leaves and globular, glossy, dark brown cones. **‘Robinson’s Gold’**, H 50–70ft (15–20m), has bright golden leaves. Z6–10

CURCUMA  
Hidden ginger, Siam tulip  
ZINGIBERACEAE

Genus of herbaceous perennials, grown for their patterned leaves and showy bracts on the flower spikes. The lower bracts form pouches from which flowers emerge, the upper bracts are without flowers, but larger and brightly colored or white. Grow in humus-rich, moist but well-drained soil in partial shade. Keep completely dry in winter. Propagate by division in spring. **C. cordata**. See *C. zedoaria*. **C. petiolata** (illus p.477). Herbaceous, clump-forming perennial. H 4ft (1.2m), S 20in (50cm). Narrowly ovate, strongly ribbed, mid-green leaves are to 2ft (60cm) long. In summer, bears a spike, to 14in (35cm) long, of lemon-yellow flowers and bright pink upper bracts. Z12 **C. zedoaria**, syn. *C. cordata* (illus. p.477). Herbaceous, clump-forming perennial. H 60in (1.5m), S 20in (50cm). Has ovate, ribbed, mid-green leaves, to 36in (90cm) long, with dark red stripes on upper sides. In mid- to late spring, bears a spike, 2–3in (5–8cm) long, of yellow flowers and pink to purple-red upper bracts. Z12

CURIO  
ASTERACEAE/COMPOSITAE

Genus of succulent herbaceous perennials, often with creeping stems, grown for its interestingly shaped succulent leaves. Needs full sun or partial shade, well-drained soil, and good ventilation when grown under cover. Propagate by seed when ripe, or by stem cuttings in summer. **C. articulata**, syn. *Baculellum articulatum*, *Kleinia articulata*, *Senecio articulatus* (Candle plant). Deciduous, spreading, perennial succulent. H 2ft (60cm), S indefinite. Branching, gray-marked, blue stems have weak joints. Bears rounded to ovate, 3–5-lobed, gray leaves and flattish heads of small, cup-shaped, yellow flowers from spring to fall. Offsets freely from stolons. Z10–11 (min. 50°F/10°C). **‘Variegatus’** (syn. S.a. ‘Variegatus’) illus. p.486. **C. rowleyana**, syn. *Senecio rowleyanus* (String-of-beads), illus. p.481.

**Curtonus paniculatus.** See *Crocosmia paniculata*.

CYANANTHUS  
CAMPANULACEAE

Genus of late summer-flowering perennials, suitable for rock gardens, walls, and troughs. Needs partial shade and humus-rich, moist but well-drained soil. Propagate by softwood cuttings in spring or by seed in fall. **C. lobatus**. Prostrate perennial. H ¾in (2cm), S 8in (20cm). Branched stems have small, wedge-shaped, dull green leaves. In late summer, each stem bears a funnel-

shaped, blue flower. Z3–9. **f. albus** illus. p.359. **C. microphyllus** illus. p.369.

CYANOTIS  
COMMELINACEAE

Genus of evergreen, creeping perennials, grown for their foliage. Needs a position in sun or partial shade, with humus-rich, well-drained soil. Propagate by tip cuttings from spring to fall. **C. kewensis** (Teddy-bear vine). Evergreen perennial forming rosettes with trailing stems. H 2in (5cm), S 12in (30cm). Clasping the stem are 2 rows of overlapping, ovate leaves, to 2in (5cm) long, dark green above, purple with velvety, brown hairs below. Propagate by clusters of 3-petaled, purplish-pink flowers are produced in axils of leaflike bracts almost all year round. Z11–12 (min. 50°F/10°C) **C. somaliensis** (Pussy ears). Evergreen, creeping perennial. H 2in (5cm), S indefinite. Small, narrow, glossy, dark green leaves with white hairs surround stems. Has purplish-blue flowers in leaf axils in winter–spring. Z11–12 (min. 50°F/10°C)

CYATHEA  
SYN. ALSOPHILA, SPHAEROPTERIS  
CYATHEACEAE

Genus of evergreen tree ferns, grown for their foliage and overall appearance. Needs a humid atmosphere, sun or partial shade and humus-rich, moisture-retentive but well-drained soil. Water potted plants freely in summer, moderately at other times. Propagate by spores in spring. **C. australis**, syn. *Alsophila australis* (Australian tree fern), illus. p.452. **C. australis of gardens.** See *C. cooperi*. **C. cooperi**, syn. *Alsophila australis* of gardens, *C. australis* of gardens. H 6–15ft (2–5m), S 10–12ft (3–4m). Evergreen, fast-growing tree fern with a slender trunk and mid-green fronds, to 12ft (4m) long, divided into lance-shaped or linear leaflets; white-scaly frond stalks do not persist. Z8–11 **C. medullaris** (Black tree fern, Mamaku). Evergreen, upright tree fern with a slender, black trunk. H 22–50ft (7–16m), S 20–40ft (6–12m). Has arching fronds, each to 22ft (7m) long, divided into small, oblong, glossy, dark green leaflets, paler beneath. Z10–11

CYCAS  
CYCADACEAE

Genus of slow-growing, evergreen, woody-stemmed palmlike cycads. Prefers a position in full light and humus-rich, well-drained soil. In frost-prone areas, grow in a temperate or warm greenhouse or as houseplants; in warmer climates, they are excellent specimen plants. Water containerized specimens moderately, less when not in full growth. Propagate in spring by seed or suckers taken from mature plants. **C. revoluta** (Japanese sago palm) illus. p.457.