

F

FABIANA

SOLANACEAE

Genus of heathlike, evergreen shrubs, grown for their foliage and flowers. Where marginally hardy, plant in a sheltered position. Requires full sun and fertile, well-drained soil. Propagate by softwood cuttings in summer.

F. imbricata ‘Prostrata’. Evergreen, low-growing, mound-forming, very dense shrub. H 3ft (1m), S 6ft (2m). Shoots are densely covered with tiny, heathlike, deep green leaves. Bears profuse, tubular, white flowers in early summer. **f. violacea** (syn. *F.i.* ‘Violacea’) illus. p.204.

FAGUS

Beech

FAGACEAE

Genus of deciduous trees, grown for their habit, foliage and fall color. Insignificant flowers are borne in late spring and hairy fruits ripen in fall to release edible, triangular nuts. Needs sun or semishade; purple-leaved forms prefer full sun, yellow-leaved forms a little shade. Grows well in any but waterlogged soil. Trim *F. sylvatica* in summer, when used as hedging. Propagate species by seed sown in fall, selected forms by budding in late summer. Bracket fungi, canker-causing fungi, aphids and beech coccus may cause problems.

F. americana. See *F. grandifolia*.

F. grandifolia, syn. *F. americana* (American beech). Deciduous, spreading tree. H and S 30ft (10m). Ovate, silky, pale green young leaves mature to dark green in summer, then turn golden-brown in fall. Z3–9

F. orientalis (Oriental beech). Deciduous, spreading tree. H 70ft (20m), S 50ft (15m). Large, ovate, wavy-edged, dark green leaves turn yellow-brown in fall. Z4–7

F. sylvatica (European beech) illus. p.64. **‘Aspleniifolia’** see *F.s.* var. *heterophylla* ‘Aspleniifolia’. **f. atropunicea** see *F.s.* **Atropurpurea Group** (syn. *F.s.* f. *atropunicea*, *F.s.* f. *purpurea*; Copper beech) is a deciduous, round-headed tree, H 80ft (25m), S 50ft (15m), with ovate, wavy-margined, purple leaves, turning a rich copper in fall.

‘Aurea Pendula’ is a deciduous, slender tree with pendulous branches. H 30ft (10m), S 5ft (1.5m). Ovate, wavy-edged, bright yellow leaves turn rich yellow and orange-brown in fall. **‘Dawyck’** illus. p.79. **‘Dawyck Gold’** is compact and columnar, with bright yellow young foliage, turning green; H 50ft (15m), S 6ft (2m). **‘Dawyck Purple’** is similar to ‘Dawyck’, but has deep purple foliage. **var. heterophylla ‘Aspleniifolia’** (syn. *F.s.* ‘Aspleniifolia’; Fern-leaved beech) illus. p.64. **‘Pendula’** (syn. *F.s.* f. ‘Pendula’; Weeping beech) illus. p.62. **f. pendula** see *F.s.* ‘Pendula’. **‘Purple Fountain’** has upright leaders with narrow, weeping branches; H 30ft (10m), S 12ft (4m).

f. purpurea see *F.s.* **Atropurpurea Group**. **‘Purpurea Pendula’**, H and S 10ft (3m),

has stiff, weeping branches and blackish-purple foliage. **‘Riversii’** illus. p.61. **‘Rohanii’** illus. p.61. **‘Zlatia’** has yellow, young foliage, later turning mid- to dark green. Z5–7

FALLOPIA

SYN. BILDERDYKIA, REYNOUTRIA

POLYGONACEAE

Genus of rhizomatous, scrambling or climbing, woody-based perennials, good for training on pergolas and deciduous trees. Grow in full sun or partial shade and moist but well-drained soil. Propagate by seed sown as soon as ripe or in spring, or by semiripe cuttings in summer, or hardwood cuttings in fall.

F. aubertii. See *F. baldschuanica*.

F. aubertii of gardens. See

F. baldschuanica.

F. baldschuanica, syn. *F. aubertii*, *F. aubertii* of gardens, *Polygonum baldschuanicum* (Mile-a-minute plant, Russian vine), illus. p.208.

FALLUGIA

ROSACEAE

Genus of one species of deciduous shrub, grown for its flowers and showy fruit clusters. Where marginally hardy, needs protection in winter. Requires a hot, sunny position and well-drained soil. Propagate by softwood cuttings taken in summer or by seed sown in fall.

F. paradoxa (Apache plume) illus. p.128.

FARFUGIUM

ASTERACEAE/COMPOSITAE

Genus of rhizomatous, evergreen perennials, grown for their foliage and daisylike flower heads. Grow in sun or semishade and in moist but well-drained soil. Propagate by division in spring or by seed in fall or spring.

F. japonicum, syn. *Ligularia tussilaginea*. Loosely clump-forming perennial. H and S 2ft (60cm). Has large, rounded, toothed, basal, mid-green leaves, above which rise woolly, branched stems bearing clusters of daisylike, pale yellow flower heads in late summer. Z7–9. **‘Aureomaculatum’** (Leopard plant) has variegated, gold-and-white leaves.

FARGESIA

POACEAE/GRAMINEAE

See also GRASSES, BAMBOOS, RUSHES, and SEDGES.

F. muriellae, syn. *Arundinaria muriellae*, *Sinarundinaria muriellae*, *Thamnocalamus spathaceus* of gardens (Umbrella bamboo). Evergreen, clump-forming bamboo. H 12ft (4m), S indefinite. Has attractive, gray young culms with loose, light brown sheaths and broad, apple-green leaves, each very long, drawn-out at its tip. Flower spikes are unimportant. Z5–9

F. nitida, syn. *Arundinaria nitida*, *Sinarundinaria nitida* (Fountain bamboo). Evergreen, clump-forming bamboo. H 15ft (5m), S indefinite. Has small, pointed, mid-green leaves on dark purple stalks and several branches at each node. Stems are often purple with close sheaths. Z5–9.

FASCICULARIA

BROMELIACEAE

Genus of one species of evergreen, rosette-forming perennial, with no or short stems, grown for its overall appearance. Prefers full light and well-drained soil. Water moderately from spring to fall, sparingly in winter. Propagate by offsets or division in spring.

F. andina. See *F. bicolor*.

F. bicolor, syn. *F. andina*. Perennial forming congested hummocks. H to 18in (45cm), S to 2ft (60cm). Has dense rosettes of linear, tapered, arching, mid- to deep green leaves; the inner leaves are bright scarlet at flowering. In summer, produces a cluster of tubular, pale blue flowers, surrounded by white bracts, at the heart of each mature rosette. Best grown at not less than 2°C (36°F). Z10–13

X FATSHEDERA

ARALIACEAE

Hybrid genus (*Fatsia japonica* ‘Moseri’ x *Hedera helix* ‘Hibernica’) of one evergreen, fall-flowering shrub, grown for its foliage. Good trained against a wall or pillar or, if supported by canes, cultivated as a houseplant. Thrives in a sunny or shaded position and in fertile, well-drained soil. Propagate by semiripe cuttings in summer.

x F. lizei (Tree ivy) illus. p.211.

‘Annemieke’ (syn. x *F.l.* ‘Lemon and Lime’) is an evergreen, mound-forming, loose-branched shrub that may also be trained as a climber. H 4–6ft (1.2–2m), S 10ft (3m). Produces deeply lobed, glossy, deep green leaves, variegated with yellow. Sprays of small, white flowers are borne in fall. **‘Lemon and Lime’** see x *F.l.* ‘Annemieke’. **‘Variegata’**, H 5ft (1.5m), or more if trained as a climber, has leaves, narrowly edged with creamy-white. Z8–11

FATSIA

ARALIACEAE

Genus of 2 or 3 species of evergreen, fall-flowering shrubs or small trees, grown for their foliage, flowers and fruits. Excellent for conservatories. Where marginally hardy, shelter from strong winds. Tolerates sun or shade and requires fertile, well-drained soil. Propagate by semiripe cuttings taken in summer, or by seed sown in fall or spring.

F. japonica, syn. *Aralia japonica*, *A. sieboldii* (Japanese aralia, Japanese fatsia). Evergreen, rounded, dense shrub. H and S 10ft (3m). Has sturdy shoots and very large, rounded, deeply lobed, glossy, dark green leaves. Dense clusters of tiny, white flowers are produced in fall, and these are followed by rounded, black fruits. Z8–10. **‘Variegata’** illus. p.144.

F. papyrifera. See *Tetrapanax papyrifer*.

FAUCARIA

Tiger jaws

AIZOACEAE

Genus of clump-forming, almost stemless, sometimes fleshy-rooted, perennial succulents with semicylindrical or 3-angled, fleshy, bright green leaves and yellow, pink, or white flowers that open

FEROCACTUS

in late afternoons in fall. Buds and dead flowers may appear orange or red. Frost tender. Requires full sun and well-drained soil. Keep dry in winter and water sparingly in spring. Propagate by seed or stem cuttings in spring or summer.

F. tigrina illus. p.495.

Feijoa. See *Acca*.

FELICIA

SYN. AGATHAEA

Blue daisy

ASTERACEAE/COMPOSITAE

Genus of annuals, perennials, evergreen subshrubs and (rarely) shrubs, grown for their daisylike, mainly blue flower heads. Requires a position in full sun and well-drained soil. Water potted plants moderately, less when not in full growth; dislikes wet conditions, particularly in low temperatures. Cut off dead flowering stems and cut back straggly shoots regularly. Propagate by seed in spring, or by greenwood cuttings in summer or early fall.

F. amelloides, syn. *Aster amelloides*, *A. capensis*, *A. coelestis* (Blue daisy). Bushy subshrub, often grown as an annual. Bears ovate to oblong, deep green leaves, and light to deep blue flower heads from summer to fall. Z8–11. **‘Santa Anita’** illus. p.157.

F. bergeriana (Kingfisher daisy). Fairly fast-growing, mat-forming annual. H and S 6in (15cm). Produces lance-shaped, hairy, gray-green leaves and small, daisylike, blue flower heads, with yellow centers, that open only in sunshine in summer and early fall. Z11

FENESTRARIA

AIZOACEAE

Genus of one species of variable, dwarf, stemless, cushion-forming, perennial succulent. Club-shaped, fleshy leaves have gray ‘windows’ in their flattened tips. Frost tender. Requires sun and very well-drained soil. Keep bone dry in winter. Propagate by seed in spring or summer.

F. aurantiaca. See *F. rhopalophylla* subsp. *aurantiaca*.

F. rhopalophylla. Clump-forming, perennial succulent. H 2in (5cm), S 8in (20cm). Forms open cushions of erect, club-shaped, glossy, glaucous to mid-green leaves, each with a flattened tip. Bears daisylike, white flowers on long stems in late summer and fall. Z13–15. (min. 41°F/5°C). **subsp. aurantiaca** (syn. *F. aurantiaca*; Baby’s toes) illus. p.495.

FEROCACTUS

CACTACEAE

Genus of slow-growing, flattened, spherical, perennial cacti, becoming columnar after many years. Frost tender. Needs full sun and very well-drained soil. Propagate by seed in spring or summer. Treat blackened areoles with systemic fungicide and ensure plants have good ventilation.

F. acanthodes. See *F. cylindraceus*.

F. chrysacanthus. Slow-growing,