

I. ‘Making Eyes’. Rhizomatous, bearded iris (Standard Dwarf). H 12–14in (30–35cm). In early spring, produces neat flowers with pale lemon-white standards, narrow, white-margined, dark purple-violet falls and yellowish-white beards. Z3–8

I. ‘Mary McIlroy’. Rhizomatous, bearded iris (Intermediate). H 16–20in (40–50cm). In early to midspring, bears bright yellow blooms, the standards veined slightly deeper yellow, with darker veins on the falls and lemon-yellow beards. Z3–8

I. ‘Matinata’. Rhizomatous, bearded iris (Tall). H 3ft (1m). In early summer, well-branched stem produces 6–9, occasionally to 12 flowers, 6in (15cm) across, that are dark purple-blue throughout. Z3–9

I. ‘Maui Moonlight’. Rhizomatous, bearded iris (Intermediate). H 26in (65cm). Has fragrant, ruffled, lemon yellow flowers in midspring. Falls have slightly darker veins and yellow beards. Z3–9

I. ‘Mer du Sud’. Rhizomatous, bearded iris (Tall). H 32in (80cm). In early summer, produces flowers with sea blue standards and falls, and light blue beards. Z3–9

I. missouriensis (Missouri flag). Very variable, rhizomatous, beardless Pacific Coast iris. H to 30in (75cm). In late spring or early summer, branched stem produces 2 or 3 pale blue, lavender, lilac, blue or white flowers, 2–3in (5–8cm) across, in each spathe. Falls are veined and usually have a yellow blaze. Z3–9

I. ‘Mountain Lake’ (illus. p.224). Rhizomatous, beardless Siberian iris. H 3ft (1m). From late spring to early summer, branched stem produces 1–4 mid-blue flowers, 2–4in (5–10cm) across, with darker veining on falls. Prefers moist soil. Z3–8

I. ‘Natascha’ illus. p.415.

I. ochroleuca. See *I. orientalis*.

I. ‘Ola Kala’ (illus. p.225). Robust, rhizomatous, bearded iris (Tall). H 3ft (1m). Produces neat, scented, rich deep yellow flowers; the falls have darker yellow bases and yellow beards. Z3–9

I. orientalis, syn. *I. ochroleuca* (illus. p.224). Rhizomatous, beardless Spuria iris. H to 3ft (90cm). In late spring, each stem, usually with one branch, bears 3–5 white flowers, 3–4in (8–10cm) across. Falls are white with yellow centers. Leaves are often present over winter. Z6–9

I. orientalis of gardens. See *I. sanguinea*.

I. ‘Orinoco Flow’. Rhizomatous, bearded iris (Intermediate). H 25in (63cm). In mid- and late spring, produces strongly ruffled white flowers, with margins heavily marked in deep purple, and navy-blue beards. Z3–9

I. pallida (Dalmatian iris). Rhizomatous, bearded iris. H 28–36in (70–90cm) or more. In late spring and early summer, produces 2–6 scented, lilac-blue flowers, 3–5in (8–12cm) across and with yellow beards, from silvery spathes on strong, branched stems. Z4–9. ‘**Aurea Variegata**’ see *I.p.* ‘**Variegata**’. ‘**Variegata**’ (syn. *I.p.* ‘**Aurea Variegata**’) has striped green and yellow leaves.

I. ‘Papillon’. Clump-forming, rhizomatous, beardless Siberian iris. H 3ft (90cm). In late spring and early summer, bears flowers with mid-blue standards and pale-blue styles. Falls are mid-blue, with a yellowish-brown throat, marked by bold, blue veins. Z3–8

I. ‘Perry’s Blue’ (illus. p.225). Robust, rhizomatous, beardless Siberian iris. H 3ft (1m). In late spring and early summer, produces neat, pale purplish-blue flowers, with noticeably deeper blue veins. Slightly twisted standards and broad, rounded falls are white margined and creamy-white near the bases, and have dark yellow markings in the throats. Z3–8

I. ‘Pixie’ illus. p.418.

I. ‘Pogo’. Rhizomatous, bearded iris (Standard Dwarf). H 12in (30cm). Bright yellow flowers are borne in early spring, Each fall has a large reddish-brown blotch, and a yellow beard. Z3–9

I. ‘Professor Blaauw’. Bulbous Xiphium iris (Dutch). H 32in (80cm), 5 6in (15cm). From spring to early summer, produces 1 or 2 rich violet-blue flowers, 2½–3in (6–8cm) across. Narrowly lance-shaped, channeled, mid-green leaves are scattered up flower stem. Z3–9

I. pseudacorus (Yellow flag; illus. p.225). Robust, rhizomatous, beardless iris. H to 6ft (2m). From early to midsummer, branched stem produces 4–12 golden-yellow flowers, 2–5in (5–12cm) across, usually with brown or violet veining and a darker yellow patch on the falls. Leaves are broad, ridged and grayish-green. Prefers semishade and thrives in a water garden. Z5–8. **var. bastardii** illus. p.445. ‘**Crème de la Crème**’ has creamy white flowers with purple markings. ‘**Roy Davidson**’ has brown-veined yellow flowers with brown patches on the falls. ‘**Variegata**’ has yellow-and-green-striped foliage in spring, often turning green before flowering.

I. pumila (Dwarf bearded iris). Rhizomatous, bearded iris. H 4–6in (10–15cm). In midspring, produces a ½in (1cm) long flower stem bearing 2 or 3 long-tubed flowers, 1–2in (2.5–5cm) across, varying from violet-purple to white, yellow or blue, with yellow or blue beards on the falls. Prefers very well-drained, slightly alkaline soil. Z4–9

I. ‘Rajah’. Rhizomatous, bearded iris (Tall). H 3ft (90cm). A striking iris, bearing flowers with yellow standards and reddish-brown falls marked with yellow stripes in early summer. The falls have a yellow beard. Z3–9

I. ‘Raspberry Blush’. Rhizomatous bearded (Intermediate). H 2ft (60cm). Produces bi-tone flowers in spring and early summer. Standards are slightly ruffled and raspberry-pink with darker veins. The falls have a dark raspberry center and paler raspberry-pink margins, with a dark orange beard. Z3–9

I. reticulata. Bulbous Reticulata iris. H 4–6in (10–15cm), 5 1½–2½in (4–6cm). In early spring, produces a solitary, fragrant, long-tubed, deep violet-purple flower, 1½–2½in (4–6cm) across, with a yellow ridge down each fall center. Narrow, squared leaves elongate after flowering time. Z5–8. ‘**Cantab**’ (illus. p.225) has clear pale blue flowers with a deep yellow ridge on each fall. ‘**Clairette**’ see *I. ‘Clairette*’. Flowers of ‘**J.S. Dijt**’ are reddish-purple with an orange ridge on each fall. ‘**Violet Beauty**’ see *I. ‘Violet Beauty*’.

I. ‘Ringo’ (illus. p.224). Vigorous, rhizomatous, bearded iris (Tall). H 3ft

(90cm). In late spring and early summer, produces lightly ruffled flowers with white standards touched purple on the midribs and dark reddish-purple falls with narrow, white margins and orange beards. Z5–8

I. x robusta ‘Gerald Darby’ (illus. p.224). Rhizomatous, beardless Species Hybrid iris. H 32in (80cm). In late spring and early summer, unbranched purplish-green stems produce up to 4 violet-blue flowers, with deep yellow signals on the falls. Leaves are stained purple at the bases. Prefers wet conditions. Z4–9

I. rosenbachiana. Bulbous Juno iris. H 4–6in (10–15cm), 5 2½in (6cm). In spring, produces 1 or 2 long-tubed flowers, 1½–2in (4–5cm) across, with small, downward-turned, rich purple standards and reddish-purple falls, each with a yellow ridge in the center. Has lance-shaped, channeled, mid-green leaves in a basal tuft. Difficult to grow; best grown in an alpine house or cold frame. Z4–9

I. ‘Ruffled Velvet’. Rhizomatous, beardless Siberian iris. H 22in (55cm). In early summer, produces 2 or 3 red-purple flowers marked with yellow. Z4–9

I. ‘Sable’. Rhizomatous, bearded iris (Tall). H 3ft (90cm). Has grayish-green foliage. In early summer, produces flowers with large, dark violet blue standards and almost black falls, with white markings near the throat. Z3–9

I. ‘Salamander Crossing’. Rhizomatous, beardless Siberian iris. H 3½ft (1.1m). Flowers from early to late spring. Pale yellow falls are speckled with lavender-blue, strongly marked with lavender-blue veins, and are a deeper yellow toward the throats. Pale lavender-blue standards have darker lavender-blue speckles. Z3–9

I. sanguinea, syn. *I. extremorientalis*, *I. orientalis* of gardens. Rhizomatous, beardless Siberian iris. H to 3ft (90cm). From late spring to early summer, each branched stem produces 2 or 3 deep purple or red-purple flowers, 2–4in (5–10cm) across, from each set of bracts. Falls are red-purple with white throats finely veined purple. Z3–9. ‘**Snow Queen**’ (illus. p.224) has pure white flowers with yellow-green marks at the bases of the falls.

I. setosa illus. p.441.

I. ‘Shepherd’s Delight’. Rhizomatous, bearded iris (Tall). H 3ft (1m). In early summer, well-branched stem produces 6–10 clear pink flowers, 6–7in (15–18cm) across, with a yellow cast. Z3–8

I. sibirica (Siberian flag; illus. p.441). Rhizomatous, beardless Siberian iris. H 20–48in (50–120cm). From late spring to early summer, branched stem produces 2 or 3 dark-veined, blue or blue-purple flowers, 2–4in (5–10cm) across, from each spathe. Prefers moist or boggy conditions. Z3–8

I. ‘Silvery Beauty’. Bulbous iris (Dutch). H 28in (70cm). Bears violet blue flowers in spring and early summer. The falls are white, with a bright yellow center. Best planted in drifts. Z5–9

I. sintenisii. Rhizomatous, beardless Spuria iris. H 12in (30cm). Has linear, dark green leaves. In late spring, bears 2 white flowers, densely veined blue-purple. Z6–8

I. sisyrinchium. See *Moraea sisyrinchium*.

I. ‘Soft Blue’ (illus. p.225). Robust,

rhizomatous, beardless Siberian iris. H 30in (75cm). In early to midspring, bears pale blue flowers; the long, arching falls have yellow basal markings and darker blue veins. Z3–9

I. spuria. Very variable, rhizomatous, beardless Spuria iris. H 20–36in (50–90cm). Strong, erect-branched stem bears 2–5 pale blue-purple, sky-blue, violet-blue, white or yellow flowers, 2–5in (5–12cm) across, in early summer. Prefers moist soil. Z6–9

I. ‘Stepping Out’. Rhizomatous, bearded iris (Tall). H 3ft (1m). Well-branched stem bears 8–11 white flowers, 5½–6in (14–15cm) across, with deep blue-purple marks in early summer. Z3–9

I. stylosa. See *I. unguicularis*.

I. ‘Superstition’. Rhizomatous, bearded iris (Tall). H 3ft (90cm). Produces striking, dark purple-brown flowers, with blue-black beards, in early summer. Z3–9

I. ‘Supreme Sultan’. Vigorous, rhizomatous, bearded iris (Tall). H 3ft (1m). In late spring and early summer, bears large, ruffled flowers, with deep golden-yellow standards, rich dark red-brown falls that are paler at the margins, and deep yellow beards. Z3–8

I. susiana (Mourning iris). Rhizomatous Oncocyclis iris. H 14–16in (35–40cm). In late spring, produces a solitary, grayish-white flower, 3–6in (8–15cm) across, heavily veined deep purple. Standards appear larger than incurved falls, which each bear a black blaze and a deep purple beard. Best in a cold frame or alpine house. Z7–9

I. ‘Sweet Musette’. Vigorous, rhizomatous, bearded iris (Tall). H 36–38in (90–95cm). In mid- to late spring, produces large, ruffled, frilly flowers, with lavender-flushed, peach-pink standards, purplish-pink falls and orange beards. Z3–8

I. tectorum (Japanese roof iris, Wall flag). Evansia iris with sturdy rhizomes. H 10–14in (25–35cm). Has fans of broadly lance-shaped, ribbed leaves. In early summer, sparsely branched stem produces 2–3 darker-veined, bright lilac flowers, ½–3in (1–8cm) across with a white crest on each fall, from each spathe. Prefers a sheltered, sunny site near a south- or west-facing wall. Z5–9

I. tenax. Rhizomatous, beardless Pacific Coast iris. H 6–12in (15–30cm). From late spring to early summer, produces 1 or 2 deep purple to lavender-blue flowers, 3–5in (8–12cm) across, often with yellow-and-white marking on falls. White, cream and yellow variants also occur. Narrow, dark green leaves are stained pink at base. Z5–9

I. ‘Thornbird’. Rhizomatous, bearded iris (Tall). H 3ft (90cm). In early summer, bears up to 7 flowers, with pale greenish-white standards and greenish-brown falls overlaid with deep violet lines. Long, horned beard is violet, tipped with mustard-yellow. Z3–9

I. ‘Titan’s Glory’. Robust, rhizomatous, well-branched, bearded iris (Tall). H to 3ft (1m). Produces very large, deep purple-blue flowers, with an almost silken texture, in midspring. Z3–8

I. ‘Tropic Night’ (illus. p.225). Strong-growing, rhizomatous, beardless Siberian iris. H 3ft (90cm). In late spring and early summer, produces deep violet-blue flowers; upright standards and rounded falls have strong, white feathering and