

flowers are borne above small clusters of strap-shaped, dark green leaves. Z5–9
L. rediviva (Bitter root) [pink form] illus. p.365; [white form] illus. p.360.

LEYCESTERIA

CAPRIFOLIACEAE

Genus of suckering, deciduous shrubs, grown for their showy flower clusters. Fully hardy to half hardy. Needs full sun and fertile, well-drained soil. Propagate by softwood cuttings in summer or by seed or division in fall.
L. formosa (Himalayan honeysuckle, Pheasant berry). Deciduous, upright shrub. H and 5 6ft (2m). Has blue-green shoots and slender, ovate, dark green leaves. In summer and early fall, small, funnel-shaped, white flowers open at tip of each pendent cluster of purplish-red bracts, followed by spherical, reddish-purple fruits. Cut weak shoots to ground level in early spring. Z7–9. **GOLDEN LANTERNS** (**Notbruce**) has soft yellow leaves.

LEYMUS

POACEAE/GRAMINEAE

See also GRASSES, BAMBOOS, RUSHES, and SEDGES.
L. arenarius, syn. *Elymus arenarius* (Lyme grass). Vigorous, spreading, herbaceous, rhizomatous, perennial grass. H to 5ft (1.5m), 5 indefinite. Has broad, glaucous leaves and stout, terminal spikes of grayish-green flowers on erect stems in late summer. Useful for binding coastal dunes. Z6–9

LIATRIS

Gay feathers

ASTERACEAE/COMPOSITAE

Genus of summer-flowering perennials with thickened, cormlike rootstocks. Fully hardy. Prefers sun and well-drained soil. Propagate by division in spring.
L. callilepis of gardens. See *L. spicata*.
L. pycnostachya (Kansas gay feather). Clump-forming perennial. H 4ft (1.2m), 5 1ft (30cm). In summer, bears tall spikes of clustered, feathery, mauve-pink flower heads. Grasslike, dark green leaves form basal tufts.
L. spicata, syn. *L. callilepis* of gardens, illus. p.438.

LIBERTIA

IRIDACEAE

Genus of rhizomatous perennials, grown for their foliage, decorative seed pods, and flowers. Borderline fully hardy to frost hardy. Needs a sheltered, sunny, or partially shaded site and well-drained soil. Propagate new plants by division in spring, or by seed in fall or spring.
L. grandiflora (New Zealand satin flower) illus. p.230.
L. ixioides. Clump-forming, rhizomatous perennial. H and 5 2ft (60cm). Produces panicles of saucer-shaped, white flowers in summer. Grasslike, dark green leaves turn orange-brown during winter. Z9–11.
‘Goldfinger’ illus. p.277.

Libocedrus chilensis. See *Austrocedrus chilensis*.

Libocedrus decurrens. See *Calocedrus decurrens*.
Libonia floribunda. See *Justicia floribunda*.

LIGULARIA

ASTERACEAE/COMPOSITAE

Genus of perennials, grown for their foliage and large, daisylike flower heads. Fully hardy. Grow in sun or partial shade and in moist but well-drained soil. Propagate by division in spring or by seed in fall or spring. Prone to slug and snail attack.
L. clivorum **‘Desdemona’**. See *L. dentata* ‘Desdemona’.
L. ‘Britt Marie Crawford’ illus. p.445.
L. dentata **‘Desdemona’** (syn. *L. clivorum* ‘Desdemona’, *Senecio clivorum* ‘Desdemona’) is a compact, clump-forming perennial. H 4ft (1.2m), 5 2ft (60cm). Heart-shaped, long-stalked, leathery, basal, dark brownish-green leaves, are almost mahogany beneath. In mid- to late summer, bears terminal clusters of large, daisylike, vivid orange flower heads on branching stems. Z4–8
L. ‘Gregynog Gold’. Clump-forming perennial. H to 6ft (1.8m), 5 3ft (1m). Leaves are large, heart-shaped, and deep green. Conical panicles of daisylike, orange-yellow flower heads are borne from mid- to late summer. Z4–8
L. przewalskii, syn. *Senecio przewalskii*, illus. p.445.
L. stenocephala. Loosely clump-forming perennial. H 5ft (1.5m), 5 3ft (1m). Has jagged-edged, round, mid-green leaves. Large heads of daisylike, yellow-orange flowers open on purplish stems from mid- to late summer. Z4–8
L. ‘The Rocket’ illus. p.219.
L. tussilaginea. See *Fartugium japonicum*.

LIGUSTRUM

Privet

OLEACEAE

Genus of deciduous, semievergreen or evergreen shrubs and trees, grown for their foliage and, in some species, flowers. Fully to frost hardy. Requires sun or partial shade, the variegated forms doing best in full sun. Thrives on any well-drained soil, including chalky soil. All except *L. lucidum* occasionally need cutting back in mid-spring to restrict growth. Propagate by semiripe cuttings in summer. ☉ All parts may cause severe discomfort if ingested.
L. japonicum (Japanese privet). Evergreen, bushy, dense shrub. H 10ft (3m), 5 8ft (2.5m). Has ovate, glossy, very dark green leaves and, from midsummer to early fall, large, conical panicles of small, tubular, white flowers with lobes. Z7–10.
‘Coriaceum’ see *L.j.* ‘Rotundifolium’.
‘Rotundifolium’, syn. *L.j.* ‘Coriaceum’, is slow-growing, and produces a dense mass of rounded, leathery leaves.
L. lucidum (Chinese privet). Evergreen, upright shrub or tree. H 30ft (10m), 5 25ft (8m). Bears large, ovate, glossy, dark green leaves. Produces large panicles of small, tubular, white flowers, with 4 lobes, in late summer and early fall. Z7–10.
‘Excelsum Superbum’ has bright green leaves, marked with pale green and yellow-edged, and small, tubular, white

flowers. **‘Tricolor’** has narrow, green and gray-green leaves with creamy white margins, pink-tinged when young.
L. ovateifolium illus. p.119. **‘Aureum’** is a vigorous, evergreen or semievergreen, upright, dense shrub. H 12ft (4m), 5 10ft (3m). Leaves are ovate, glossy, and mid-green, broadly edged with bright yellow. Dense panicles of small, rather unpleasantly scented, tubular, white flowers, with 4 lobes, are borne in mid-summer and are followed by spherical, black fruits. Cut back hedges to 1ft (30cm) after planting and prune hard for first 2 years; then trim as necessary during the growing season. **‘Vicaryi’**, syn. *L. ‘Vicaryi’*, illus. p.140. Z5–9
L. sinense illus. p.112.
L. ‘Vicaryi’. See *L. ovateifolium* ‘Vicaryi’.
L. vulgare. Deciduous or semievergreen, bushy shrub. H and 5 10ft (3m). Leaves are narrowly lance-shaped and dark green. Bears panicles of small, strongly scented, tubular, white flowers, with 4 lobes, from early to midsummer; then spherical, black fruits. Cut back hedges to 1ft (30cm) after planting and prune hard for the first 2 years; then trim as necessary during the growing season. Z4–8

LILIUM

Lily

LILIACEAE

Genus of mainly summer-flowering bulbs, grown for their often fragrant, brightly colored flowers. Each fleshy-scaled bulb produces one unbranched, leafy stem; in some cases roots develop on the stem just above the bulb. Glossy, mid- to dark green leaves are mostly lance-shaped or linear, to 9in (22cm) long, and arranged in whorls or spirals, or are scattered alternately up the stems. Flowers, usually several per stem, are mainly trumpet- to bowl-shaped or with the 6 petals strongly reflexed to form a turkscap shape. (Petals of *Lilium* are known botanically as perianth segments.)
Three categories of flower size—small, medium, and large—are used in the descriptions below. For turkscap, bowl-, cup- and star-shaped flowers: small is up to 2in (5cm) across; medium is 2–3in (5–7cm) across; large is over 3in (7cm) across. For trumpet- and funnel-shaped flowers: small is up to 3in (7cm) long; medium is 3–4in (7–10cm) long; large is over 4in (10cm) long. Most lilies vary between 3ft (1m) and 6ft (2m) in height, although a few may reach 10ft (3m).
They do not spread widely—mostly to 1ft (30cm)—therefore only height measurements are given below. Fully hardy to half hardy. Need sun and well-drained soil, unless otherwise stated. Propagate by seed in fall or spring, by bulb scales in summer, or by stem bulbils (where present) in fall. Virus, fungal diseases, and lily beetle may cause problems. Lilies are classified into 9 divisions. They may also be further classified into subdivisions according to their flower aspect: **(a)** for upward-facing flowers; **(b)** outward-facing; **(c)** downward-facing. Also, they are classified according to flower form: **(a)** trumpet-shaped; **(b)** bowl-shaped; **(c)** predominantly flat (albeit sometimes

with recurved petal tips); **(d)** turkscap. For example, ‘African Queen’ Vlb-c/a is a Trumpet/Aurelian hybrid with flowers that are outward- to downward-facing and trumpet-shaped. See also feature panel pp.388–391.

Division I (Asiatic hybrids)

These lilies are derived from various Asiatic species, including *L. bulbiferum*, *L. cernuum*, *L. concolor*, *L. davidii*, *L. lancifolium* and *L. maculatum*. The flowers are borne in racemes or umbels, and are usually unscented. The leaves are narrowly ovate and arranged alternatively.

Division II (Martagon hybrids)

Derived primarily from *L. hansonii* and *L. martagon*, these lilies produce racemes of turkscap, sometimes scented flowers, and have whorls of elliptic leaves.

Division III (Euro-Caucasian hybrids)

Derived from *L. candidum* and other European species, except *L. martagon*, these lilies produce sometimes scented, mostly turkscap flowers, singly or in umbels or racemes. Leaves are elliptic, and spirally arranged or scattered.

Division IV (American hybrids)

Derived from American species, these lilies bear racemes of sometimes scented, mostly turkscap, but occasionally funnel-shaped flowers, and have whorls of lance-shaped to elliptic leaves.

Division V (Longiflorum hybrids)

Derived from *L. formosanum* and *L. longiflorum*, these lilies bear racemes or umbels of large, often sweetly scented, trumpet- or funnel-shaped flowers, in some only 2 or 3 per stem. The scattered leaves are linear to narrowly lance-shaped.

Division VI (Trumpet and Aurelian hybrids)

Derived from Asiatic species, including *L. regale*, *L. henryi* and *L. sargentiae*, these lilies bear racemes or umbels of usually scented flowers. Leaves are elliptic to linear, and alternate or spirally arranged.

Division VII (Oriental hybrids)

Derived from E. Asian species, such as *L. auratum*, *L. japonicum* and *L. speciosum*, as well as their hybrids with *L. henryi*, these lilies have flowers borne in racemes or panicles, and are often scented. Leaves are lance-shaped and alternate.

Division VIII (Other hybrids)

These include the many Interdivisional hybrids which have become increasingly popular in recent years, such as Longiflorum/Asiatic hybrids (LA) and Orienpet (Oriental/Trumpet) hybrids (OT).

Division IX All true species or their selected forms.

L. ‘African Queen’ (illus. p.391). Mid- to late summer-flowering, vigorous Division Vlb-c/a lily. H 5–6ft (1.5–2m). Bears 3 large fragrant flowers, brownish-purple outside and apricot-orange on the inside, with recurved petals. Z4–8
L. ‘Altari’ (illus. p.388). Mid- to late summer-flowering, Division VIIIa-b/b (OT) lily. H 3–4ft (1–1.2m). Bears highly scented, white flowers with cranberry-red inner throats. Z4–8
L. amabile. Mid-summer-flowering Division IXc/d lily with stem roots. H 1–3ft (30–90cm). Scattered leaves are lance-shaped. Bears up to 10 unpleasant-