

when in growth, moderately at other times. Tip prune young plants to promote branching; cut back hard after flowering. Raise softwood cuttings annually in late spring. Red spider mite may cause problems. **R. indica**, syn. *R. trigyna* (Yellow flax), illus. p.459. **R. trigyna**. See *R. indica*.

RESEDA
Mignonette
RESEDACEAE

Genus of annuals and biennials with flowers that attract bees and are suitable for cutting. Fully hardy. Grow in a sunny position and in any fertile, well-drained soil. Deadhead regularly to prolong flowering. Propagate by sowing seed outdoors in spring or early fall. **R. odorata** (Mignonette) illus. p.300.

RETAMA
PAPILIONACEAE/LEGUMINOSAE

Genus of deciduous shrubs grown for their willowy, dark green or silky gray stems and pealike, white or yellow flowers. Frost hardy. Needs full sun, sharply drained soil, and a sheltered site against a south- or west-facing wall. Propagate from seed in a cold frame or under glass or by semiripe cuttings in summer. **R. monosperma**, syn. *Genista monosperma*. Deciduous, almost leafless, graceful, arching shrub. H to 12ft (4m), S 5ft (1.5m). Slender, silky-gray shoots bear clusters of small, very fragrant, white flowers in early spring. Has a few linear leaves, which soon fall. Grow against a south- or west-facing wall. Z9–11

Reynoutria. See *Fallopia*.

RHAMNUS
Buckthorn
RHAMNACEAE

Genus of deciduous or evergreen, usually thorny, shrubs and trees, with inconspicuous flowers, grown mainly for their foliage and fruits. Fully to frost hardy. Requires sun or partial shade and fertile soil. Propagate deciduous species by seed in fall, evergreen species by semiripe cuttings in summer. ☉ All parts may cause severe discomfort if ingested. **R. alaternus** (Italian buckthorn). ‘**Argenteovariegata**’ is an evergreen, bushy shrub. H and S 10ft (3m). Has ovate, leathery, glossy, gray-green leaves margined creamy-white. Tiny, yellowish-green flowers are produced from early to midsummer and followed by spherical red, then black fruits. H5

R. imeretina. Deciduous, spreading, open shrub. H 10ft (3m), S 15ft (5m). Stout shoots bear large, broadly oblong, prominently veined, dark green leaves that turn bronze-purple in fall. Small, green flowers are produced in summer. Z6–8

RHAPHIOLEPIS
ROSACEAE

Genus of evergreen shrubs and trees, grown for their foliage and fragrant, apple-blossomlike flowers. Frost hardy to half hardy. In most areas does best against a

sunny, sheltered wall; *R. umbellata* is the most hardy. Needs sun and fertile, well-drained soil. Propagate by semiripe cuttings in late summer. **R. x delacourii ‘Coates’ Crimson’**. Evergreen, rounded shrub. H 6ft (2m), S 8ft (2.5m). Clusters of fragrant, star-shaped, deep pink flowers, produced in spring or summer, are set off by the ovate, leathery, dark green leaves. Z8–10

R. indica (Indian hawthorn). Evergreen, bushy shrub. H 5ft (1.5m), S 6ft (2m). Produces clusters of fragrant, star-shaped, white flowers, flushed with pink, in spring or early summer amid narrowly lance-shaped, glossy, dark green leaves. Z9–11

R. japonica. See *R. umbellata*. **R. ovata**. See *R. umbellata*. **R. umbellata**, syn. *R. japonica*, *R. ovata*, illus. p.150.

RHAPIS
ARECACEAE/PALMAE

Genus of evergreen fan palms, grown for their foliage and overall appearance. May have tiny, yellow flowers in summer. Frost tender. In frost-prone areas, grow in a temperate or warm greenhouse, or as houseplants. In frost-free areas, grow use in a shady border. Needs partial shade and humus-rich, well-drained soil. Water containerized specimens freely when growing, moderately at other times. Propagate by seed, suckers, or division in spring. Susceptible to red spider mite. **R. excelsa**, syn. *R. flabelliformis* (Bamboo palm, Slender lady palm), illus. p.458. **R. flabelliformis**. See *R. excelsa*.

Rhazya orientalis. See *Amsonia orientalis*.

RHEUM
Rhubarb
POLYGONACEAE

Genus of rhizomatous, often tough or woody, perennials, grown for their foliage and overall appearance. Includes the edible rhubarb and various ornamental plants. Some species are extremely large and require plenty of space. Fully hardy. Prefers sun or partial shade and deep, rich, well-drained soil. Propagate by division in spring or by seed in fall ☉ Leaves may cause severe discomfort if ingested. **R. ‘Ace of Hearts’**, syn. *R. ‘Ace of Spades’*. Clump-forming perennial. H to 4ft (1.2m), S 3ft (90cm). Has elongated, heart-shaped, dark green leaves, to 14in (35cm) long, red-veined above, purple-veined beneath. Bears long panicles of numerous, tiny, star-shaped, very pale pink to white flowers in mid- to late summer. Z4–9

R. ‘Ace of Spades’. See *R. ‘Ace of Hearts’*. **R. nobile**. Clump-forming perennial. H to 6ft (2m), S 2ft (60cm). Leaves are oblong to ovate, leathery, basal, mid-green, 2ft (60cm) long. In late summer, produces long stems and conical spikes of large, overlapping, pale cream bracts that hide insignificant green flowers. Z6–9

R. palmatum. Clump-forming perennial. H to 8ft (2.5m), S to 6ft (1.8m). Has 2–2½ft (60–75cm) long, rounded, 5-lobed, mid-green leaves. In early summer, bears broad

panicles of small, creamy-green to deep red flowers. Z5–9. ‘**Atrosanguineum**’ illus. p.439. ‘**Bowles Crimson**’ has darker red flowers, and leaves that are crimson beneath. Leaves of ‘**Hadspen Crimson**’ emerge crimson, then turn green in summer; bears carmine-red flowers.

Rhipsalidopsis gaertneri. See *Hatiora gaertneri*. **Rhipsalidopsis rosea**. See *Hatiora rosea*.

RHIPSALIS
Mistletoe cactus
CACTACEAE

Genus of mostly epiphytic, perennial cacti with usually pendent, variously formed stems. Small, diurnal flowers are followed by spherical, translucent berries. Frost tender. Needs partial shade and rich, well-drained soil. Prefers 80% relative humidity—higher than for most cacti. Give only occasional, very light watering in winter. Propagate by seed or stem cuttings in spring or summer. **R. capilliformis**. See *R. teres*. **R. cereuscula** (Coral cactus) illus. p.483. **R. clavata**. See *R. gaertneri*. **R. crispata**. Bushy, then pendent, perennial cactus. H 2ft (60cm), S indefinite. Has leaflike, elliptic to oblong, pale green stem segments, to 5in (12cm) long, with undulating edges that bear short, funnel-shaped, cream or pale yellow flowers, to ½in (1cm) across, with recurved tips, in winter–spring. Flowers are followed by white, sometimes red-flushed, berries. Z11–12 (min. 45–54°F/7–12°C)

R. floccosa var. tucumanensis illus. p.493.

R. gaertneri, syn. *R. clavata*, *Hatiora clavata*. Pendent, perennial, epiphytic cactus. H 2ft (60cm), S 3ft (1m). Has multi-branched, cylindrical, dark green stems that widen toward tips. Profuse, terminal, bell-shaped, white flowers, ¾in (1.5cm) across, are produced in late winter and early spring on plants over 1ft (30cm) high. Z11–12 (min. 45–54°F/7–12°C)

R. paradoxa (Chain cactus). Bushy, then pendent, perennial cactus. H 3ft (1m), S indefinite. Triangular, green stems have segments alternately set at different angles. Short, funnel-shaped, white flowers, ¾in (2cm) across, with recurved tips, in winter–spring are followed by red berries. Z11–12 (min. 45–54°F/7–12°C)

R. salicornioides. See *Hatiora salicornioides*.

R. teres, syn. *R. capilliformis*. Pendent, perennial cactus. H 16in (40cm) or more, S 1ft (30cm). Has freely branching, cylindrical, pale green stems. In winter–spring, produces short, funnel-shaped, white flowers, to ½in (1cm) across, with recurved tips, followed by white berries. Z11–12 (min. 45–54°F/7–12°C)

R. warmingiana. See *Lepismium warmingianum*.

RHODANTHE
SYN. ACROCLINIUM
Strawflower
ASTERACEAE/COMPOSITAE

Genus of drought-tolerant subshrubs, annuals, and perennials grown for their

RHODOCHITON

daisylike, papery flower heads, which are good for cutting and drying. Half hardy. Grow in sun and in poor, very well-drained soil. Propagate by seed sown outdoors in mid-spring. Aphids may cause problems. **R. chlorocephala subsp. rosea**, syn. *Acroclinium roseum*, *Helipterum roseum*, illus. p.303. **R. manglesii**, syn. *Helipterum manglesii*. Moderately fast-growing, erect annual. H 2ft (60cm), S 6in (15cm). Has pointed-ovate, grayish-green leaves and bears daisylike, papery, red, pink, or white flower heads, in summer and early fall. Z10–11

RHODANTHEMUM
ASTERACEAE/COMPOSITAE

Genus of mat-forming, often rhizomatous perennials and subshrubs, grown for their solitary, large, daisylike, white flower heads, surrounded by prominent, usually green bracts. Most are hardy to 14°F (–10°C) in well-drained soil. Needs full sun and moderately fertile, very well-drained soil. Propagate by seed in spring or by softwood cuttings in summer. **R. hosmariense**, syn. *Chrysanthemum hosmariense*, *Pyrethropsis hosmariense*, illus. p.332.

RHODIOLA
CRASSULACEAE

Genus of perennials, some dioecious, with thick, fleshy rhizomes producing scaly, brown basal leaves and stiffly erect stems that bear triangular-ovate to lance-shaped, fleshy, gray-green leaves. Star-shaped flowers have prominent stamens, and may be unisexual or bisexual. Good for a rock garden or the front of a border. Fully hardy. Grow in full sun and moderately fertile soil. Propagate by seed in spring or fall, divide rhizomes in spring or early summer or take leaf cuttings in summer.

R. heterodonta, syn. *Sedum heterodontum*, *S. rosea* var. *heterodontum*, illus. p.268. **R. rosea**, syn. *Sedum rosea* (Roseroot). Clump-forming perennial. H 2–12in (5–30cm), S 8in (20cm). Stems are clothed with ovate to inversely lance-shaped, toothed, fleshy, glaucous leaves. In late spring or early summer, pink buds on dense, terminal heads open to small, star-shaped, greenish-, yellowish- or purplish-white flowers. Z4–8

Rhodocactus grandifolius. See *Pereskia grandifolia*.

RHODOCHITON
PLANTAGINACEAE

Genus of deciduous, perennial climbers, grown for their unusual flowers, which have pendent, long-tubed corollas, with 5 rounded segments, and inflated calyces. Best when grown as an annual. May be planted against fences and trellises or used as groundcover. Frost tender. Grow in sun and in any well-drained soil. Propagate by seed in early spring. **R. atrosanguineus**, syn. *R. volubilis*, illus. p.203. **R. volubilis**. See *R. atrosanguineus*.