

S. x boydii ‘Hindhead Seedling’ illus. p.356; cultivation group 2.

S. ‘Brookside’. See *S. burseriana* ‘Brookside’.

S. brunoniana. See *S. brunonis*.

S. brunonis, syn. *S. brunoniana*. Semi-evergreen, rosetted perennial. H 4in (10cm), 5 8in (20cm). Small, soft green rosettes of lance-shaped, rigid leaves produce many long, threadlike, red runners. Many of the rosettes die down to large terminal buds in winter. Short racemes of 5-petaled, spreading, pale yellow flowers are borne in late spring and summer on 2–3in (5–8cm) stems. Cultivation group 1. Z8–10

S. burseriana illus. p.348. **‘Brookside’** (syn. *S. ‘Brookside’*) is a slow-growing, evergreen perennial forming a hard cushion. H 1–2in (2.5–5cm), S to 4in (10cm). Has broadly linear, spiky, gray-green leaves. In early spring, bears upturned, rounded, shallowly cup-shaped, bright yellow flowers on short, red stems. Flowers of **‘Crenata’** (syn. *S. ‘Crenata’*) have fringed, white petals and red sepals. **‘Gloria’** (syn. *S. ‘Gloria’*) has dark reddish-brown stems, each bearing 1 or 2 flowers, with red sepals and white petals, in late spring. Cultivation group 3. Z5–9

S. callosa, syn. *S. linguata*, illus. p.337; cultivation group 4.

S. cochlearis. Evergreen, rosetted perennial. H 8in (20cm), S 10in (25cm). Has spoon-shaped, green leaves with white-encrusted edges. Produces loose panicles of rounded, white flowers, often with red-spotted petals, in early summer. Z5–9. **‘Minor’**, H and S 5in (12cm), has smaller leaf rosettes and loose panicles of red-spotted, white flowers on red stems. Ideal for a trough. Cultivation group 4.

S. cortusifolia var. fortunei. See *S. fortunei*.

S. cotyledon. Evergreen perennial. H and S to 1ft (30cm). Has large, pale green rosettes of leaves, which die after flowering. In late spring–early summer, bears arching, conical panicles of white, cup-shaped flowers, sometimes strongly marked red inside. Cultivation group 2. Z3–9

S. ‘Cranbourne’. See *S. x anglica* ‘Cranbourne’.

S. ‘Crenata’. See *S. burseriana* ‘Crenata’.

S. cuneifolia illus. p.337; cultivation group 1.

S. exarata subsp. moschata ‘Cloth of Gold’ illus. p.377; cultivation group 1.

S. federici-augusti subsp. grisebachii ‘Wisley Variety’, syn. *S. grisebachii* ‘Wisley Variety’, illus. p.353; cultivation group 4. Z5–9

S. ferdinandi-coburgi. Evergreen, cushion-forming perennial. H and S 6in (15cm). Forms rosettes of linear, spiny, glaucous green leaves and, in early spring, bears racemes of open cup-shaped, rich yellow flowers on stems 1¼–4in (3–10cm) long. Cultivation group 3. Z5–9

S. fortunei, syn. *S. cortusifolia* var. *fortunei*. Semievergreen or herbaceous, clump-forming perennial. H and S 1ft (30cm). Rounded, 5- or 7-lobed, fleshy, green or brownish-green leaves are red beneath. In fall, bears panicles of tiny, mothlike, white flowers, with 4 equal-sized petals and one elongated petal, on upright stems. Propagate by division in spring. Z6–9. **‘Rubrifolia’** has dark red flower

stems and dark reddish-green leaves, beetroot-red beneath. Cultivation group 1.

S. x geum. Evergreen, mat-forming perennial. H 6–8in (15–20cm), S 1ft (30cm). Has shallow-rooted rosettes of spoon-shaped, hairy leaves. In summer, star-shaped, pink-spotted, white flowers, deep pink in bud, are borne on loose panicles on slender stems. Cultivation group 1. Z6–8

S. ‘Gloria’. See *S. burseriana* ‘Gloria’.

S. granulata (Fair maids of France, Meadow saxifrage) illus. p.332. **‘Plena’** is a clump-forming perennial. H 9–15in (23–38cm), S to 6in (15cm) or more. Loses its kidney-shaped, glossy, pale to mid-green leaves soon after flowering. Has a loose panicle of large, rounded, double, white flowers in late spring or early summer. Bulbils or resting buds form at base of foliage. Cultivation group 1. Z6–8

S. ‘Gregor Mendel’. See *S. x apiculata* ‘Gregor Mendel’.

S. grisebachii ‘Wisley Variety’. See *S. federici-augustii* subsp. *grisebachii* ‘Wisley Variety’.

S. hirsuta. Evergreen, mound-forming perennial. H 6–8in (15–20cm), S 8in (20cm). Has rosettes of round, hairy leaves. Bears loose panicles of tiny, star-shaped, white flowers, often yellow-spotted at the base of petals, in late spring and early summer. Cultivation group 1. Z5–9

S. ‘Irvingii’. See *S. x irvingii* ‘Walter Irving’.

S. x irvingii ‘Jenkinsiae’, syn. *S. ‘Jenkinsiae’*, illus. p.350; cultivation group 2.

‘Walter Irving’ (syn. *S. ‘Irvingii’*) is a slow-growing, evergreen, dense, mound-forming perennial. H ¾in (2cm), S 3in (8cm). Has minute leaves in neat rosettes. In early spring, bears stemless, cup-shaped, lilac-pink flowers. Cultivation group 3. Z5–9

S. ‘Jenkinsiae’. See *S. x irvingii* ‘Jenkinsiae’.

S. linguata. See *S. callosa*.

S. longifolia. Rosetted perennial. H 2ft (60cm), S 8–10in (20–25cm). Long, narrow, lime-encrusted leaves form attractive rosettes that, after 3–4 years, develop long, arching, conical to cylindrical panicles of numerous, rounded, 5-petaled, white flowers in late spring and summer. After flowering, rosettes die and no daughter rosettes are formed, so propagate by seed in spring or fall. Cultivation group 4. Z5–8

S. oppositifolia (Purple mountain saxifrage) illus. p.353. **‘Ruth Draper’** is an evergreen, loose mat-forming perennial. H 1–2in (2.5–5cm), S 6in (15cm). Has small, opposite, oblong to ovate, white-flecked, dark green leaves closely set along prostrate stems. Large, cup-shaped, deep purple-pink flowers are borne in early spring, just above foliage. Prefers peaty soil. Cultivation group 1. Z5–8

S. paniculata. Evergreen, tightly rosetted perennial. H 6–12in (15–30cm), S 8in (20cm). In summer, bears loose panicles of rounded, usually white flowers, with or without purplish-red spots, on upright stems above rosettes of oblong to ovate, lime-encrusted leaves. Very variable in size. Pale yellow or pale pink forms also occur. Cultivation group 4. Z5–8

S. sancta illus. p.358; cultivation group 3.

S. sarmentosa. See *S. stolonifera*.

S. scardica illus. p.348; cultivation group 3.

S. sempervivum illus. p.353; cultivation group 4.

S. Southside Seedling Group illus. p.338; cultivation group 3 or 4.

S. stolonifera, syn. *S. sarmentosa* (Mother of thousands). Evergreen, prostrate perennial with runners. H 6in (15cm) or more, S 1ft (30cm) or more. Has large, rounded, shallowly lobed, hairy, silver-veined, olive-green leaves that are reddish-purple beneath. Loose panicles of tiny, mothlike, white flowers, each with 4 equal-sized petals and one elongated petal, are borne in summer on slender, upright stems. Provides good ground cover. Z10–11. **‘Tricolor’** (syn. *S. ‘Tricolor’*; Strawberry geranium) has green-and-red leaves marked silver. Cultivation group 1.

S. stribnyi illus. p.354; cultivation group 3.

S. ‘Tricolor’. See *S. stolonifera* ‘Tricolor’.

S. ‘Tumbling Waters’ illus. p.333; cultivation group 4.

S. x urbium (London pride). Evergreen, rosetted, spreading perennial. H 1ft (30cm), S indefinite. Has spoon-shaped, toothed, leathery, green leaves. Flower stems bear tiny, star-shaped, at times pink-flushed, white flowers, with red spots, in summer. Useful as groundcover. Cultivation group 1. Z4–9

SCABIOSA

Scabiosa

CAPRIFOLIACEAE

Genus of annuals, biennials, and perennials, some of which are evergreen, with flower heads that are good for cutting. Fully to frost hardy. Prefers sun and fertile, well-drained, alkaline soil. Propagate annuals by seed in spring and perennials by cuttings of young, basal growths in summer, by seed in fall, or by division in early spring.

S. arvensis. See *nautia arvensis*.

S. atropurpurea (Sweet scabious). Moderately fast-growing, upright, bushy annual. H to 3ft (1m), S 8–12in (20–30cm). Has lance-shaped, lobed, mid-green leaves. Domed heads of pincushionlike, scented, deep crimson flower heads, 2in (5cm) across, are produced on wiry stems in summer and early fall. Tall forms, H 3ft (1m), and dwarf, H 1½ft (45cm), are available with flower heads in shades of blue, purple, red, pink or white. Z4–9

S. caucasica ‘Clive Greaves’ illus. p.271. **‘Miss Willmott’** is a clump-forming perennial. H and S 2ft (60cm). Large, creamy-white flower heads, with pincushionlike centers, are produced throughout summer. Light green leaves are lance-shaped at base of plant and segmented on stems. H4

S. columbaria subsp. ochroleuca, syn. *S. ochroleuca*. Clump-forming perennial. H and S 3ft (1m). In late summer, branching stems bear many heads of frilled, sulphur-yellow flower heads with pincushionlike centers. Has narrowly ovate, toothed, gray-green leaves. Z4–9

S. gigantea. See *Cephalaria gigantea*.

S. graminifolia. Evergreen, clump-forming perennial, often with a woody base. H and S 6–10in (15–25cm). Has tufts of narrow, grasslike, pointed, silver-haired leaves. In summer, produces stiff stems with spherical, bluish-violet to lilac flower heads like pincushions. Suits a rock garden. Resents disturbance. Z4–9

SCHEFFLERA

S. lucida illus. p.341.

S. ochroleuca. See *S. columbaria* subsp. *ochroleuca*.

S. rumelica. See *nautia macedonica*.

S. tartarica. See *Cephalaria gigantea*.

SCADOXUS

AMARYLLIDACEAE

Genus of bulbous and rhizomatous perennials, grown for their spectacular, mainly spherical, umbels of red flowers. Frost tender. Needs a position in partial shade and humus-rich, well-drained soil. Reduce watering in winter. Propagate by seed or offsets in spring.

S. multiflorus, syn. *Haemanthus multiflorus*. Summer-flowering bulb. H to 3ft (1m), S 1ft (30cm). Has broadly lance-shaped to ovate, semierect, basal leaves that form a “stem” of sheathed leaf stalks. Bears a spherical head, 4–10in (10–25cm) across, of up to 200 narrow-petaled, red flowers, followed by small orange berries. Z11–12 (min. 50–59°F/10–15°C).

subsp. katherinae (syn. *Haemanthus katherinae*; Blood flower) illus. p.386.

S. puniceus, syn. *Haemanthus magnificus*, *H. natalensis*, *H. puniceus* (Royal paintbrush). Spring- and summer-flowering bulb. H 20in (50cm), S 6in (15cm). Has elliptic, semierect leaves in a basal cluster. Leaf bases are joined, forming a false stem. Flower stem bears up to 100 tiny, tubular, greenish-white to pink or orange-red flowers in a conical umbel, surrounded by a whorl of red bracts, then red berries. Z11–12 (min. 50–59°F/10–15°C)

SCAEVOLA

GOODENIACEAE

Genus of short-lived, mainly evergreen perennials, but also scrambling climbers, shrubs, and small trees, grown for summer display, usually in containers. Frost tender. Grow in moist but well-drained, fertile compost in sun or partial shade. Propagate by softwood cuttings in late spring or summer.

S. aemula ‘Little Wonder’. Evergreen, trailing, bushy perennial, grown as an annual. H 1ft (30cm), S 16in (40cm). Has spoon-shaped, rich green leaves. In summer, produces 5-lobed, blue flowers each with a yellow ring round the white throat. Z10–11 (min. 41°F/5°C)

SCHEFFLERA

SYN. BRASSAIA, HEPTAPLEURUM

ARALIACEAE

Genus of evergreen shrubs, trees, and climbers grown mainly for their handsome foliage. Frost hardy to frost tender. Grows in any fertile, well-drained but moisture-retentive soil and in full light or partial shade. Water potted specimens freely when in full growth, moderately at other times. Pruning is tolerated if needed. Propagate by air-layering in spring, by semiripe cuttings in summer, or by seed as soon as ripe, in late summer.

S. delavayi. Small, bushy, broad-headed, evergreen tree. H 12–25ft (4–8m), S 6–12ft (2–4m). Stalks, to 2ft (60cm) long, bear leaves, composed of usually 5 dark green leaflets, 4–14in (10–35cm) long, densely covered in yellow-brown or gray-white