

TIBOUCHINA

semiripe cuttings in late spring–summer. **T. semidecandra** of gardens. See *T. urvilleana*. **T. urvilleana**, syn. *T. semidecandra* of gardens (Glory bush), illus. p.457.

TIGRIDIA

SYN. RIGIDELLA

IRIDACEAE

Genus of summer-flowering bulbs, grown for their highly colorful but short-lived flowers, rather irislike in shape, with 3 large, outer petals. Frost hardy to frost tender. Needs sun and well-drained soil, with ample water in summer. Plant in spring. Lift in fall; then partially dry bulbs and store in peat or sand at 46–54°F (8–12°C). Propagate by seed in spring. **T. pavonia** (Peacock flower, Tiger flower) illus. p.413.

TILIA

Lime, Linden

MALVACEAE

Genus of deciduous trees, grown for their foliage, small, fragrant, cup-shaped flowers, and stately habit. Flowers attract bees, but are toxic to them in some cases. Flowers are followed by dry, nutlike fruits. Fully hardy. Requires sun or partial shade and fertile, well-drained soil. Propagate species by seed in fall, selected forms and hybrids by grafting in late summer. Except for *T. x euchlora*, trees are usually attacked by aphids, which cover growth and the ground beneath with sticky honeydew. ①The nectar of

T x europaea, *T. x euchlora*, *T. tomentosa* and *T. tomentosa* ‘Petiolaris’ may be toxic, especially to bumblebees.

T. americana (American lime, Basswood). Deciduous, spreading tree. H 80ft (25m), S 40ft (12m). Has large, rounded, sharply toothed, glossy, dark green leaves and produces small, yellowish-white flowers in summer. Z2–8

T. cordata (Small-leaved lime). Deciduous, spreading tree. H 100ft (30m), S 40ft (12m). Has small, glossy, dark green leaves and produces small, yellowish-white flowers in midsummer. Z3–7.

‘Greenspire’, S 25ft (8m), is very vigorous and pyramidal in habit, even when young. **‘Winter Orange’** has red buds and orange young shoots in winter, and butter-yellow fall color; stool or pollard for best effect.

T. x euchlora (Caucasian lime, Crimean lime). Deciduous, spreading tree with lower branches that become pendent with age. H 70ft (20m), S 30ft (10m). Rounded, very glossy, deep green leaves turn yellow in fall. Bears small, yellowish-white flowers, which may be toxic to bees, in summer. Relatively pest-free. Z3–8

T. x europaea, syn. *T. x vulgaris* (Common lime). Vigorous, deciduous, spreading tree. H 120ft (35m), S 50ft (15m). Trunk develops many burs. Has rounded, dark green leaves. Small, yellowish-white flowers, which may be toxic to bees, are borne in summer. Periodically remove shoots from burs at base. Z3–7

T. henryana. Deciduous, spreading tree. H and S 30ft (10m). Broadly heart-shaped, glossy, bright green leaves, fringed with

long teeth, are often tinged red when young. Produces an abundance of small, creamy-white flowers in fall. Z7–9

T. mongolica (Mongolian lime). Deciduous, spreading, graceful tree. H 50ft (15m), S 40ft (12m). Young shoots are red. Heart-shaped, coarsely toothed, glossy, dark green leaves turn yellow in fall. Small, yellowish-white flowers are borne in summer. Z3–6

T. oliveri illus. p.63. **T. petiolaris**. See *T. tomentosa* ‘Petiolaris’.

T. ‘Petiolaris’. See *T. tomentosa* ‘Petiolaris’.

T. platyphyllos (Broad-leaved lime, Large-leaved lime). Deciduous, spreading tree. H 100ft (30m), S 70ft (20m). Has rounded, dark green leaves and bears small, dull yellowish-white flowers in midsummer. Z4–6. **‘Rubra’** (Red-twigged lime) illus. p.63.

T. tomentosa (European white lime, Silver lime). Deciduous, spreading tree. H 80ft (25m), S 70ft (20m). Leaves are large, rounded, sharply toothed, dark green above and white beneath. Very fragrant, small white flowers, which may be toxic to bees, are borne in late summer. Z4–7. **‘Petiolaris’** (syn. *T. petiolaris*, *T. ‘Petiolaris’*; Pendent silver lime) illus. p.64. **T. x vulgaris**. See *T. x europaea*.

TILLANDSIA

BROMELIACEAE

Genus of evergreen, epiphytic perennials, often rosette-forming, some with branching stems and spirally arranged leaves, all grown for their flowers or overall appearance. Frost tender. Requires partial shade. Provide a rooting medium of equal parts humus-rich soil and either sphagnum moss or bark. May also be grown on slabs of bark or sections of trees. Using soft water, water moderately in summer, sparingly at other times; spray plants grown on bark or tree sections with water several times a week from mid-spring to mid-fall. Propagate by offsets or division during spring. See also feature panel p.471.

T. argentea (illus. p.471). Evergreen, basal-rosetted, epiphytic perennial. H and S 4–6in (10–15cm). Very narrow, almost threadlike leaves, covered with white scales, are produced in dense, near-spherical rosettes, each with a fleshy, bulblike base. In summer, bears small, loose racemes of tubular, red flowers. Z10–12 (min. 45°F/7°C)

T. caput-medusae. Evergreen, basal-rosetted, epiphytic perennial. H and S 6in (15cm) or more. Linear, channelled, twisted and rolled, incurved leaves, covered in gray scales, develop in loose rosettes that have hollow, bulblike bases. In summer, spikes of tubular, violet-blue flowers are produced above the foliage. Z10–12 (min. 45°F/7°C)

T. cyanea. Evergreen, basal-rosetted, epiphytic perennial. H and S 10in (25cm). Forms dense rosettes of linear, pointed, channelled, arching, usually deep green leaves. In summer, broadly ovate, bladelike spikes of pansy-shaped, deep purple-blue flowers, emerging from pink or red bracts, are produced among foliage. Z11–12 (min. 50°F/10°C)

T. fasciculata. Evergreen, basal-rosetted, epiphytic perennial. H and S 1ft (30cm) or more. Has dense rosettes of narrowly triangular, tapering, arching, mid-green leaves. In summer, flat spikes of tubular, purple-blue flowers emerge from red or reddish-yellow bracts, just above leaf tips. Bracts require strong light to develop reddish tones. Z10–12 (min. 45°F/7°C)

T. ionantha (Sky plant). Evergreen, clump-forming, basal-rosetted, epiphytic perennial. H and S 5in (12cm). Linear, incurved, arching leaves, covered in gray scales, are produced in dense rosettes; the inner leaves turn red at flowering time. Spikes of tubular, violet-blue flowers, emerging in summer from narrow, white bracts, are borne just above foliage. Z10–12 (min. 45°F/7°C)

T. lindenii (Blue-flowered torch; illus. p.471). Evergreen, basal-rosetted, epiphytic perennial. H and S 16in (40cm). Linear, pointed, channelled, arching, mid-green leaves, with red-brown lines, form dense rosettes. In summer, produces bladelike spikes of widely pansy-shaped, deep blue flowers, emerging from sometimes pink-tinted, green bracts, which are borne just above leaves. Z11–12 (min. 50°F/10°C)

T. recurvata. Evergreen, basal-rosetted, epiphytic perennial. H and S 4–8in (10–20cm). Has long, loose, stemlike rosettes of linear, arching to recurved leaves, densely covered in silvery-gray scales. In summer, bears short, dense spikes of small, tubular, pale blue or pale green flowers above the leaves. Z10–12 (min. 45°F/7°C)

T. stricta (illus. p.471). Evergreen, clump-forming, basal-rosetted, epiphytic perennial. H and S 8–12in (20–30cm). Narrowly triangular, tapering, arching, mid-green leaves, usually with gray scales, are produced in dense rosettes. Large, tubular, blue flowers emerge from pendant, conelike spikes of brightred bracts, usually in summer. Z10–12 (min. 45°F/7°C)

T. usneoides (Spanish moss). Evergreen, pendent, epiphytic perennial. H 3ft (1m) or more, S 4–8in (10–20cm). Slender, branched, pendant stems bear linear, incurved leaves, densely covered in silvery-white scales. Inconspicuous, tubular, greenish-yellow or pale blue flowers, hidden among foliage, are borne in summer. Z10–12 (min. 45°F/7°C)

TITANOPSIS

AIZOACEAE

Genus of short-stemmed, basal-rosetted, perennial succulents eventually forming small, dense clumps. Produces 6–8 opposite pairs of fleshy, triangular leaves, ¼–1¼in (2–3cm) long, narrow at stems and expanding to straight tips. Frost tender. Requires a position in sun and well-drained soil. Propagate by seed in spring or summer.

T. calcarea illus. p.496. **T. schwantesii**. Clump-forming, perennial succulent. H 1¼in (3cm), S 4in (10cm). Has a basal rosette of triangular, gray-blue leaves, covered with small, wartlike, yellow-brown tubercles. Bears daisylike, light yellow flowers, ¾in (2cm) across, in summer–fall. Z10–11 (min. 41°F/5°C)

TITHONIA

ASTERACEAE/COMPOSITAE

Genus of annuals. Half hardy. Grow in sun and in fertile, well-drained soil. Provide support and deadhead regularly. Propagate by seed sown under glass in late winter or early spring.

T. rotundifolia (Mexican sunflower). **‘Fiesta del Sol’** Robust, bushy annual. H 2–3ft (60–90cm), S 1ft (30cm). Has long, entire or occasionally 3-lobed, toothed leaves, 8–30cm (3–12in) long, hairy beneath. Bears vibrant, orange flowers from late summer to fall. **‘Torch’** illus. p.324. Z10–11

TOLMIEA

SAXIFRAGACEAE

Genus of one species of perennial that is sometimes semievergreen and is grown as groundcover. Suitable for cool woodland gardens. Fully hardy. Prefers a position in shade and requires well-drained, neutral to acidic soil. Propagate by division in spring or by seed in fall.

T. menziesii (Pick-a-back-plant, Youth-on-age). Mat-forming perennial, sometimes semievergreen. H 18–24in (45–60cm), S 1ft (30cm) or more. Young plantlets develop where ivy-shaped, mid-green leaves join stem. Spikes of tiny, nodding, tubular to bell-shaped, green and chocolate-brown flowers are borne in spring. Z7–9

TOLPIS

ASTERACEAE/COMPOSITAE

Genus of summer-flowering annuals and perennials. Frost hardy. Grow in sun and in fertile, well-drained soil. Propagate by seed sown outdoors in spring.

T. barbata. Moderately fast-growing, upright, branching annual. H 1½–2ft (45–60cm), S 1ft (30cm). Has lance-shaped, serrated, mid-green leaves. Daisylike, bright yellow flower heads, 1in (2.5cm) or more across, with maroon centers, are produced in summer. Z3–10

TOONA

MELIACEAE

Genus of deciduous or semievergreen trees, grown for their foliage, fall color and flowers. Fully hardy. Prefers full sun; requires fertile, well-drained soil. Propagate by seed in fall, or root cuttings in winter.

T. sinensis, syn. *Cedrela sinensis*, illus. p.73.

TORENIA

LINDERNIACEAE

Genus of annuals and perennials. Frost tender. Often grown as summer bedding. Grow in partial shade and in a sheltered position in fertile, well-drained soil. Pinch out growing shoots of young plants to encourage a busy habit. Propagate by seed sown under glass in early spring. **T. fournieri** (Wishbone flower). Moderately fast-growing, erect, branching annual. H 1ft (30cm), S 8in (20cm). Has long-stalked, serrated, light green leaves.