

In summer and early fall, bears abundant lilac-blue flowers, the lower lips deep purple, the throats marked yellow. Z10–12 (min. 41°F/5°C). **Moon Series BLUE MOON (‘Dantmoon’)** (syn. *T. Moon Series BLUE MOON*), H 6–8in (15–20cm), S 1½ft (45cm), is dense and trailing with ovate, toothed, dark green leaves. Bears tubular, flared, 2-tone blue flowers with dark throats. **T. Moon Series BLUE MOON (‘Dantmoon’)**. See *T. fournieri* Moon Series BLUE MOON (‘Dantmoon’).

TORREYA
TAXACEAE
See also CONIFERS.
T. californica (California nutmeg) illus. p.101.
Tovara. See *Persicaria*.

TOWNSENDIA
ASTERACEAE/COMPOSITAE
Genus of evergreen, short-lived perennials and annuals, grown for their daisylike flowers. Suit alpine houses, as they dislike winter wet. Fully hardy. Needs sun and moist soil. Propagate by seed in fall. **T. grandiflora** illus. p.369. **T. parryi**. Evergreen, basal-rosetted, short-lived perennial. H 3–6in (7–15cm), S 2in (5cm). In late spring, produces daisylike, lavender or violet-blue flower heads, with bright yellow centers, above spoon-shaped leaves. Z4–8

TOXICODENDRON
ANACARDIACEAE
Genus of deciduous trees, shrubs, and woody vines, primarily grown for their foliage; many species have rich fall color. Fully to frost hardy. Requires sun and well-drained soil. Propagate by semiripe cuttings in summer, seed in fall or root cuttings in winter.

T. succedaneum, syn. *Rhus succedanea* (Wax tree). Deciduous, spreading tree. H and S 30ft (10m). Large, glossy, dark green leaves, each made up of 9–15 ovate leaflets, turn red in fall. Bears dense clusters of tiny, yellow-green flowers in summer. Female flowers develop into tiny, spherical, black or brownish fruits. ☉Contact with the foliage may cause skin irritation. Z4–9 **T. vernicifluum**, syn. *Rhus verniciflua* (Varnish tree). Deciduous, spreading tree. H 50ft (15m), S 30ft (10m). Large, glossy, bright green leaves, with 7–13 ovate leaflets, redden in fall. Bears dense clusters of tiny, yellow-green flowers in summer, followed by berrylike, brownish-yellow fruits. ☉All parts are highly toxic if ingested; contact with the sap may severely irritate the skin. Z8–10

TRACHELIUM
SYN. DIOSPHAERA
CAMPANULACEAE
Genus of small perennials, useful for rock gardens and borders. Some are good in alpine houses. Flowers of half-hardy species are ideal for cutting. Borderline

fully hardy to half hardy, but protect fully hardy species under glass in winter as they resent damp conditions. Grow in a sunny, sheltered site in fertile, very well-drained soil (*T. asperuloides* prefers lime-rich soil). Propagate by seed in early or mid-spring or by softwood cuttings in early summer. **T. asperuloides**, syn. *Diosphaera asperuloides*, illus. p.369. **T. caeruleum** (Throatwort) illus. p.313.

TRACHELOSPERMUM
APOCYNACEAE
Genus of evergreen, woody-stemmed, twining climbers with stems that exude milky sap when cut. Frost hardy. Grow in any well-drained soil and in sun or partial shade. Propagate by seed in spring, by layering in summer, or by semiripe cuttings in late summer or fall. **T. asiaticum** illus. p.195. **T. jasminoides** (Confederate jasmine, Star jasmine) illus. p.195. **‘Variegatum’** is a woody, evergreen, twining climber. H 10–15ft (3–5m). Has ovate, glossy, dark green leaves splashed and margined creamy white, often suffused red and pink in winter. In mid- and late summer, bears cymes of pure white flowers. Z8–11

TRACHYCARPUS
ARECACEAE/PALMAE
Genus of evergreen, summer-flowering palms, grown for their architectural habit, foliage and flowers. Fully to frost hardy. Needs full sun and does best if sheltered from strong, cold winds, especially when young. Needs fertile, well-drained soil. Propagate by seed in fall or spring. **T. fortunei** (Chusan palm, Windmill palm) illus. p.80. **T. wagnerianus**. Dwarf, evergreen palm. H 10–20ft (3–6m), S 6–10ft (2–3m). Has smaller, more rigid, leathery leaves, on shorter stalks than *T. fortunei*, and the segments edged with white hairs. Wind and drought tolerant. Z6–10

TRACHYMENE
APIACEAE/UMBELLIFERAE
Genus of summer-flowering annuals, biennials and perennials. Half hardy. Grow in a sunny, sheltered position and in fertile, well-drained soil. Support with sticks. Propagate by seed sown under glass in early spring. **T. coerulea**, syn. *Didiscus coeruleus* (Blue lace flower). Moderately fast-growing, upright, branching annual. H 1½ft (45cm), S 8in (20cm). Has deeply divided, pale green leaves. Spherical heads, to 2in (5cm) across, of tiny, blue flowers are produced in summer. Excellent for cutting. Z10–11

TRADESCANTIA
Spiderwort
COMMELINACEAE
Genus of perennials, some of which are evergreen, grown for their flowers or ornamental foliage. Fully hardy to frost tender. Grow in fertile, moist to dry soil and in sun or partial shade. Cut back or repropagate trailing species when they become straggly. Propagate hardy species

by division or seed, and frost-tender species by tip cuttings in spring, summer or fall. Hardy species are useful in shady gardens. ☉Contact with the foliage may cause skin irritation. **T. albiflora**. See *T. fluminensis*. **T. Andersoniana Group ‘J.C. Weguelin’**, syn. *T. ‘J.C. Weguelin’*. Clump-forming perennial. H to 2ft (60cm), S 1½ft (45cm). Has narrowly lance-shaped, fleshy, green leaves, 6–12in (15–30cm) long. In summer, produces clusters of 3-petaled, lavender-blue flowers, 1in (2.5cm) or more across, surrounded by 2 leaflike bracts. H6. **‘Osprey’** (syn. *T. ‘Osprey’*) illus. p.263. **‘Purple Dome’** (syn. *T. ‘Purple Dome’*) illus. p.269. **T. blossfeldiana**. See *T. cerinthoides*. **T. cerinthoides**, syn. *T. blossfeldiana*. Evergreen, creeping perennial. H 2in (5cm), S indefinite. Narrowly ovate, fleshy, stem-clasping leaves, to 4in (10cm) long, are glossy, dark green above, purple with long, white hairs below. Intermittently produces clusters of tiny, pink flowers, with white centers, surrounded by 2 leaflike bracts. Z10–12 (min. 41–45°F/5–7°C). Leaves of **‘Variegata’** have long, cream stripes. **T. fluminensis**, syn. *T. albiflora* (Wandering Jew). Evergreen perennial with trailing, rooting stems. H 2in (5cm), S to 2ft (60cm) or more. Ovate, fleshy leaves, 1½in (4cm) long, that clasp the stem, are glossy green above, sometimes tinged purple below. Intermittently has clusters of tiny, white flowers enclosed in 2 leaflike bracts. Z10–12 (min. 41–45°F/5–7°C). **‘Albovittata’** illus. p.468. **‘Aurea’** (syn. *T.f. ‘Variegata’*) H 1ft (30cm), S indefinite, has striped creamy-white leaves. **‘Variegata’** see *T.f. ‘Aurea’*. **T. ‘J.C. Weguelin’**. See *T. Andersoniana Group ‘J.C. Weguelin’*. **T. navicularis**. See *Callisia navicularis*. **T. ‘Osprey’**. See *T. Andersoniana Group ‘Osprey’*. **T. pallida ‘Purpurea’**, syn. *T.p. ‘Purple Heart’*, *Setcreasea purpurea*, illus. p.310. **T. pexata**. See *T. sillamontana*. **T. ‘Purple Dome’**. See *T. Andersoniana Group ‘Purple Dome’*. **T. purpusii**. See *T. zebrina ‘Purpusii’*. **T. sillamontana**, syn. *T. pexata*, *T. velutina*, illus. p.469. **T. spathacea**, syn. *Rhoeo discolor*, *R. spathacea* (Boat lily, Moses-in-the-cradle). Evergreen, clump-forming perennial. H 20in (50cm), S 10in (25cm). Rosette of lance-shaped, fleshy leaves, to 1ft (30cm) long, is green above, purple below. Bears tiny, white flowers, enclosed in boat-shaped, leaflike bracts, year-round. Z10–12 (min. 41–45°F/5–7°C). **‘Vittata’** produces leaves striped longitudinally with pale yellow. **T. velutina**. See *T. sillamontana*. **T. zebrina**, syn. *Zebrina pendula* (Silver inch plant), illus. p.469. **‘Purpusii’** (syn. *T. purpusii*) is a strong-growing, evergreen, trailing or mat-forming perennial. H 4in (10cm), S indefinite. Has elliptic, purple-tinged, bluish-green leaves and tiny, shallowly cup-shaped, pink flowers. **‘Quadricolor’** has leaves striped green, pink, red and white. Z10–12 (min. 41–45°F/5–7°C).

TRICYRTIS

TRAPA
Water chestnut
LYTHRACEAE/TRAPACEAE
Genus of annual floating water plants, grown for their foliage and flowers. Half hardy to frost tender. Requires sun. Propagate in spring from seed gathered in fall and stored frost-free, in water, or damp moss. **T. natans** (Water chestnut) illus. p.443.

Trichocereus bridgesii. See *Echinopsis lageniformis*. **Trichocereus candicans**. See *Echinopsis candicans*. **Trichocereus spachianus**. See *Echinopsis spachiana*.

TRICHODIADEMA
AIZOACEAE
Genus of bushy, perennial succulents with woody or tuberous roots and cylindrical to semi-cylindrical leaves. Frost tender. Needs a sunny position and well-drained soil. Propagate by seed or stem cuttings in spring or summer. **T. densum**. Tufted, perennial succulent. H 4in (10cm), S 8in (20cm). Cylindrical, pale green leaves are each ½–¾in (1–2cm) long and tipped with clusters of white bristles. Roots and prostrate, green stem are both fleshy and form caudex. The stem tip bears daisylike, cerise-pink flower heads, 1¼in (3cm) across, in summer. Z10–12 (min. 41°F/5°C) **T. mirabile** illus. p.481.

TRICHOSANTHES
CUCURBITACEAE
Genus of annual and evergreen, perennial, tendrill climbers, grown for their fruits and overall appearance. Frost tender. Needs full sun or partial shade and humus-rich soil. Water freely when in growth, less in cool weather. Provide support. Propagate by seed in spring at 70°F (21°C) or above. **T. anguina**. See *T. cucumerina* var. *anguina*.

T. cucumerina var. *anguina*, syn. *T. anguina* (Snake gourd). Erect to spreading, annual, tendrill climber. H 10–15ft (3–5m). Broadly ovate to almost triangular, sometimes shallowly 3- to 5-lobed, mid-to pale green leaves are to 8in (20cm) long. In summer, bears 5-petaled, white flowers, 1–2in (2.5–5cm) across, with heavily fringed petals; females are solitary, the males in racemes. Cylindrical fruits, 2ft (60cm) or rarely to 6ft (2m) long, often twisted or coiled, are green-and-white striped and ripen to dull orange. Z12 (min. 59°F/15°C)

Trichosma suavis. See *Eria coronaria*. **Tricospidaria lanceolata**. See *Crinodendron hookerianum*.

TRICYRTIS
Toad lily
LILIACEAE/CONVALLARIACEAE
Genus of late summer- and fall-flowering, rhizomatous perennials. Fully hardy.