



Mollusks

The second-largest phylum of invertebrates is the mollusks. Mollusks range from filter-feeding bivalves and grazing gastropods to highly intelligent cephalopods. All mollusks share a common body plan. The main muscle is the "foot," which is used for locomotion in snails. In cephalopods, the foot is divided into tentacles, while bivalves use it to move and dig.




◁ Bivalve

Mollusks in this class have two hinged shells, or valves, made from calcium carbonate. They include clams and mussels. Bivalves are exclusively aquatic, and most filter food from the water using hairlike cilia on their gills.



△ Cephalopod

This class includes octopuses, squid, and the nautilus. All but the latter have evolved out of their shells. They catch food with suckered—and in many cases clawed—tentacles that surround a beaklike mouth. They squirt a jet of water from a funnel near their mouth, called the siphon, to move.



△ Gastropod

This class of mollusks includes snails, slugs, winkles, and limpets. They have one shell—although this can be either reduced in size or absent completely in slugs. Snails and slugs are the only mollusks to live on land, although they require damp habitats. Snails breathe using a lunglike cavity in the mantle.

Worms

Worms are simple animals. They all lack legs, but can live in a wide range of habitats from the deep sea to inside the bodies of other animals. About half of the nematodes, also known as roundworms, are intestinal parasites, while the rest live in soil. The platyhelminthes, or flatworms, are parasitic or aquatic. They do not have intestines, and absorb food through their skin.

▷ Annelid

Also known as segmented worms, the Annelid phylum includes ragworms living in the ocean, oligochaetes such as earthworms on land, and leeches, which can live in freshwater or on land. Small, hairlike structures called setae help earthworms to burrow and sense their environment, while a series of pseudohearts pumps blood around their bodies.

