

Arctocebus calabarensis

Angwantibo



Length 9–10 in (25–35 cm)
Tail $\frac{3}{8}$ in (1 cm)
Weight 8–17 oz (225–475 g)
Location W. Africa
Social unit Group
Status Least concern




One of only 2 *Arctocebus* species, this one is orange to yellow on its upperparts and buff beneath. It climbs carefully and deliberately using all 4 equal-length limbs. The second toe is tiny and the first widely separated from the other 3, giving a clamplike grip. Solitary and nocturnal, the angwantibo eats small creatures such as caterpillars, plus a few types of fruit. The other species, the golden angwantibo, lives farther south.



Loris lydekkerianus

Gray slender loris



Length 8–10 $\frac{1}{4}$ in (21–26 cm)
Tail None
Weight 3–13 oz (180–294 g)
Location S., E., S.W. India, Sri Lanka
Social unit Individual/Pair
Status Least concern



FOUR-WAY GRIP
On each foot the big toe opposes the other 4 toes for a pincerlike grip. The gray slender loris can even sleep holding onto branches.

This small, slim primate moves with great deliberation, gripping twigs with all 4 feet—until it suddenly smells, sees, and, with its front legs, snatches a small creature such as an insect or lizard. It also eats soft leaves and buds, fruit, and birds' eggs. By day, the gray slender loris curls up in a tree-hole, dense leaf nest, or a similarly secure place. Its fur color ranges from yellow-gray to dark brown on the back, with silver-gray underparts.

REARING A FAMILY

The female slender loris reaches sexual maturity at 10 months and is receptive to the male twice a year thereafter. After a gestation period of 157–162 days, she bears 1–2 young, who are weaned after 148 days.



Nycticebus coucang

Sunda slow loris



Length 12–13 $\frac{1}{2}$ in (30–34 cm)
Tail $\frac{3}{8}$ – $\frac{3}{4}$ in (1–2 cm)
Weight 22–30 oz (635–850 g)
Location S.E. Asia
Social unit Variable
Status Vulnerable



Aptly named because of its lack of speed, the Sunda slow loris has gripping hands like the gray slender loris (see left) and is also a nocturnal tree-dweller. It creeps carefully toward its prey, then lunges with its front limbs. The loris lives alone or in pairs or groups, and the adult male chases other males from its territory, which is marked by urine. The dense, soft fur is brown with a white face and underparts, and dark eye rings and ears.

Perodicticus potto

West African potto



Length 12 in (30 cm)
Tail 1 $\frac{1}{2}$ –2 $\frac{1}{4}$ in (4–6 cm)
Weight 1 $\frac{3}{4}$ –2 $\frac{1}{4}$ lb (0.85–1 kg)
Location W. and C. Africa
Social unit Variable
Status Least concern



A careful, nocturnal climber, the West African potto has very mobile limb joints and reaches out at any angle to bridge gaps between branches. Its fur may be gray, brown, or red; the eyes and ears are small; and the diet consists mainly

of fruit, sap, gum, and small animals. A potto can remain immobile for hours to escape attention. If attacked, it will tuck down its head and batter the enemy with the “shield” of horny skin covering spiny bones on the upper back. The gestation period is about 200 days, and the potential life span is over 25 years.



Otolemur crassicaudatus


Brown greater galago



Length 10–16 in (25–40 cm)
Tail 13 $\frac{1}{2}$ –19 $\frac{1}{2}$ in (34–49 cm)
Weight 2 $\frac{1}{4}$ –4 $\frac{1}{2}$ lb (1–2 kg)
Location C., E., and southern Africa
Social unit Variable
Status Least concern



The largest galago, this species locates insects at night using its huge eyes and ears, and snatches them by hand in a split second. It also scrapes gum and sap with its comblike, protruding lower incisor and canine teeth. Deriving its alternative name of bushbaby from its childlike wails, it lives in small family groups of a male–female pair or 2 females, with young. Compared to closely related species, it tends to run on all fours more frequently and to leap in an upright position less often.



Euoticus elegantulus

Western needle-clawed galago



Length 7 $\frac{1}{2}$ –9 in (19–23 cm)
Tail 10–12 $\frac{1}{2}$ in (26–32 cm)
Weight 10–13 oz (270–360 g)
Location W. Africa
Social unit Individual/Pair
Status Least concern



Resembling other galagos in form, this species has an orange back, gray underside, pink hands and feet, and a long, gray, white-tipped tail. The oval eyes are rimmed by pale fur, and thin, sharp claws grip well for climbing. Like all galagos and lorises, the second toe on the rear foot has an upward-facing “toilet” claw, for scratching and combing fur. The specially enlarged front teeth scrape wood and bark to obtain gum and sap, which make up three-quarters of the diet, along with a variety of fruit and insects. It may make 1,000 “gum scrapes” per night.