

*Cacajao calvus*

Bald uakari



Location N.W. South America

Length 15–22½ in (38–57 cm)

Tail 5½–7½ in (14–18.5 cm)


Weight 5–7¼ lb (2.3–3.5 kg)

Social unit Group

Status Vulnerable



Bald uakaris prefer “blackwater,” part-flooded forests along small rivers, lakes, and swamps, rather than forests fringing large rivers. They forage by day in trees for seeds, fruit, flowers, and small animals, in a large troop of males and females, usually 10–20 but sometimes up to 200. The troop may mix with similar primates, such as squirrel monkeys, to feed.



HAIRLESS FACE

The bald uakari has a hairless face and forehead, the skin color varying from pink to deep red.

short tail in relation to body

COLOR FORMS



Bald uakari subspecies have a variety of fur colors, leading to different names: white uakari (above) in northwestern Brazil; golden uakari on the Brazil–Peru border; red uakari on the Colombia–Brazil border; and pale-backed red uakari further east.

*Aotus lemurinus*

Lemurine night monkey



Location N.W. South America

Length 12–12½ in (30–32 cm)

Tail 13½–13¾ in (34–34.5 cm)

Weight 29–36½ oz (800–1,050 g)

Social unit Pair

Status Vulnerable



Formerly regarded as one species, genetic studies show there may be 11 species of Lemurine night monkeys, also called douroucoulis, or owl monkeys. They are the only nocturnal monkeys, eating a mixed diet of fruit, leaves, and insects as they clamber cautiously through the branches. They live in male–female pairs and communicate by scents in urine and chest gland secretions. The single young is born after a gestation of 120 days. Weaning takes 8 months and, since juveniles may also stay with parents, close-knit family groups of 4–5 develop.



speckled gray fur on back

dark, bushy tail tip

yellow or gray underparts

*Callicebus aureipalatii*

Golden palace monkey



Location C. South America

Length 11½–12½ in (29–32 cm)

Tail 19–20¾ in (48–52.5 cm)

Weight 32–36 oz (900–1,000 g)

Social unit Group

Status Least concern



*Sapajus paella*

Guianan brown capuchin



Location N. South America

Length 15–18 in (38–46 cm)

Tail 15–15½ in (38–39 cm)

Weight 2¾–11 lb (1.3–4.8 kg)

Social unit Group

Status Least concern



*Callicebus moloch*

Red-bellied titi



Location N. South America

Length 10½–17 in (27–43 cm)

Tail 14–22 in (35–55 cm)


Weight 25–43 oz (700–1,200 g)

Social unit Pair

Status Least concern




The 30 or more titi species all have thick, soft fur, chunky bodies, short limbs, and—unlike other New World monkeys of their size—ears almost hidden in fur. They eat mainly fruit, leaves, seeds, and grubs. The red-bellied titi monkey’s back is speckled brown, the underparts mainly orange. It relies on its drab coloration and slow movements for camouflage in trees near swamps and pools. Female and male form a close pair-bond and defend a territory of 15–30 acres (6–12 hectares). They intertwine tails and sing a “duet” just after dawn to maintain their family and pair-bonds, and to proclaim their territory.



A colorful monkey, with orange-gold coat, white-tipped tail, and reddish hands and feet, this species was named after the organization that maintains the Madidi National Park in Bolivia. It is also known as the Madidi titi. Here, the species was discovered in 2004—in the forest of the Andean foothills and floodplain. It is likely that it ranges into southern Peru, too. Groups have a home range of between 42 and 62 acres (17 and 25 hectares) and consume a wide range of rain forest plants. Although much of its population is within a protected area and is not under significant pressure from hunting, the species may be affected by petroleum exploitation, road-building, and hydroelectric projects.

Capuchins are often regarded as the most adept New World monkeys, and the Guianan brown capuchin uses a variety of tools, such as stones to crack hard nuts. It also eats fruit, insects, and some vertebrates, such as frogs, lizards, and even small bats. This species, also known as the tufted capuchin from the furry “horns” above each ear, has the widest range of any New World monkey. Mixed groups of 10–20 are usual as members are not sexually mature until 7 years old—later than most monkeys of similar size.



prehensile tail often curled at tip