

Cebus olivaceus

Weeper capuchin



Location N.E. South America

Length 14½–18 in (37–46 cm)
Tail 16–22 in (40–55 cm)
Weight 5½–8¾ lb (2.3–4.2 kg)
Social unit Group
Status Least concern



Like the other 21 or so capuchin species, the weeper has a robust build with relatively short limbs, and a prehensile tail. There is no bare skin on the tail's underside, as in many



Saimiri boliviensis

Black-capped squirrel monkey



Location W. to C. South America

Length 10½–12½ in (27–32 cm)
Tail 15–16½ in (38–42 cm)
Weight 25–36 oz (700–1,000 g)
Social unit Group
Status Least concern



No other New World monkeys form such large, active troops as the 7 species of squirrel monkey. They regularly number 35–65—but can reach 200 or more—moving noisily by day with twitters



SQUIRREL FEATURES
The black-capped squirrel monkey has a small, white face, a large forehead, dark head crown above, black nose tip and muzzle, and well-furred ears.

Saguinus imperator

Emperor tamarin



Location W. South America


Length 9–10 in (23–26 cm)
Tail 14–16½ in (35–42 cm)
Weight 14–20 oz (400–550 g)
Social unit Group
Status Least concern



Marmosets and tamarins form a distinct group of about 47 American primate species. They are similar to other New World monkeys but differ in body chemistry, have claws rather than nails, and bear 2 rather than 1 offspring. Identified by its flowing white mustache, this species feeds on fruit in the wet season, flower nectar and tree sap in the dry season, and insects, especially crickets, all year round. It often forms a mixed troop with related species such as the saddleback tamarin. Each species responds to the other's alarm calls if a predator is detected nearby.

MUSTACHE

The white, curly-ended mustache of both males and females contrasts with the black face, speckled red- or gray-brown body, and red-orange tail fur.



TAMARIN BABY CARE



Tamarins and marmosets have relatively long gestation periods for such small mammals, the emperor tamarin's being 140–145 days. There are almost always 2 offspring, and the father carries these, as here, except when they are being suckled by their mother.

and clucks to disturb their food of various small creatures. They may follow in the path of other monkeys to obtain insects in their wake. They also consume fruit and seeds. The mature male becomes “fatted” around the shoulders in the breeding season and competes aggressively—the winner mating with the most females.

FAMILY MEAL



Within a squirrel monkey troop, members coalesce into peer subgroups: adult males, pregnant females, females with young, and juveniles. If an individual locates food, subgroup members quickly gather around to exploit the find.

Callimico goeldii

Goeldi's marmoset



Location N.W. South America

Length 9 in (19–25 cm)
Tail 10–12½ in (26–35 cm)
Weight 21 oz (355–366 g)
Social unit Group
Status Vulnerable



Goeldi's marmoset is larger than most marmosets and tamarins. Its long fur is black, and it has a “cape” of longer hair on its head and neck. Unlike similar species, it has wisdom teeth. It eats sap and gum (digging its incisor teeth into bark to make these flow), and fruit, insects, and small vertebrates such as lizards. The species forms stable, close-knit groups of up to 2–12, mainly male–female pairs and their young, and keeps to dense vegetation such as creeper-tangled bamboo.

