

Erythrocebus patas

Patas monkey



Length 19–35 in
(48–88 cm)
Tail 25–28 in
(63–72 cm)
Weight 8¼–29 lb
(4–13 kg)
Location W. to E. Africa
Social unit Group
Status Least concern



This is one of the fastest monkeys when it is running. It has a long, slim body, lengthy limbs, hands, and feet, and short digits. The white moustache and beard contrast with the darker face; the red back contrasts with the white limbs and underside. It lives in troops with an average of 15 members, and the single male stays on the periphery, to act as decoy, drawing predators while the females and young hide.

Cercopithecus neglectus


De Brazza's monkey



Length 17½–22 in
(44–56 cm)
Tail 23–28 in
(59–70 cm)
Weight 7–12 lb
(3.2–5.5 kg)
Location C. to E. Africa
Social unit Pair
Status Least concern



One of the most terrestrial of the 23 *Cercopithecus* species, De Brazza's monkey is also the only one of the genus that forms male–female pair bonds. Widespread yet inconspicuous over its large range, it marks its territory with saliva and scent, yet avoids rather than challenges any intruders. It has blue-white upper lip and chin fur, and a thin, white thigh stripe. It uses deep, booming calls to communicate, and eats mainly seeds and fruit.



Rungwecebus kipunji

Kipunji




Length 33½–35½ in
(85–90 cm)
Tail 39½–43½ in
(100–110 cm)
Weight 22–35 lb
(10–16 kg)
Location E. Africa
Social unit Group
Status Critically endangered



Discovered in 2003, this omnivorous tree-dwelling primate from the Udzungwa Mountains and Southern Highlands of Tanzania was originally thought to be a mangabey (a type of forest monkey)—but later shown to be more related to ground-dwelling baboons. It may even

have originated by crossbreeding between the two groups. It has a barking call, reminiscent of the grunts of baboons. Two isolated populations together inhabit scarcely 38 square miles (100 square km), and the species is threatened by deforestation and hunting.



Papio anubis

Olive baboon



Length 20–35 in
(50–90 cm)
Tail 15–23½ in
(38–60 cm)
Weight 31–66 lb
(14–30 kg)
Location W. to E. Africa
Social unit Group
Status Least concern



One of the largest baboons, and common across west and northeast Africa, the male may be twice the female's weight. Both sexes have a thick, gray ruff around the cheeks. They live in mixed groups of 1–120, that tend to be smaller in West Africa (7–40) than those in East Africa (30–120). They eat fruit, leaves, insects, lizards, and sometimes larger prey such as gazelle fawns.


GROUP ORDER



Female offspring tend to have a position directly below their mother, which depends on her position in the hierarchy. They don't take their place at the base unless the female has the lowest rank.

Macaca munzala

Arunachal macaque



Length 23 in (58 cm)
Tail 2¼ in (26.4 cm)
Weight 31 lb (14 kg)
Social unit Group
Status Endangered

Location S. Asia



Restricted to the Indian Arunachal Pradesh state of eastern Himalaya, this macaque was scientifically described in 2005 on the basis of photographs, and differs from related species in having a dark crown patch and pale neck collar. It occurs between 6,650 and 11,483 ft (2,000–3,500 m) in coniferous and broadleaf forests, as well as agricultural land. Like other high altitude monkeys it is robustly built to withstand the cold. Although densely populated in places and relatively tolerant of humans, the Arunachal macaque may be exposed to retaliatory hunting when it raids crops: this could threaten its long-term survival. So far the species is known with certainty from no more than 5 locations within an area of about 1,042 square miles (2,700 square km)—but it may also occur in adjacent areas of Bhutan and Tibet. Field data suggest that there are less than 600 known individuals.

Macaca nigra


Celebes crested macaque



Length 17½–22½ in
(44–57 cm)
Tail ½–1 in (1.5–2.5 cm)
Weight 12–29 lb
(5.5–13 kg)
Location S.E. Asia
(northern Sulawesi)
Social unit Group
Status Critically endangered



The all-black fur, very short tail, and homeland of Celebes (now Sulawesi) give this species the alternative name of Celebes black ape. A crest runs from the forehead back over the crown. Usually flat, it rises when the animal is aroused. This species forms huge, mixed-sex troops of more than 100, yet is generally an inconspicuous, quiet, fruit-eating forest dweller.



Papio papio

Guinea baboon



Length 14–34 in
(35–86 cm)
Tail 17–28 in
(43–70 cm)
Weight 22–57 lb
(10–26 kg)
Location W. Africa
Social unit Group
Status Near threatened



The male of the smallest baboon species has proportionally the longest mane, reaching almost to his rump. Even for a baboon, the diet is

varied: from tough roots to juicy grubs and eggs, and sometimes farm crops. Troops number up to 200, although about 40 is more usual, with several males gathering their own harem of females.



PRIMATE "DOG"

Typically powerful and doglike, the olive baboon is speckled olive-green, with a black face and rump.