

Hylobates moloch

Moloch gibbon




Length 18–25 in
(45–64 cm)
Tail None
Weight 13–15 lb
(6–7 kg)
Social unit Pair
Status Endangered

Location S.E. Asia



Pale eyebrows, cheeks, and beard merge with the mainly silvery fur of this gibbon. Its diet is fruit, some leaves, and sometimes nectar and grubs. Like most other gibbons, the family group (female, male, offspring) use calls to defend their territory. Unlike the lar gibbon (p.151), the male and female do not “duet” together.

dark
blue-gray
head cap



Pan paniscus

Bonobo



Length 28–33 in
(70–83 cm)
Tail None
Weight Up to 95 lb
(43 kg)
Social unit Group
Status Endangered


Location C. Africa




The bonobo or pygmy chimpanzee is only slightly smaller than the common chimpanzee (see below), but has a slimmer body and relatively longer and more slender limbs. It feeds chiefly on the ground, mainly on fruit and seeds, but also on leaves, flowers, fungi, eggs, and small animals. This ape can live in groups of up to 120 but is usually found in smaller groups, as it

forages and grooms. Sexual relations are common between males, females, and young in various combinations, and may be used to ease social tensions. Females are dominant and leave their family groups when mature; males tend to stay.

FACE AND FUR
The skin of the pygmy chimpanzee is mostly black, even on a juvenile's face. The fur on the crown has a central “parting.”



LONG INFANCY



The bonobo is born after 8 months' gestation, and suckled by its mother for up to 3 years. The female continues to protect, groom, and share nests with her offspring for another year or 2. The interval between births is about 5 years.

Pan troglodytes

Common chimpanzee



Length 28–38 in
(70–96 cm)
Tail None
Weight 44–200 lb
(20–70 kg)
Social unit Group
Status Endangered


Location W. to C. Africa



Common chimpanzees live in communities of up to 150. Subgroup composition varies almost hourly for activities such as grooming, feeding, traveling, and defending the territory. This last task is usually carried out by adult male parties, who may attack and kill stray chimps from other communities. Most daylight hours are spent eating—mainly fruit and leaves, but also flowers and seeds. Raiding parties sometimes cooperate to kill and eat animal prey, such as monkeys, birds, and small antelopes. Social bonds may last years, but there are no long-term male–female bonds for reproduction. The single young (rarely twins), born after a gestation period of 8 months, is fed, carried, and groomed by its mother for 3–4 years. It also learns her feeding techniques. Common chimpanzees not only use


tools but also make them—for example, stripping side branches from a twig, which it uses to scoop out termites from their nest. The two chimpanzee species are our closest living relatives, and their intelligence, range of emotions, and communication and learning skills have made them valuable to animal trainers, collectors, and researchers. They are also killed for the bushmeat trade.

bare skin on
face darkens
with age




VOCAL CHIMPANZEES

The common chimpanzee makes more than 30 distinct calls, including the pant-hoot (shown right), which consists of shrieks and roars that can be heard up to 1¼ miles (2km) away. It is used in many situations and is the most common adult sound. It is thought to identify the caller within its community and solicit information from other members.



FACIAL FEATURES
The common chimpanzee uses a wide variety of expressions, clearly visible on the hairless facial area. In particular, it uses its flexible, protrusile lips to make grimacelike “smiles” that actually signify fear.

NIGHT NESTS



Each adult common chimpanzee builds a new, individual tree nest each night for sleeping. (Rarely, an old nest is renovated or reused.) The common chimpanzee bends over and intertwines many branches to make a firm, leafy platform, usually 10–33 ft (3–10 m) above the forest floor, away from ground-based predators. Infants generally sleep in their mothers' nests until they are about 5–6 years of age.