

Kobus thomasi

Ugandan kob



Location C. and E. Africa

Length 5¼–6 ft
(1.6–1.8 m)Tail 4–6 in
(10–15 cm)Weight 60–120 lb
(130–265 kg)

Social unit Group

Status Least concern

The Ugandan kob is a graceful but strong grazing antelope, pale cinnamon to brown-black, with white facial and throat markings, and black leg stripes and feet. The male has ringed, lyre-shaped horns. Kobs live at very high densities, and males compete for a patch of ground (known as the lek), perhaps only 50 ft (15 m) across. The winner of this contest gains mating rights over many females.

*Kobus vardonii*

Puku



Location W. to E. Africa


Length 4¼–5 ft
(1.3–1.5 m)Tail 11–13 in
(28–32 cm)Weight 105–200 lb
(48–91 kg)

Social unit Individual/Group

Status Near threatened

The puku resembles the kob (see left) in its breeding system: it has leks at high population densities; territories at lower ones. The long coat is a uniform golden

yellow, and the horns are about 20 in (50 cm) in length. The puku grazes morning and evening and, like other plains antelopes, flees rapidly from danger.

*Redunca bohor*

Bohor reedbuck



Location C. and E. Africa

Length 3¼–4¼ ft
(1–1.3 m)Tail 7–7½ in
(18–20 cm)Weight 79–120 lb
(36–55 kg)

Social unit Variable

Status Least concern

This small, fawn, lightweight savanna antelope has white underparts, throat, and eye-rings. A conspicuous gray patch under the ear marks a scent

gland. The bohor reedbuck feeds on grasses and tender reed shoots. Small groups, of a female and young, or horned males, merge into dry-season herds.

*Hippotragus equinus*

Roan antelope




Location W., C., and E. Africa

Length 6½–7¼ ft
(2–2.1 m)Tail 23½–29½ in
(60–75 cm)Weight 470–620 lb
(215–280 kg)

Social unit Individual/Group

Status Least concern

The roan antelope is red- to brown-coated, with white underneath and black and white facial markings. Both sexes are horned and maned. They survive on poor grass growth and need to drink 2–3 times daily. Herds consist of 6–20 females and young which range over the territory of a single or several males.

*Hippotragus niger*

Sable antelope



Location E. to S.E. Africa

Length 6¼–8¾ ft
(1.9–2.7 m)Tail 14½–30 in
(37–76 cm)Weight 330–660 lb
(150–300 kg)


Social unit Group

Status Least concern

Similar to the roan antelope (see above right) in many respects, the sable antelope gathers in herds of 100 or more during the dry season, when

ADULT MALE

Mature males are black but with the same facial pattern as females. The lower jaw and muzzle are white. Males have larger, more curved horns than females.



SABLE CALF

Calf coloration is similar to that of adult females—chestnut or sorrel. It is born after 240–280 days' gestation, and initially is kept away from the herd.

browsing replaces the usual grazing. In the wet season, herds split into bachelor groups of 2–12 males, while dominant males occupy territories and mate with the females there. After an initial concealment, calves form nursery groups, only joining their mothers to suckle.

Oryx dammah

Scimitar-horned oryx



Location N. Africa

Length 5¼–5¾ ft
(1.5–1.7 m)Tail 14½–17¼ in
(37–44 cm)Weight 395–440 lb
(180–200 kg)

Social unit Group

Status Extinct in the wild

Specialized for deserts, arid plains, and rocky hills, the scimitar-horned oryx has many physiological adaptations to conserve body water. Its kidneys are very efficient and it sweats only when its body temperature exceeds 116° F (46° C). The large hooves spread the stocky body's weight on soft sand. This oryx feeds on a wide variety of plants in the early morning and evening, and on moonlit nights, resting by day in any available shade. It forms nomadic mixed herds

of 20–40, males displaying and tussling to mate with females. After 222–253 days' gestation, the mother leaves the herd to calve, but returns within hours. Young are weaned by 14 weeks and are sexually mature by 2 years. This species was hunted almost to extinction, surviving only on a reserve in north-central Chad. Captive bred animals being held in Tunisia are part of a planned reintroduction program.

