


Bats



During sleep, toes have special locking mechanism to prevent falling.

The bat hangs from a branch with its strong, hooked claws.

Bats can grip many types of surfaces.

Wings are made of skin and bone.

Clawed thumb on the edge of the wing

Mexican fruit bat eats mostly fruit but will also eat insects.



Funnel-eared bat lives in South America and roosts in caves.



Mouse-tailed bat lives in Africa and Asia.



Short-tailed leaf-nosed bat eats bananas in South America.



New World fruit bat helps to spread American tree seeds.



Short-tailed leaf-nosed bat has an acute sense of smell.



Franquet's fruit bat is an African bat that eats ripe fruit.



Yellow-shouldered bat feeds on nectar, pollen, and fruit.



Lesser horseshoe bat feeds exclusively on insects.



Proboscis bat roosts in trees by streams; its fur looks like lichen.



Spear-nosed bat has a well-developed sense of smell.



Common vampire bat has saliva that prevents its host's blood clotting.



New World fruit bat can produce young twice a year.



Borneo fruit bat has a distinct long and fox-like muzzle.



Noctule bat flies over fields and meadows, hunting for large beetles.



White-lined bat often roosts in the trunks of trees.



White line, which helps to disguise bat in foliage.

This species of bat uses its wings just two weeks after birth.

Wings are folded in while the bat hangs upside down on branch.

White-lined bat hanging

This bat has a strong sense of smell to locate fruit to eat. It also eats insects.