


» RED ALGAE

**PTEROCADIELLA CAPILLACEA**

With feathery branches that taper towards the tip, this red seaweed, found in pools worldwide, often develops a Christmas tree-like appearance.

**SPOROLITHON PTYCHOIDES**

This red algae forms a crust using calcareous deposits in its cell walls. It is found worldwide in rock pools and on current-swept rocky seabeds.

**CERAMIUM VIRGATUM**

This small red seaweed grows worldwide on rocks and other seaweeds. From a tiny holdfast it grows into filament-like fronds with forked tips.

**GELIDIUM PUSILLUM**

Growing worldwide from an extensive creeping base that incorporates shells and small snails, this turf-forming red seaweed has flattened, leaf-like fronds.

20 cm
8 inflat, leaf-like
structure**MELANAMANSIA FIMBRIFOLIA**

Found in N. America and Australia, this seaweed grows on sediment-covered reefs at depths of up to 55 m (180 ft).


blades on
frond**CHONDRIA DASYPHYLLA**

This species, found worldwide, has feathery fronds ending in club-like branchlets that sprout spore-bearing branchlets and urn-shaped fruit bodies.

10–21 cm
4–8 1/2 in20 cm
8 in**BROAD WEED***Lenormandiopsis nozawae*

On both sides of this broad-bladed seaweed, found in temperate areas, are clusters of spore-bearing bodies, themselves home to tiny, parasitic algae.

stolon

8.5 cm
3 1/4 in**GELIDIELLA ACEROSA**


An important source of agar, used in the food and pharmaceutical industries, this red seaweed, found in India, has a slender, cylindrical shoot, known as a stolon.

35 cm
14 in**LEATHER WEED***Ptilophora leliaertii*

First described in 2004 after its discovery on a reef off the coast of South Africa, this red seaweed has feathery compound branches.

15 cm
6 in**BUSH WEED***Ahnfeltia* sp.

A source of the gelatinous substance agar, used in petri dishes for cell culture, this northern-hemisphere seaweed forms dense tufts of fronds.

7–22 cm
2 3/4–8 3/4 in**LAURENCIA OBTUSA**

This tropical species is a source of halogenated terpenoids – chemical defences against grazing crabs and urchins – which also make useful anti-fouling agents.

30 cm
12 in**SEA BEECH***Delesseria sanguinea*

Known for its characteristic beech-like leaves, this red seaweed grows in the understory of kelp forests in Europe.

2–10 cm
3/4–4 in**POLYSIPHONIA LANOSA**

Growing on other seaweeds in tufts like pom-poms in the northern hemisphere, this red alga has branching filaments made up of long tubular cells.