




GODETIA
Clarkia amoena
F: Onagraceae


This annual is native to coastal hills of western N. America. Cultivated in gardens, its flower has four broad petals and forms a dry capsule, containing numerous seeds.



FUCHSIA FULGENS
F: Onagraceae
Typically found clinging to rocks or tree branches by its long, tuberous roots, this deciduous shrub grows near water in the mountains of Mexico.




HARDY FUCHSIA
Fuchsia magellanica
F: Onagraceae
This widely planted deciduous, ornamental perennial originates from Chile and Argentina, where it usually grows with its roots in water.



ROSEBAY WILLOWHERB
Epilobium angustifolium
F: Onagraceae
Widespread in the northern hemisphere, this herbaceous perennial proliferates by spreading its roots and wind-dispersed seeds. Some taxonomists put it in the genus *Chamerion* or *Chamaenerion*.

large, purple-pink flower



GREAT WILLOWHERB
Epilobium hirsutum
F: Onagraceae

This very hairy perennial forms extensive patches from its underground stems. It is native to most of Europe, N. Africa, and parts of Asia.



50 cm
20 in

SHOWY EVENING PRIMROSE
Oenothera speciosa
F: Onagraceae

This herbaceous, smooth-stemmed perennial is from S.E. USA and Mexico. Its white flowers become pink with age and close in full sun.



1.5 m
5 ft


EVENING PRIMROSE
Oenothera biennis
F: Onagraceae

This biennial from eastern N. America has bluish green leaves in a basal rosette and erect stems bearing flowers that open in the evening. The seed oil has medicinal properties.



stigma has four lobes

characteristic four petals



1 m
3 1/4 ft

SEEDBOX
Ludwigia alternifolia
F: Onagraceae

Native to eastern N. America, this is a plant of damp places. It has box-shaped seed pods borne on angled stems.

ENCHANTER'S NIGHTSHADE
Circaea lutetiana
F: Onagraceae


This woodland plant is found in Europe, N. Africa, and the Middle East. Flowers of two sepals and two notched petals form small, round, hairy fruits.

CELASTRALES

The two families in this order are characterized by a nectar-secreting disc in the flower. The Lepidobotryaceae comprises just two tropical tree species, but the Celastraceae or spindle tree family has 1,200 species, mainly of vines, shrubs, and trees. Genetic analysis is yet to determine how the Celastrales order relates to other rosids.

SPINDLE TREE
Euonymus europaeus
F: Celastraceae

Native to Europe, the wood of this tree was used to make spindles – spikes used to spin thread. Its fruits ripen to red and split into four to reveal orange-coated, poisonous seeds.



6 m
20 ft

nectar disc

elliptical leaf

GRASS OF PARNASSUS
Parnassia palustris
F: Celastraceae

This rosette plant grows in bogs in the northern temperate zone. Each flower is borne on a stem, clasped by a stalkless leaf.



30 cm
12 in