


OXALIDALES

This order contains about 2,000 species, grouped within seven families. Of these, the Cephalotaceae contains only one species – the carnivorous Albany pitcher plant. The Cunoniaceae are woody plants that produce woody fruit capsules containing small seeds. The wood sorrel family (Oxalidaceae), is the largest with 800 species within five genera. They are characterized by divided leaves that exhibit or open out by day and then close up at night.



ALBANY PITCHER PLANT

Cephalotus follicularis
F: Cephalotaceae

This carnivorous plant is native to the coast of S.W. Australia. It has basal, oval leaves, and a liquid-filled pitcher that traps prey but it is not a member of the pitcher family (see p.194).

20 cm
8 in

flower has
five petals

4 m
13 ft

35 cm
14 in

PINK SORREL

Oxalis articulata
F: Oxalidaceae

Growing from a swollen rhizome, or root stalk, this plant from S. America forms leafy mounds topped with clusters of flowers. These produce seed capsules that explode.

FLOWER
CLUSTER

CHRISTMAS BUSH

Ceratopetalum gummiferum
F: Cunoniaceae

From coastal E. Australia, this shrub produces insignificant white flowers in spring. The pink and red sepals enlarge in winter, enclosing the fruit.

12 m
40 ft

BLACK WATTLE

Callicoma serratifolia
F: Cunoniaceae

The wood of this shrubby tree from the east coast of Australia was used by early settlers to make wattle-and-daub shelters. Its young foliage is bronze.

RIPE
FRUIT

FLOWER
SPRIG

STAR FRUIT

Averrhoa carambola
F: Oxalidaceae

This bushy tree, native to S.E. Asia, is widely cultivated for its edible star-shaped fruits. It can flower four times a year.

15 m/50 ft

LEAFLET
CLUSTER

ROSALES

This order of plants contains nine families, among them the Rosaceae (rose family), Cannabaceae (hemp family), Moraceae (mulberry family), Rhamnaceae (buckthorn family), Ulmaceae (elm family), and Urticaceae (nettle family). Members of the Rosales are often grown for their fruits or other products.

Plants in this order tend to have five sepals and numerous stamens. Mostly insect-pollinated, they are often thorny or hairy.



oil-rich berry

SEA BUCKTHORN

Hippophae rhamnoides
F: Elaeagnaceae

Widespread through Asia and Europe, this shrub bears tiny flowers before the leaves. The small, bright-orange berries are rich in vitamin C.

10 m
33 ft