




LOVE-LIES-BLEEDING
Amaranthus caudatus
F: Amaranthaceae
Believed to originate in S. America, this annual plant has long been cultivated for its edible leaves and seeds.



POLPALA
Aerva lanata
F: Amaranthaceae

With catkin-like flower clusters, this perennial from tropical regions of Asia and Africa grows on open, disturbed ground.



ACHYRANTHOS BIDENTATA
F: Amaranthaceae
Forest edges, streamsides, and moist shady places in China, Japan, India, and Nepal are home to this species.

ANNUAL SEABLITE
Suaeda maritima
F: Amaranthaceae

Mainly coastal, inhabiting salt marshes in Europe, but growing inland in parts of Asia and N. America, annual seablite has green leaves which turn red.



HEART-SHAPED LEAF

TALL, LEAFY STEM

SEA BEET
Beta vulgaris
F: Amaranthaceae

The wild ancestor of beetroots, sea beet is a fleshy inhabitant of bare ground by the sea in parts of Europe, N. Africa, and Asia.



SEA PURSLANE
Atriplex portulacoides
F: Amaranthaceae

Salt marshes, especially edges of tidal channels and pools, are home to sea purslane, a sprawling silvery species from S. Europe and the Mediterranean.



COMMON GLASSWORT
Salicornia europaea
F: Amaranthaceae

Common glasswort is a bushy species of muddy salt marshes in W. Europe. Its succulent stems are sometimes cooked as a vegetable.



MEXICAN TEA
Dysphania ambrosioides
F: Amaranthaceae

A short-lived inhabitant of cultivated and waste ground in tropical America, this aromatic species is used as a seasoning and in tea.



PLUMED COCKSCOMB
Celosia argentea
F: Amaranthaceae

This showy plant is native to dry slopes and stony ground in tropical Africa, but is widely planted elsewhere for its colourful foliage.



GARDEN ORACHE
Atriplex hortensis
F: Amaranthaceae

Probably originally from seashores in S.W. Asia, this spinach-like species has long been cultivated for its edible leaves.



EDIBLE SALAD LEAF

1.2 m
4 ft

2 m
6 1/2 ft