




sheath
around
flower

50 cm
20 in

CHILDING PINK
Petrorhagia nanteuilii
F: Caryophyllaceae


The flowers of childing pink open one at a time. This species inhabits dry grassy places on sandy soils in W. Europe.



1 m
3 1/4 ft

SOAPWORT
Saponaria officinalis
F: Caryophyllaceae


Once used to make soap, this species is found along streams and damp ground in S. Europe and W. Asia.



50 cm
20 in

COMMON CHICKWEED
Stellaria media
F: Caryophyllaceae

This sprawling species is found on cultivated and open ground worldwide and is sometimes used as a salad vegetable.



ROUND-LEAVED SUNDEW
Drosera rotundifolia
F: Droseraceae

Inhabiting open boggy moorland and heathland in Europe, N. Asia, and N. America, this insectivorous species has leaves with numerous sticky hairs covered in insect-dissolving enzymes.

leaf hairs
trap insects

10 cm
4 in



white flowers
surrounded by bracts


brightly coloured
bract

trap triggered by hairs
on leaf surface

8 m/26 ft

BOUGAINVILLEA
Bougainvillea glabra
F: Nyctaginaceae

What seem to be petals on bougainvillea are actually bracts. This slow-growing evergreen climber from Brazil is widely cultivated.



when triggered,
leaf snaps shut,
trapping insect

"teeth" prevent
insect from escaping

15 cm
6 in

nectar glands on
lid attract insects



30 cm/12 in

SEA HEATH
Frankenia laevis
F: Frankeniaceae

A mat-forming species, sea heath grows on bare sandy ground in the drier parts of salt marshes in W. Europe and N.W. Africa.



FOUR O'CLOCK FLOWER
Mirabilis jalapa
F: Nyctaginaceae

From dry open habitats in tropical C. and S. America, this species has fragrant flowers which open during the late afternoon.

1 m/3 1/4 ft



insects slither
off slippery rim
into pitcher

fluid in base of pitcher
digests insects

pitcher
grows from
tendrils at
end of leaf

12 m
40 ft

PITCHER PLANT
Nepenthes stenophylla
F: Nepenthaceae

In the forests of Borneo, this scrambling vine gets extra nutrients by trapping insects in its pitchers.