



LEWISIA BRACHYCALYX
F: Montiaceae

Native to damp stony meadows in mountainous areas of S.W. USA, this species has a basal rosette of fleshy leaves.

4 cm
1½ in



SPRING BEAUTY
Claytonia perfoliata
F: Montiaceae

Also known as miner's lettuce, this native of cultivated and bare ground in western N. America, Mexico, and Guatemala has two fused leaves beneath the flowers.

30 cm
12 in

FRINGED REDMAIDS
Calandrinia ciliata
F: Montiaceae

Native to grasslands from S. Canada to Argentina, this species has escaped to the Falkland Islands and is also grown in gardens.

30 cm
12 in



CEYLON LEADWORT
Plumbago zeylanica
F: Plumbaginaceae

Growing mainly in open, human-created habitats, this scrambling shrub is found throughout tropical and subtropical regions.



1.5 m
5 ft



25 cm
10 in

SEA PINK
Armeria maritima
F: Plumbaginaceae

In much of W. Europe, thrift, or sea pink, inhabits coastal rocks and cliffs, salt marshes, and mountains. It is a cushion-forming species.



3 m
10 ft

TAMARISK
Tamarix gallica
F: Tamaricaceae


Usually coastal, tamarisk also grows in saline areas inland. It originates from S. Europe, N. Africa, and the Canary Islands, and is planted elsewhere.



2 m
6½ ft

JOJOBA
Simmondsia chinensis
F: Simmondsiaceae

Deserts in Arizona and California, USA, and Mexico are home to jojoba, a species often cultivated for its oil.



40 cm
16 in

STATICE
Limonium sinuatum
F: Plumbaginaceae

Statice has winged branching stems and is found in rocky and sandy coastal habitats and in saline areas inland in Mediterranean countries.

SANTALES

Found mainly in the tropics and subtropics, the seven families in the order Santales include several important timber trees. They also feature many parasitic and semi-parasitic plants, notably the 900 species in the showy mistletoe family (Loranthaceae) of the southern hemisphere. These live attached to other plants, from which they obtain all or most of the water and nutrients they need for growth.

10 m/33 ft



FIRE TREE
Nuytsia floribunda
F: Loranthaceae

This species is semi-parasitic, deriving moisture and nutrients from the roots of surrounding plants in woodland in S.W. Australia.



berry can be
poisonous

1 m
3¼ ft

9 m
30 ft



SANDALWOOD
Santalum album
F: Santalaceae

Cultivated for its wood and fragrant oil, this semi-parasitic tree grows in dry, rocky areas in Asia and Australia.

MISTLETOE
Viscum album
F: Santalaceae

Mistletoe forms spherical masses on tree branches. A semi-parasitic species with white berries, it inhabits much of Europe, N. Africa, and Asia.

1.2 m
4 ft

OSYRIS
Osyris alba
F: Santalaceae

This broom-like semi-parasitic species with fragrant flowers is native to S. Europe, N. Africa, and S.W. Asia, inhabiting dry, rocky places.

