

YELLOW RATTLE
Rhinanthus minor
F: Orobanchaceae

A partial parasite whose roots tap grasses for nutrition, this variable, yellow-flowered annual grows in grasslands in northern temperate regions.


50 cm
20 in



PURPLE TOOTHWORT
Lathraea clandestina
F: Orobanchaceae

Lacking green leaves, this parasitic W. European perennial taps the roots of willows and poplars for food. Its flowerheads emerge directly from underground shoots.

8 cm
3 in



flowers attract
butterflies for
pollination



opposite, oval
shiny leaf

3.7 m
12 ft

FLAMING GLORYBOWER
Clerodendrum splendens
F: Lamiaceae

This African vine, with clusters of tubular, red flowers, twines round trees in its native forest habitat, or trellises in gardens.

**BODINIER'S
BEAUTYBERRY**
Callicarpa bodinieri
F: Lamiaceae

Related to American beautyberry, this Chinese ornamental shrub is grown in gardens for its decorative purple berries, which are bitter but not poisonous.

3 m
10 ft



VERVAIN
Verbena officinalis
F: Verbenaceae

Widely distributed in rough grassland in temperate and tropical regions, this stiffly hairy perennial has slender spikes of two-lipped, lilac flowers.


1 m
3 1/4 ft



LANTANA
Lantana camara
F: Verbenaceae

The tubular flowers of this prickly shrub open yellow or orange then turn red. Native to C. and S. America, it is one of the most invasive weeds in warm regions.

1.5 m
5 ft




SOLANALES

The potato family (Solanaceae), an economically important family of up to 4,000 species, dominates the order Solanales. Many species contain poisonous alkaloids. The bindweed family (Convolvulaceae) includes tropical climbers and low-growing herbs. The three other families in this order are the Hydrolaceae, with one genus found only in the Americas; five African trees in the Montiniaceae; and a pan-tropical herb in the Sphenocleaceae.

GREAT BINDWEED
Calystegia sylvaticus
F: Convolvulaceae

A plant of hedgerows and waste ground, even in cities, around the temperate northern hemisphere, this large-flowered perennial spreads by far-creeping rhizomes (underground plant stems).

3 m
10 ft



trumpet-shaped
flower with
white or
yellow centre

MORNING GLORY
Ipomoea tricolor
F: Convolvulaceae

This twining herbaceous climber from Central America has lobed leaves and funnel-shaped flowers that open only in the morning.



three-
lobed leaf

4 m
13 ft



COMMON DODDER
Cuscuta epithymum
F: Convolvulaceae

A Eurasian annual forming a dense network of threadlike, twining stems, parasitic dodder taps into gorse, heather, and other species for food.

60 cm
23 1/2 in