

AGARICALES

Many of the most familiar mushrooms and toadstools belong to this order. They include fungi with fleshy, not woody, fruitbodies (the body supporting the spore-forming cells). Many have caps and stems with gills; some also have pores. Other forms include bird's-nest, brackets, crusts, truffle-like, and puffballs. Most live on leaf litter, soil, or wood; others are parasitic or live in association with plant roots (mycorrhizal).



5–12 cm
2–4 3/4 in

YELLOW STAINER
Agaricus xanthodermus
F: Agaricaceae

This species has a flattened cap centre, chrome-yellow stains on the stem base, and an unpleasant odour. It is common in Eurasia and western N. America.



3–8 cm
1 1/4–3 1/4 in

BLUSHING DAPPERLING
Leucoagaricus badhamii
F: Agaricaceae

Found in Eurasia, this rare fungus grows in woods and gardens with rich soils. White when young, it bruises blood-red when handled, and then blackens.



5–8 cm
2–3 1/4 in

WHITE DAPPERLING
Leucoagaricus leucothites
F: Agaricaceae


Common in meadows and grassy roadsides in Eurasia and N. America, this fungus's ivory fruitbody turns grey-brown with age. Its gills are white to pale pink.



1–5 cm
3/8–2 in

PLANTPOT DAPPERLING
Leucocoprinus birnbaumii
F: Agaricaceae

Common worldwide, this species grows on soils of potted plants. Its features include a delicate golden-yellow cap and a slender stem with a ring.



YOUNG CAP

LAWYER'S WIG
Coprinus comatus
F: Agaricaceae

This fungus is common on disturbed soils and pathsides in Eurasia and N. America. Its tall, shaggy cap is unmistakable, and dissolves into inky fluid.




MATURE CAP

2–6 cm
3/4–2 1/4 in

THE PRINCE
Agaricus augustus
F: Agaricaceae

A large, orange-brown, scaly cap and a white, woolly stem with a floppy ring characterize this common species from Eurasia and N. America.



4–10 cm
1 1/2–4 in



8–15 cm
3 1/4–6 in

FIELD MUSHROOM
Agaricus campestris
F: Agaricaceae

Common in meadows of Eurasia and N. America, this fungus has a rounded cap; bright pink gills turning brown with maturity; and a short stem with a small ring.




5–10 cm
2–4 in

CULTIVATED MUSHROOM
Agaricus bisporus
F: Agaricaceae

Selling in millions, this familiar fungus is common worldwide. Its cap varies from white to dark brown with scales.

FRECKLED DAPPERLING
Echinoderma asperum
F: Agaricaceae

From woods and gardens in Eurasia and N. America, this uncommon fungus's brown cap has pyramidal scales that rub off. The brown stem has a ring.



5–15 cm
2–6 in

HORSE MUSHROOM
Agaricus arvensis
F: Agaricaceae

Common in parks in Eurasia and N. America, this species stains dull brassy yellow. The pendent ring on its stem has a cogwheel-like pattern on its underside.




7–15 cm
2 1/4–6 in



1–4 cm
3/8–1 1/2 in

STINKING DAPPERLING
Lepiota cristata
F: Agaricaceae

A rubber-like smell is a characteristic feature of this species of *Lepiota* (most common in woods and meadows) from Eurasia and N. America.



4–11 cm
1 1/2–4 1/2 in

ORANGE-GIRDLED DAPPERLING
Lepiota ignivolva
F: Agaricaceae

Found in Eurasia, this uncommon species has an orange-margined ring set low down on the club-shaped, white stem.



INNER LAYER

GREY PUFFBALL
Bovista plumbea
F: Agaricaceae

This ball-shaped fungus is smooth when young. As it matures, its skin flakes to reveal a papery inner layer. It is common in Eurasia and N. America.




OUTER SKIN



GIANT PUFFBALL
Calvatia gigantea
F: Agaricaceae

Common in hedgerows, fields, and gardens in Eurasia and N. America, this mushroom's large, smooth, white fruitbody with a white to yellow interior is unmistakable.



5–15 cm
2–6 in

SHAGGY PARASOL
Chlorophyllum rhacodes
F: Agaricaceae

Common in Eurasia and N. America, this fungus has a shaggy, brown cap; thick, double-edged ring; reddening flesh; and swollen stem base.

bulbous stem base