



contrasting
brown scales


10–30 cm
4–12 in

PARASOL

Macrolepiota procera
F: Agaricaceae

Common in meadows in Eurasia and N. America, this tall species has a scaly cap and snakeskin-patterned stem with a thick ring.

snakeskin
pattern on
the stem



1–2.5 cm
 $\frac{3}{8}$ –1 in

STUMP PUFFBALL

Apioperdon pyriforme
F: Agaricaceae

Common in Eurasia and N. America, this pear-shaped species grows on wood. It has prominent cords at its base, and is firm when young.



6–10 cm
2 $\frac{1}{4}$ –4 in

WINTER STALKBALL

Tulostoma brumale
F: Agaricaceae

Typically found in sandy soils in dune slacks in Eurasia and N. America, this fungus has tiny, whitish yellow, round heads on slender, pale brown stalks.



3–5 cm
1 $\frac{1}{4}$ –2 in

SANDY STILTBALL

Battarreoides digueti
F: Tulostomataceae

Found in very dry, sandy soils in N. America, this tall, stalked species emerges from a leathery "egg". Its brown cap contains the spores.



6–15 cm
2 $\frac{1}{4}$ –6 in

POPLAR FIELDCAP

Cyclocybe cylindracea
F: Bolbitiaceae

Found in Eurasia, this rare species grows on poplar wood. The cap cracks when dry and the stem has a ring.



1–3 cm
 $\frac{3}{8}$ –1 $\frac{1}{4}$ in

SPINY PUFFBALL

Lycoperdon echinatum
F: Agaricaceae

This species is found in beech woods in Eurasia and N. America. It has groups of long spines that meet at their tips. Its spores are purplish brown.




1–3.5 cm
 $\frac{3}{8}$ –1 $\frac{1}{2}$ in

COMMON FIELDCAP

Agrocybe pediades
F: Bolbitiaceae

Usually found in lawns in Eurasia, this fungus's smooth, yellowish cap and slender stem have a mealy smell and no veil.



1–1.5 cm
 $\frac{3}{8}$ – $\frac{1}{2}$ in

MILKY CONECAP

Conocybe apala
F: Bolbitiaceae

This species is frequently found in lawns in Eurasia and N. America. Its conical, ivory cap with orange-brown gills sits on a tall stem.



2–4.5 cm
 $\frac{3}{4}$ –1 $\frac{3}{4}$ in

MEADOW PUFFBALL

Lycoperdon pratense
F: Agaricaceae

This short-stemmed species has an internal membrane separating the spore-mass from the stem. It is common in meadows throughout Eurasia.


5–10 cm
2–4 in



PESTLE PUFFBALL

Lycoperdon excipuliforme
F: Agaricaceae

One of the tallest puffballs, this fungus has a buff-white stem that is left behind once the spores are released. It is found in Eurasia.




2–4 cm
 $\frac{3}{4}$ –1 $\frac{1}{2}$ in

COMMON PUFFBALL

Lycoperdon perlatum
F: Agaricaceae

The most common white puffball, this species is found in Eurasia and N. America. Its granular spines rub off to leave circular scars in a regular pattern.



2–5 cm
 $\frac{3}{4}$ –2 in

STINKING PUFFBALL

Lycoperdon foetidum
F: Agaricaceae

Common in acidic heaths and woods in Eurasia, this yellowish brown puffball has dark brown spines. Its flesh has an unpleasant smell.



3–7 cm
1 $\frac{1}{4}$ –2 $\frac{3}{4}$ in

SPRING FIELDCAP

Agrocybe praecox
F: Bolbitiaceae

Common in Eurasia and N. America during spring, this fungus's cap may have marginal veil fragments. The stem may have a fragile ring.



1–3 cm
 $\frac{3}{8}$ –1 $\frac{1}{4}$ in

YELLOW FIELDCAP

Bolbitis titubans
F: Bolbitiaceae

Growing in meadows, this fungus is common in Eurasia and N. America. It has a delicate, sticky cap that lasts only a day or so.