



» AGARICALES




CAESAR'S MUSHROOM
Amanita caesarea
F: Amanitaceae

Found under oaks, mainly from the Mediterranean to C. Eurasia, this mushroom has an orange cap and stem, and a white volva.




DEATH CAP
Amanita phalloides
F: Amanitaceae

Common in Eurasia and parts of N. America, this species has a fragile ring, large volva, and sickly smell. Its cap may be greenish, yellowish, or white.



FALSE DEATHCAP
Amanita citrina
F: Amanitaceae

Found in Eurasia and eastern N. America, this species has a stem with a distinctly rimmed, rounded bulb, and flesh with a potato odour. Its cap is white or pale yellow.



TAWNY GRISETTE
Amanita fulva
F: Amanitaceae

Common in woods in Eurasia and N. America, this fungus may have veil patches. The stem has a white volva (bag-like remains of an enclosing veil).




BLUSHER
Amanita rubescens
F: Amanitaceae

This species is common in mixed woods in Eurasia and N. America. Its cap varies from cream to brown, and its flesh bruises pinkish red.



FLY AGARIC
Amanita muscaria
F: Amanitaceae

Common in Eurasia and N. America, this distinctive mushroom is especially common under birches. Its white warts may wash off after rain.



DESTROYING ANGEL
Amanita virosa
F: Amanitaceae

Found throughout N. Eurasia, this rare fungus is pure white with a slightly sticky, bell-shaped cap and a white volva.



PANTHER CAP
Amanita pantherina
F: Amanitaceae

This species has pure white warts (remains of its protective veil), a ring on its stem, and a ridged bulb. It is found in Eurasia and N. America.



GOLDEN SPINDLES
Clavulinopsis corniculata
F: Clavariaceae

This golden-yellow club with antler-like branches is relatively common in acidic, unimproved meadows and grassy clearings in woods in Eurasia.

YELLOW CLUB
Clavulinopsis helvola
F: Clavariaceae

Common in grassland in Eurasia and N. America, this is one of several yellow clubs. It can only be reliably identified by its spiny spores.



club often flattened



VIOLET CORAL
Clavaria zollingeri
F: Clavariaceae

A rare species found in mossy meadows and woodlands in Eurasia and N. America, this species has violet, coral-like branches.



WHITE SPINDLES
Clavaria fragilis
F: Clavariaceae

This fungus forms dense clusters of pure white, simple clubs in woods and meadows in Eurasia.



WOOD PINKGILL
Entoloma rhodopolium
F: Entolomataceae

Found in Eurasia, this pale grey to grey-brown species has a cap with a domed centre and slender stem. It may smell of bleach.




MOUSEPREE PINKGILL
Entoloma incanum
F: Entolomataceae

Smelling of mice, this pinkgill has a hollow stem and a bright grass-green cap that fades to brown with age. It grows in unimproved meadows in Eurasia.



BLUE EDGE PINKGILL
Entoloma serrulatum
F: Entolomataceae

This blue-black capped fungus has serrated pinkish gills with dark edges. It is found in meadows and parks in Eurasia and N. America.



LILAC PINKGILL
Entoloma porphyrophaeum
F: Entolomataceae

Occasionally found in meadows, this fungus grows in Eurasia. It has a grey-purple fibrous cap and stem, and pinkish gills.



THE MILLER
Clitopilus prunulus
F: Entolomataceae

Common in mixed woods in Eurasia and N. America, this fungus has a cap that changes shape with age, starting convex then becoming depressed.