



5–20 cm
2–8 in

contrasting
cream gills

5–20 cm
2–8 in

WOOD BLEWIT
Lepista nuda
F: Tricholomataceae

Common in mixed woods in Eurasia and N. America, this species has a violet cap that fades to brown, but the stem and gills remain violet.

BLUE LEGS
Lepista personata
F: Tricholomataceae
The blue legs has a buff cap and a bright blue-lilac stem with perfumed flesh. It is common in open meadows in Eurasia.



8–20 cm
3¼–8 in

CLOUDED FUNNEL
Clitocybe nebularis
F: Tricholomataceae

This fleshy mushroom has crowded gills that often run down the stem. It frequently grows in large circles in Eurasia and N. America.



3–6 cm
1¼–2¼ in

ANISEED FUNNEL
Clitocybe odora
F: Tricholomataceae

Common in Eurasia and N. America, this fungus is unmistakable due to its strong aniseed odour. Its sea-green colour fades as it matures to greyish green.



2–6 cm
¾–2¼ in

IVORY FUNNEL
Clitocybe dealbata
F: Tricholomataceae

This species grows in rings in lawns and meadows in Eurasia and N. America. It has a frosted appearance and slightly decurrent gills.



2–8 cm
¾–3¼ in

SULPHUR KNIGHT
Tricholoma sulphureum
F: Tricholomataceae

Characterized by its bright sulphur-yellow colour and repulsive smell of coal gas, this fungus is common in mixed woods in Eurasia and parts of N. America.




inrolled cap (curls
inward at margins)

12–40 cm
4¾–16 in

short stem

GIANT FUNNEL
Leucopaxillus giganteus
F: Tricholomataceae
Often found in huge fairy rings in Eurasia and N. America, this giant fungus is white, with decurrent gills, an inrolled cap margin, and a short stem.



5–12 cm
2–4¾ in

DINGY AGARIC
Tricholoma portentosum
F: Tricholomataceae

Usually found with conifers in northern N. America and N. Eurasia, this species has a smooth, bell-shaped cap and a stout, whitish stem that flushes yellowish.



4–10 cm
1½–4 in

SOAPY KNIGHT
Tricholoma saponaceum
F: Tricholomataceae

Variable in colour – grey-brown, greenish or pinkish grey to mottled – with a soapy smell, this species is common in mixed woods in Eurasia and N. America.

PIPE CLUB
Typhula fistulosa
F: Typhulaceae

Often found in large groups in Eurasia and parts of N. America, this species has tall, slender clubs attached to fallen branches and hardwood litter.

5–25 cm/2–10 in