


» AGARICALES



5–12 cm
2–4¾ in

THE GYPSY
Cortinarius caperatus
F: Cortinariaceae


From N. Eurasia and N. America, this species, found in conifer woods, has a frosted cap with a white veil, and a stem with a distinct sheathing ring.



3–6 cm
1¼–2¼ in

DAPPLED WEBCAP
Cortinarius bolaris
F: Cortinariaceae

Frequent in beech woods in Eurasia, this fungus has tiny reddish brown scales on its cap and stem. The flesh stains yellow-orange in the stem.



5–8 cm
2–3¼ in

BLUE AND BUFF WEBCAP
Cortinarius malachius
F: Cortinariaceae

Found under pines in N. Eurasia, this uncommon species has a pale brown cap with a lilac tint, and a stem with bands of white veil.



6–10 cm
2¼–4 in

ELEGANT WEBCAP
Cortinarius elegantissimus
F: Cortinariaceae

Found throughout Eurasia in beech woods on chalky soil, this fungus has a vivid orange-yellow cap and a yellow stem with a large, marginate bulb.



5–10 cm
2–4 in

BLUE AND BUFF WEBCAP
Cortinarius malachius
F: Cortinariaceae

Found under pines in N. Eurasia, this uncommon species has a pale brown cap with a lilac tint, and a stem with bands of white veil.



3–8 cm
1¼–3¼ in

SCALY WEBCAP
Cortinarius pholideus
F: Cortinariaceae

This uncommon species is found under birches in Eurasia. It is easily recognized by its scaly, brown cap and stem, and violet gills and stem apex when young.

raised point at
cap centre

scaly ring

6–10 cm
2¼–4 in

PEARLY WEBCAP
Cortinarius alboviolaceus
F: Cortinariaceae

This species is common in mixed woods in Eurasia and N. America. Its silvery white fruitbody has a lilac tint. The gills mature to cinnamon-brown.



5–12 cm
2–4¾ in

RED AND OLIVE WEBCAP
Cortinarius rufolivaceus
F: Cortinariaceae

Found in beech woods throughout Eurasia, this uncommon species has a distinctively copper-coloured cap – with a pink or olive-green margin. The bulbous stem is violet to greenish yellow.

ORANGE WEBCAP
Cortinarius mucosus
F: Cortinariaceae

Restricted to N. Eurasia and N. America, this species is common in pine woods. Its orange-brown cap and stout, white stem are very slimy. The gills are rust-brown.



1–3 cm
¾–1¼ in



4–10 cm
1½–4 in

**BITTER
BIGFOOT WEBCAP**
Cortinarius sodagnitus
F: Cortinariaceae

Found mainly in S. England and Mediterranean Eurasia, this uncommon species occurs in beech woods on chalky soil. The bulb on the stem is large.



3–7 cm
1¼–2¾ in


SPLENDID WEBCAP
Cortinarius splendens
F: Cortinariaceae

This rare species has a golden-yellow cap, yellow flesh, a sulphur-yellow veil, and a bulbous stem. It mainly occurs in beech woods in Eurasia.



PIXIE WEBCAP
Cortinarius flexipes
F: Cortinariaceae

An odour of pelargoniums and delicate white hairs on the pointed cap characterize this species. It is common under birches in N. Eurasia and N. America.



minute
scales
on cap

ring zone

club-shaped
stem

6–15 cm
2¼–6 in


VIOLET WEBCAP
Cortinarius violaceus
F: Cortinariaceae

Found in mixed woodland throughout Eurasia and N. America, this rare species has a distinctive intensely violet-coloured cap and stem.

RED BANDED WEBCAP
Cortinarius armillatus
F: Cortinariaceae

The cinnabar-red bands of veil on the club-shaped stem distinguish this common species of birch woods in Eurasia and N. America.

club-shaped stem



8–15 cm
3¼–6 in

BIRCH WEBCAP
Cortinarius triumphans
F: Cortinariaceae

Common under birches in Eurasia, this fungus is yellow-orange overall, with prominent girdles of yellow veil on the stem.