

CAP (UPPER SIDE)

2–7 cm
¾–2¾ in

CAP (UNDERSIDE)

PEELING OYSTERLING*Crepidotus mollis*
F: Crepidotaceae

This fungus occurs in Eurasia and N. America. Its small, fan-shaped, pale cap has a gelatinous skin, which peels easily. The stem is absent or very short.

VARIABLE OYSTERLING
Crepidotus variabilis
F: Crepidotaceae

One of several similar species in Eurasia and N. America, its cap is dry with fine fibres. It usually lacks a stem.

0.5–3 cm
3/16–1¼ in**ROOTING POISONPIE**
Hebeloma radicosum
F: Hymenogastraceae

The deeply rooting stem with a large ring, buff cap with flat scales, and strong smell of marzipan characterize this Eurasian species.

0.3–0.8 cm
¼–¾ in

CAP NOT FULLY EXPANDED

5–12 cm
2–4¾ in

attached to wood

DEADLY GALERINA
Galerina marginata
F: Hymenogastraceae

This species has an orange-brown cap and a small ring on the stem. It grows on fallen timber in Eurasia and N. America.

1–5 cm
¾–2 in


red band

3–7 cm
1¼–2¾ in**STAR FIBRECAP**
Inocybe asterospora
F: Inocybaceae


A flattened bulb at the stem's base and star-shaped spores distinguish this small, brown fibrous-capped species from Eurasia.

1–4.5 cm
¾–1¾ in**TORN FIBRECAP**
Inocybe lacera
F: Inocybaceae

Found in Eurasia and N. America, this fungus is distinguished by its cylindrical spores. It has a scaly fibrous cap and a slender brown stem.

4–9 cm
1½–3½ in**POISONPIE**
Hebeloma crustuliniforme
F: Hymenogastraceae

From Eurasia and N. America, this species strongly smells of radish. Its cap is ivory-white to buff, and slimy when wet. In damp weather, the gills exude droplets.



radial fibres

stout stem

3–9 cm
1¼–3½ in**DEADLY FIBRECAP**
Inocybe erubescens
F: Inocybaceae

Found in mixed woodland on chalk in Eurasia, this uncommon species has a fibrous cap that discolours with age, and stout stem that bruises red.

3–7 cm
1¼–2¾ in**SPLIT FIBRECAP**
Inocybe rimosa
F: Inocybaceae

Common in mixed woodland in Eurasia and N. America, its pointed fibrous cap is straw-yellow and the stem is tall and slender.



WHITE FORM

1–4 cm
¾–1½ in**WHITE FIBRECAP**
Inocybe geophylla
F: Inocybaceae

This is one of the commonest *Inocybe* species in woods in Eurasia and N. America. It has a conical, silky cap and a slender stem.

0.8–4 cm
5/16–1½ in**LILAC LEG FIBRECAP**
Inocybe griseolilacina
F: Inocybaceae

A common species of beech woods in Eurasia, it has a scaly cap and a pale violet, slender stem.

VIOLET FORM