

» AGARICALES



gills are
broadly
attached

2.5–6 cm
1–2¼ in

MEADOW WAXCAP
Cuphophyllus pratensis
F: Hygrophoraceae

One of the larger meadow waxcaps, it is found in Eurasia and N. America. It has a stout stem and a fleshy cap with gills running down the stem.



1.5–5 cm
½–2 in

SNOWY WAXCAP
Cuphophyllus virgineus
F: Hygrophoraceae

The most common waxcap in all types of grassland in Eurasia, its waxy cap and slender stem are translucent. It has decurrent gills.



1–4 cm
¾–1½ in

green stem apex

PARROT WAXCAP
Gliophorus psittacinus
F: Hygrophoraceae

This fungus occurs in Eurasia and N. America. Its slimy cap is vivid green to orange and the upper part of the stem is bright green.




fleshy, waxy,
pink gills

3–7 cm
1¼–2¾ in

PINK WAXCAP
Porpolomopsis calyptriformis
F: Hygrophoraceae

This highly recognizable but rare fungus has a pointed pink cap, fragile pale stem, and waxy gills. Found in Eurasia, it grows in unimproved meadows.



1–5 cm
¾–2 in

BLACKENING WAXCAP
Hygrocybe conica
F: Hygrophoraceae

Quite common in grasslands and woodlands in Eurasia and N. America, this species has a conical reddish orange cap and fibrous stem that blacken with age or bruising.



1.5–6 cm
½–2¼ in

SCARLET WAXCAP
Hygrocybe coccinea
F: Hygrophoraceae

The scarlet, waxy cap, gills, and stem make this a very striking fungus of unimproved grasslands in Eurasia and N. America.



4–12 cm
1½–4¾ in

pale yellow
margin

CRIMSON WAXCAP
Hygrocybe punicea
F: Hygrophoraceae

Found in unimproved grassland in Eurasia and N. America, this uncommon species is the largest blood-red waxcap. Its stem is dry and fibrous with a white base.



1.5–7 cm
½–2¾ in

GOLDEN WAXCAP
Hygrocybe chlorophana
F: Hygrophoraceae


Found in Eurasia, this is the commonest waxcap found in meadows. It has a bright yellow-orange, slightly sticky cap.



4–8 cm
1½–3¼ in

CLUB FOOT
Ampulloclitocybe clavipes
F: Hygrophoraceae

Common in mixed woods in late autumn, this species from Eurasia and N. America has decurrent gills and a swollen, spongy base.



3–8 cm
1¼–3¼ in

IVORY WOODWAX
Hygrophorus eburneus
F: Hygrophoraceae

This species is found in beech woods in Eurasia and N. America. It has a sticky cap and stem, thick gills, and a flowery smell.



2–5 cm
¾–2 in

**AMETHYST
DECEIVER**
Laccaria amethystina
F: Hydnangiaceae


This highly recognizable, slender fungus from Eurasia and N. America has an intense violet coloration when fresh, and powdery spores dusting the gills.



1–5 cm
¾–2 in

DECEIVER
Laccaria laccata
F: Hydnangiaceae

Variable in colour – from brick-red to flesh-pink – with a dry cap and thick gills, this species is abundant in temperate woods in Eurasia and N. America.



0.5–2 cm
3/16–¾ in

CONIFERCONE CAP
Baeospora myosura
F: Cyphellaceae

Found in Eurasia and N. America, this is one of a few fungi growing mostly on pine cones. It has crowded, narrow gills and a velvety stem.

pine cone



3–5 cm
1¼–2 in

**HERALD OF
WINTER**
Hygrophorus hypothejus
F: Hygrophoraceae

Appearing in pine woods after frost in Eurasia and N. America, this sticky fungus has an olive-brown cap and yellow stem.