


» AGARICALES

SILVERLEAF FUNGUS
Chondrostereum purpureum
F: Cyphellaceae

Common on cherry and plum trees, this fungus is found in Eurasia and N. America. Violet below when young, it darkens to purple-brown with age.

**ORANGE BONNET**
Mycena acicula
F: Mycenaceae

Common in leaf litter and debris in broadleaf woods in Eurasia and N. America, this fungus has a tiny, translucent cap that is striped almost to the centre.

**YELLOWLEG BONNET**
Mycena epipterygia
F: Mycenaceae

Occurring in Eurasia and N. America, this fungus grows on acidic soils in woods and heaths. Its cap and stem have a sticky, peelable layer.

**CLUSTERED BONNET**
Mycena inclinata
F: Mycenaceae


Found in dense clusters on wood in Eurasia and N. America, this fungus has a cap with toothed margins and a strong soapy smell.

**OLIVE OYSTERLING**
Sarcomyxa serotina
F: Mycenaceae

From Eurasia and N. America, this mushroom fruits in winter, often near water on hardwood tree trunks. Its cap is slimy when wet.

**SPINDLE TOUGHSHANK**
Gymnopus fusipes
F: Omphalotaceae

Abundant in Eurasian oak woods from early summer onwards, this fungus forms large clumps of tough fruitbodies on tree roots.

**WOOD WOOLLYFOOT**
Gymnopus peronatus
F: Omphalotaceae

The common name of this species from Eurasia refers to the stiff, fuzzy hairs at the base of the stem.

**SPOTTED TOUGHSHANK**
Rhodocollybia maculata
F: Omphalotaceae

Common in mixed woods across Eurasia and N. America, this fungus has a white cap, stem, and gills. The crowded gills stain rust-red with age.



3–6 cm
1¼–2¼ in

**BUTTER CAP**
Rhodocollybia butyracea
F: Omphalotaceae

This fungus is abundant in woods in Eurasia and N. America. Varying from blackish or reddish brown to dark ochre, its cap is greasy to the touch.

**JACK O'LANTERN**
Omphalotus illudens
F: Omphalotaceae

Found across Eurasia and N. America, this bright orange, poisonous fungus is famous for gills that glow in the dark with an eerie greenish light.

**SAFFRONDROP BONNET**
Mycena crocata
F: Mycenaceae

Common in woods with chalky soils in Eurasia, this species exudes a bright saffron-orange juice when it is broken.

**COMMON BONNET**
Mycena galericulata
F: Mycenaceae


Abundant in temperate woods in Eurasia and N. America, this fungus is variable in colour. Its pinkish grey gills have linking veins and cross-ridges.



domed
cap centre

ROSY BONNET
Mycena rosea
F: Mycenaceae

Found in beech woods in Eurasia, this common species has a robust pink cap and stem, and smells strongly of radish.



club-shaped
stem