




3–10 cm
1¼–4 in

HONEY FUNGUS

Armillaria lutea

F: Physalacriaceae


Found in woods across Eurasia, this species is usually found on the ground. It is a weak parasite on the trees around which it grows.

BRANCHING OYSTER

Pleurotus cornucopiae

F: Pleurotaceae

Growing in clusters on fallen logs, usually of elm, this fungus has trumpet-shaped caps with gills running down the short, frequently branching stems. It occurs in Eurasia and N. America.




2.5–10 cm
1–4 in

WRINKLED PEACH

Rhodotus palmatus

F: Physalacriaceae

Found on fallen logs of mainly elm, this uncommon species of Eurasia and N. America has an unusual, peach-pink, wrinkled cap and a fruity odour.




1–6 cm
¾–2¼ in

VELVET SHANK

Flammulina velutipes


F: Physalacriaceae

Growing throughout winter in Eurasia and N. America, this fungus is characterized by its often sticky cap and velvety stem.



4–12 cm
1½–4¾ in

tiny glistening granules



2–3 cm
¾–1¼ in

GLISTENING INKCAP

Coprinellus micaceus

F: Psathyrellaceae

Usually found in clusters, this mushroom's rounded, grooved caps are dusted with mica-like flakes of veil. It is common on wood in Eurasia and N. America.



1.5–7 cm
½–2¾ in

PALE BRITTLESTEM

Psathyrella candolleana

F: Psathyrellaceae

Usually found in clusters on woody debris in early summer, this fungus has fragile, slender stems and pale or buff caps. It occurs in Eurasia and N. America.



0.8–4 cm
⅓–1½ in

CLUSTERED BRITTLESTEM

Psathyrella multipedata

F: Psathyrellaceae

The densely clustered stems of this species are joined at the base. It is usually found in open grass across Eurasia.



2–15 cm
¾–6 in

PORCELAIN FUNGUS

Mucidula mucida

F: Physalacriaceae

Usually found on beech logs in Eurasia, this fungus has grey-white caps that are slimy when wet. The tough stem has a thin ring.



2.5–10 cm
1–4 in

ROOTING SHANK

Hymenopellis radicata

F: Physalacriaceae

Found across Eurasia and N. America, this deep-rooted fungus has a stiff, tall stem. The cap is slimy when wet, and the gills are widely spaced.

caps often overlap



6–20 cm
2¼–8 in

OYSTER MUSHROOM

Pleurotus ostreatus

F: Pleurotaceae

From Eurasia and N. America, this species is found on trees and logs. Its shelf-like caps vary from blue-green to pale buff. The stem is almost absent.



0.5–1 cm
⅓–⅝ in

FAIRIES' BONNETS

Coprinellus disseminatus

F: Psathyrellaceae

Often found in clusters on rotted stumps in Eurasia and N. America, it has tiny umbrella-like caps with deeply fluted grooves. The gills are blackish when mature.



2.5–7.5 cm
1–3 in

WEeping WIDOW

Lacrymaria lacrymabunda

F: Psathyrellaceae

Common on pathsides and disturbed soils in Eurasia and N. America, it is named after its black gills that weep droplets from their edges.



2.5–8 cm
1–3¼ in

COMMON INKCAP

Coprinopsis atramentaria

F: Psathyrellaceae

Occurring in Eurasia and N. America, this species has egg-shaped caps that dissolve to a blackish fluid as the spores are released.

MAGPIE INKCAP

Coprinopsis picacea

F: Psathyrellaceae

Found on chalky soils in Eurasian woodlands, this uncommon species has white, fluffy scales that contrast with the dark grey-brown cap.



white veil breaks into patches

5–8 cm
2–3¼ in