

» AGARICALES



4–10 cm
1½–4 in

DEER SHIELD
Pluteus cervinus
F: Pluteaceae

Found in Eurasia and N. America, this species is variable in colour, with a radially fibrous cap, often dome-centred; pink gills not attached to the stem; and a fibrous stem.



1–6 cm
¾–2¼ in

YELLOW SHIELD
Pluteus chrysophlebius
F: Pluteaceae

A golden to greenish yellow cap, yellow gills turning pink, and a whitish stem characterize this species. Found in Eurasia, it grows on decayed wood.



2.5–8 cm
1–3¼ in

WILLOW SHIELD
Pluteus salicinus
F: Pluteaceae

Common in broadleaved woodlands in Eurasia, this species can be identified by the bluish grey staining at the base of its slender stem.



10–25 cm
4–10 in

BEEFSTEAK FUNGUS
Fistulina hepatica
F: Schizophyllaceae
Resembling a fleshy steak, with blood-like red juice, this species is particularly common in warmer parts of Eurasia and N. America.



SILKY ROSEGILL
Volvariella bombycina
F: Pluteaceae

10–25 cm
4–10 in

Growing on deciduous trees in Eurasia and N. America, this rare species has a white to pale lemon cap and a whitish stem with a bag-like thin veil enclosing the base of the stem.



CAP (UNDERSIDE)

COMMON PORECRUST
Schizophyllum commune
F: Schizophyllaceae
Found in Eurasia and N. America, this fan-shaped fungus is identified by the gill-like spore surface on the underside of the cap.



CAP (UPPER SIDE)



6–14 cm
2¼–5½ in

STUBBLE ROSEGILL
Volvopluteus gloiocephalus
F: Pluteaceae

Quite common in fields and old stubble, and on woodchip mulch, in Eurasia and N. America, this species has a sticky, grey cap.



3–7 cm
1¼–2¾ in

CONIFER TUFT
Hypholoma capnoides
F: Strophariaceae


Found in Eurasia and N. America, this uncommon species occurs on conifer wood. Its whitish gills mature to greyish lilac.



3–7 cm
1¼–2¾ in

SULPHUR TUFT
Hypholoma fasciculare
F: Strophariaceae

Abundant in temperate woods in Eurasia and N. America, this species has greenish yellow gills that mature to dark purple, and are easy to recognize in the field.



5–10 cm
2–4 in

BRICK TUFT
Hypholoma lateritium
F: Strophariaceae

This species is characterized by a fleshy cap with veil fragments and pale yellow gills that mature to lavender. It grows on hardwoods in Eurasia and N. America.