

**SHEATHED WOODTUFT**

Kuehneromyces mutabilis
F: Strophariaceae

Often confused with the deadly *Galerina marginata*, this species can be identified by its sticky cap, scaly stem, and brown gills. It occurs in Eurasia and N. America.

2–7 cm
¾–2¾ in

ring on stem

**REDLEAD ROUNDHEAD**

Leratiomyces ceras
F: Strophariaceae

Previously known as *Stropharia aurantiaca*, this species has a bright red cap and reddish flushed stem. It grows among woodchips in Eurasia and N. America.


1.5–6 cm
½–2¼ in

**ALDER SCALYCAP**

Pholiota alnicola
F: Strophariaceae

Found in Eurasia, this species is characterized by a sticky cap and tufted habit. Despite its common name, it is usually seen at the base of birches.

3–7 cm
1¼–2¼ in



3–12 cm
1¼–4¾ in

GOLDEN SCALYCAP

Pholiota aurivella
F: Strophariaceae

This rather common species is found on beech logs or trunks in Eurasia and N. America. Its sticky golden caps have dark orange-brown scales.



0.5–4 cm
¼–1½ in

DUNG ROUNDHEAD

Protostropharia semiglobata
F: Strophariaceae

Occurring on animal dung or in grass where animals graze, this species is found in Eurasia and N. America. It has a sticky cap and a slender stem with a ring.



5–15 cm
2–6 in

SHAGGY SCALYCAP

Pholiota squarrosa
F: Strophariaceae

Growing in Eurasia and N. America, this fungus has a distinctive dry, sharply scaly cap and stem; pale yellow gills; and a smell of corn or radish.



0.5–2 cm
¼–¾ in

LIBERTY CAP

Psilocybe semilanceata
F: Strophariaceae

This well-known agaric has a conical, dull yellow cap with a distinct point. Common in meadows in late autumn, it occurs in Eurasia and N. America.



3–7 cm
1¼–2¾ in

BLUE-GREEN SLIMEHEAD

Stropharia cyanea
F: Strophariaceae

This fungus has a blue-green cap that fades to yellow. It is found in Eurasia and N. America.



6–15 cm
2¼–6 in

WHITELACED SHANK

Megacollybia platyphylla
F: Incertae sedis

Found in Eurasia and N. America, this species has a radially fibrous, pale grey-brown cap; widely spaced, deep gills; and a stem base with root-like strands.

scaly, furry surface



4–7 cm
1½–2¾ in

COMMON CAVALIER

Melanoleuca polioleuca
F: Incertae sedis

Common in grass in Eurasia, this species has a grey-brown cap with white gills. Its stem has blackish flesh at the base.




5–10 cm
2–4 in

PLUMS AND CUSTARD

Tricholomopsis rutilans
F: Incertae sedis

Common on pine stumps, this species has a reddish purple cap and stem, and golden-yellow gills. It grows in Eurasia and N. America.



5–15 cm
2–6 in

SPECTACULAR RUSTGILL

Gymnopilus junonius
F: Incertae sedis

Occurring in Eurasia, this fungus is found in clumps, usually at the base of a tree. It has a dry cap with crowded, shallow yellowish gills.

1–6 cm
¾–2¼ in



1–4 cm
¾–1½ in

PETTICOAT MOTTLEGILL

Panaeolus papilionaceus
F: Incertae sedis

The minute toothed remains of the veil at the cap margin and mottled black gills characterize this species. It is found in Eurasia and N. America.



3–7 cm
1¼–2¾ in

GOBLET

Pseudoclitocybe cyathiformis
F: Incertae sedis

This distinctive species has a very dark coloration and a tall, fibrous stem. Found in Eurasia, it is common in late autumn and winter.



5–20 cm
2–8 in

TROOPING FUNNEL

Infundibulicybe geotropa
F: Incertae sedis

This fungus has a funnel-shaped, fleshy cap; a tall stem; and pale leather-brown colouring. It occurs in Eurasia.

EGGHEAD MOTTLEGILL

Panaeolus semiovatus
F: Incertae sedis

Found on animal dung, this species occurs in Eurasia and N. America. It is characterized by a sticky grey cap and a ring around its tall stem.

CINNABAR POWDERCAP

Cystoderma cinnabarina
F: Incertae sedis

Found in Eurasia and N. America, this species has a brick-red cap and pale cream gills. The cap and stem have a granular surface.

root attaches to conifer wood